

Effect of cobalt on synthesis of extracellular alkaline phosphatase production from *Bacillus* sp.

Abstract

Various species of *Bacillus* isolated from soil were characterized and tested for their ability to produce extracellular alkaline phosphatase. Six strains of *Bacillus* demonstrated significant activity as compared to *Bacillus subtilis* 6633 in optimized *Bacillus* medium ATCC 552. Highest levels of extracellular alkaline phosphatase activity were observed with 1% whey, in the presence 0.1mM CoCl₂ (10.4 U/ml/min). This trend was observed in five of the six strains tested.

Keywords: alkaline phosphatase, *Bacillus*

Volume 2 Issue 6 - 2015

Dipak Vora, Kainaz Irani

Department of Microbiology, Ramnarain Ruia College, Mumbai University, India

Correspondence: Dipak Vora, Department of Microbiology, Ramnarain Ruia College, Mumbai University, India, Email knazirani@gmail.com

Received: November 04, 2014 | **Published:** December 02, 2015

Introduction

Alkaline phosphatase, APase (EC 3.1.3.1), a metalloenzyme, catalyses the non-specific hydrolysis of phosphate monoesters¹ and has been extensively studied in various organisms ranging from bacteria to mammals.²⁻⁴ In the bacteria alkaline phosphatase is found in the periplasmic space⁵ and is induced under low phosphate concentrations suggesting its involvement in phosphate metabolism.⁶ Phosphatases are thought to function as phosphate scavengers under condition of phosphate depletion. Further, addition of stimulants such as detergents, metal-salts, and various other compounds enhance the activity of the enzyme.^{7,8} Very few studies have been carried out on extracellular phosphatases.⁹ This investigation aims at determining the role of metal ions in the production of extracellular alkaline phosphatase in *Bacillus* species. The morphological properties and taxonomic characteristics of the organisms isolated from soil were studied.¹⁰ and biochemical analysis was processed by probabilistic identification software, PIBWIN.¹¹ Six strains of *bacillus* screened for phosphatases showed much higher activity as compared to *Bacillus subtilis* 6633.¹²

Bacillus medium ATCC 552 having the following composition was used (g/l): Peptone, 10g; Glucose, 10%; Beef extract, 3g, NaCl, 5g; K₂HPO₄, 5g; K₂HPO₄, pH adjusted to 7.6 and sterilized by autoclaving. The cultures were grown in 100ml of medium in 250ml Erlenmeyer flasks after inoculation with 5ml of overnight culture and this was incubated at ambient temperature on a rotary shaker having a throw of 2.5-3.81cms at 200rpm. Aliquots (5ml) were withdrawn at

4h intervals and centrifuged at 3000xg for 10 minute. The supernatant obtained was assayed for the enzyme.

Alkaline phosphatase activity was assayed as described by Sawhney & Singh,¹³ taking into consideration the optimum pH and temperature of the *bacillus* cultures. One unit of alkaline phosphatase activity was defined as the amount of enzyme that liberates 1μmol of p-nitrophenol/min under defined conditions. All reagents were obtained from Loba Chemie Pvt Ltd.

The medium was supplemented with (1% w/v) carbon sources such as lactose, maltose, malt extract, starch, sucrose or whey. A control flask of *Bacillus* medium ATCC 552+1% glucose was also maintained. Whey sample was prepared from Cow's milk (obtained locally) and the lactose content determined.¹⁴ The final medium contained whey ≈1% lactose. Total enzyme activity (U) was analyzed at different time intervals during incubation.

Further the isolates were grown in medium prepared in double distilled water and supplemented with metal salts (0.1mM chloride form) such as Ba, Ca, Co, Cu, Fe, K, Li, Mg, Mn, Ni, Zn and extracellular alkaline phosphatase activity determined.

The APase activity varied among the *Bacillus* sp. and maximum activity was obtained with the medium supplemented with 1% whey as carbon source (Table I). Further, addition of 0.1mM CoCl₂ to the production medium enhanced extracellular alkaline phosphatase maximally in all the *bacillus* species studied except, *Taxon 18* (Table 2). Higher concentration of metal ions suppressed APase activity right across upto 24h.

Table I Effect of Carbon sources on alkaline phosphatase production in *Bacillus* sp

Carbon Source (1%)	<i>B. badius</i> (U/ml)	<i>B. cereus</i> (U/ml)	<i>B. firmus</i> (U/ml)	<i>B. licheniformis</i> (U/ml)	<i>B. smithii</i> (U/ml)	<i>Taxon</i> (U/ml)
Glucose*	71.2	46	69.3	32.1	25	16.9
Lactose	14.9	18.8	14.5	40	15.3	5.2
Maltose	44.4	42	63.4	31.7	51.2	45
Malt extract	7.8	2.4	1.4	7.8	10.4	6.5

Table continued

Carbon Source (1%)	<i>B. badius</i> (U/ml)	<i>B. cereus</i> (U/ml)	<i>B. firmus</i> (U/ml)	<i>B. licheniformis</i> (U/ml)	<i>B. smithii</i> (U/ml)	Taxon (U/ml)
Starch	3.1	3.1	6.9	6.2	5.1	1.1
Sucrose	7.3	2.9	6.4	6.3	6.6	3.8
Whey	107.9	111.1	87.2	91.6	184	161

*Control.

Table 2 Alkaline phosphatase activity in presence of metal ions. Time of incubation (h)

Organism	Metal ions (0.1 mM)	Activity (U/ml)
<i>Bacillus badius</i>	CoCl ₂	252 (16h)
<i>Bacillus cereus</i>	CoCl ₂	126 (16h)
<i>Bacillus firmus</i>	CoCl ₂	157 (16h)
<i>Bacillus licheniformis</i>	CoCl ₂	205 (12h)
<i>Bacillus smithii</i>	CoCl ₂	146 (16h)
Taxon 18	MnCl ₂	108 (20h)

Our results suggest that the stimulation after addition of cobalt on APase production in *Bacillus* is not restricted to *Bacillus licheniformis*¹⁵ indicating that the regulatory mechanism among all the *Bacillus* species may be similar.

Acknowledgments

None.

Conflicts of interest

Authors declare that there is no conflict of interest.

References

- Coleman J. Characterization of the *Escherichia coli* gene for 1-acyl-sn-glycerol-3-phosphate acyltransferase (plsC). *Mol Gen Genet.* 1992;232(2):295–303.
- Junior AB, GuimarAes LH, Terenzi HF, et al. Purification and biochemical characterization of thermostable alkaline phosphatases produced by *Rhizopus microsporus* var. *rhizopodiformis*. *Folia Microbiol (Praha)*. 2008;53(6):509–516.
- Simão AMI, Beloti MM, Rosa AL, et al. Culture of osteogenic cells from human alveolar bone: A useful source of alkaline phosphatase. *Cell Biol Int.* 2007;31(11):1405–1413.
- Sugahara T, Konno Y, Ohta H, et al. Purification and properties of two membrane alkaline phosphatases from *Bacillus subtilis* 168. *J Bacteriol.* 1991;173(5):1824–1826.
- Campbell LL, Hulett-cowling FM. Purification and properties of an alkaline phosphate of *Bacillus licheniformis*. *J Biochem.* 1971;10:1364–1371.
- Hulett FM. The signal-transduction network for Pho regulation in *Bacillus subtilis*. *Mol Microbiol.* 1996;19(5):933–939.
- Sharipova MR, Balaban NP, Nekhotyaeva NV, et al. A novel *Bacillus intermedius* extracellular alkaline phosphatase: isolation, physico-chemical and catalytic characteristics. *Biochem Mol Biol Int.* 1996;38(4):753–761.
- Oh WS, Im YS, Yeon KY, et al. Phosphate and carbon source regulation of alkaline phosphatase and phospholipase in *Vibrio vulnificus*. *J Microbiol.* 2007;45(4):311–317.
- Sharipova MR, Balaban NP, Mardanova AM, et al. Isolation and properties of extracellular alkaline phosphatase from *Bacillus intermedius*. *Biochem (Mosc)*. 1998;63(10):1178–1182.
- Buchanan RE, Gibbons NE. *Bergey's manual of determinative bacteriology*. Williams and Wilkins: Baltimore, USA; 1974. 1246 p.
- Bryant TN. PIBWIN- Software probabilistic identification. *J Appl Microbiol.* 2004;97(6):1326–1327.
- Barber M, Kuper SW. Identification of *Staphylococcus pyogenes* by the phosphatase reaction. *J Pathol Bacteriol.* 1951;63(1):65–68.
- Sawhney SK, Singh R. *Introductory Practical Biochemistry*. Norosa Publishing House: New Delhi, India; 2002.
- Teles FFF, Young CK, Stull JW. A method for rapid determination of lactose. *J Dairy Sci.* 1978;61(4):506–508.
- Spencer DB, Chen CP, Hulett FM. Effect of cobalt on synthesis and activation of *Bacillus licheniformis* phosphatase. *J Bacteriol.* 1981;145(2):926–933.