

Table 7 Antimicrobial susceptibility pattern of bacterial isolates

Isolates	ANTIBIOTICS																
	OFX	PEF	CPX	AU	CN	S	CEP	NA	SXT	PN	NB	RD	E	APX	LEV	AML	CH
	S (%)	S (%)	S (%)	S (%)	S (%)	S (%)	S (%)	S (%)	S (%)	S (%)	S (%)	S (%)	S (%)	S (%)	S (%)	S (%)	S (%)
<i>P. mirabilis</i> (17)	10(58.8)	8(47.1)	13(76.5)	11(64.7)	17(100)	17(100)	17(100)	16(94.0)	16(94.0)	17(100)	NA	NA	NA	NA	NA	NA	NA
<i>E. coli</i> (22)	18(81.8)	11(50.0)	16(72.7)	10(45.5)	14(63.6)	20(90.9)	13(59.1)	7(31.8)	10(45.5)	9(40.9)	NA	NA	NA	NA	NA	NA	NA
<i>P. aeruginosa</i> (8)	8(100)	7(87.5)	8(100)	4(50.0)	6(75.0)	8(100)	4(50.0)	2(9.1)	6(75.0)	4(50.0)	NA	NA	NA	NA	NA	NA	NA
<i>K. pneumonia</i> (10)	7(70.0)	8(80.0)	8(80.0)	3(30.0)	8(80.0)	9(90.0)	7(70.0)	6(60.0)	7(70.0)	6(60.0)	NA	NA	NA	NA	NA	NA	NA
<i>S. aureus</i> (24)	NA	NA	17(70.8)	NA	23(95.8)	22(92.0)	NA	NA	NA	NA	13(54.2)	0	3(12.5)	12(50.0)	3(12.5)	11(45.8)	5(20.8)
Total (81)	43(53.1)	34(42.0)	62(76.5)	28(34.6)	68(84.0)	76(94.0)	41(50.6)	31(38.3)	39(48.1)	36(44.4)	13(16.0)	0	3(3.7)	12(14.8)	3(3.7)	11(13.6)	5(6.2)

Keywords: OFX, Tarivid; PEF, Reflacine; CPX, Ciprofloxacin; AU, Augmentin; CN, Gentamycin; S, Streptomycin; CEP, Ceporex; NA, Nalidixic acid; SXT, Septrin; PN, Ampicillin; NB, Norfloxacin; RD, Rifampicin; E, Erythromycin; APX, Ampiclox; LEV, Levofloxacin; AML, Amoxicillin; CH, Chloramphenicol; NA, not applicable; S (%), sensitivity in percentage