

FMRI analysis of the motor network functional connections at rest and with motor load in healthy people and patients with STBI

Introduction

Severe traumatic brain injury (STBI) is accompanied by motor disorders in 2/3 of the victims - often in combination with cognitive impairment and depression of consciousness.¹ An objective diagnosis of the motor sphere in patients with SBI at different severity of motor disorders is among the actual ones.

Work objective

To investigate the possibility of the motor corticospinal tract network's functional activity in normal and STBI estimating based on a comparison of fMRI connectivity in the motor act and at rest.

Method

Observation groups: 18 patients with STBI (main) and 15 healthy volunteers (control). Movement defect in the form of hemiparesis were evaluated on a scale of muscle strength.² 3T fMRI recorded at rest and passive right hand finger clenching (by experimenter). The validity of this motor test using is shown in our previous publications.^{3,4} The CONN program carried out a group analysis of the cortico-spinal tract's motor network brain connectivity (BC) - between the regions of interest (ROI) in the motor cortex of the left (1) and right (2) hemispheres, supplementary motor cortex (3), and cerebellum right hemisphere (4).

Three of these brain regions are involved in the fMRI response, to both the active and the passive movement of the arm (Figure 1).^{3,4} The intensity of BC between the selected points was assessed, as well as the level of their significance.




Figure 1 Averaged fMRI responses of healthy right-handed at the active and passive movement of the right hand.³ NI7, $t=3.69$, $p<0.001$. Digits indicate the selected ROI.

Results

In Figure 2 shows histograms of the BC intensity distribution in predetermined motor fMRI network of healthy people and patients with various degrees of right-sided hemiparesis after STBI in passive motor test (Figure 2A) and at rest (Figure 2B). Both the features of similarity and the differences in their spatial organization are noted. In both tests close order of the BC intensity values in the majority of ROI pairs of the same name was observed, as well as the tendency to weaken them from normal to rough hemiparesis.

At rest these changes are practically linear in most pairs analyzed (4 of 6). Less evident this orientation for the connections of the left and right motor cortex with the right hemisphere of the cerebellum. In the passive motor test for only one connection from 6 (between left

Volume 8 Issue 2 - 2018

EV Sharova,¹ TS Mukhina,^{1,3} GN Boldyreva,¹ LA Zhavoronkova,¹ MV Chelyapina,¹ AS Smirnov,² EV Alexandrova,² IN Pronin²

¹Institute of Higher Nervous Activity and Neurophysiology of RAS, Russia

²NN Burdenko National Medical Research Center of Neurosurgery of the Ministry of Health of the Russian Federation, Russia

³National Research Nuclear University MEPhI (Moscow Engineering Physics Institute), Russia

Correspondence: EV Sharova, Institute of Higher Nervous Activity and Neurophysiology of RAS, Russia, Email ESharova@nsi.ru

Received: February 13, 2018 | **Published:** March 16, 2018

and right motor cortex, 1-2) values decrease linearly with increasing hemiparesis. BC of the supplementary motor cortex with right motor, as well as with cerebellum, are the highest in easy hemiparesis in comparison with other samples of observations, and the connections of the right and left motor cortex with the cerebellum, on the contrary, are weakened. The nonlinearity of these changes correlates with the previously identified topographic features of motor fMRI responses at different degrees of posttraumatic hemiparesis,⁵ reflecting, in our opinion, the variational mechanisms of motor defect compensation.




Figure 2 Histograms of the fMRI motor network's BC intensity in healthy people and patients with post-traumatic hemiparesis. A - a passive motor test, B - resting state, I - norm (N=15), II - easy hemiparesis (N=8), III - rough hemiparesis (N=10). Horizontal axis - the ROI couples. Vertical axis - BC intensity. BC significance level by one-sample t-test: * 0,05<math>p<0,15</math>; ** 0,01<math>p<0,05</math>; ***

Conclusion

Both in rest and in a passive motor sample, changes of the BC between the symmetrical motor cortical zones are associated with the severity of posttraumatic hemiparesis most clearly. Evaluation of the motor network fMRI paretic arm at rest has a greater diagnostic potential. In the passive movement of the hand intergroup changes of the motor network BC spatial organization as hemiparesis increases can more reflect the compensatory possibilities of the studied system.

Disclosure

Supported by RFFI № 18-013-00355

Conflict of interest

The authors declare there is no conflict of interest.

Acknowledgements

None.

References

1. Potapov AA, Likhтерman LB, Kravchuk AD, et al. Traumatic brain injury: problems and prospects. *Voprosy neurokhirurgii im.N.N.Burdenko*. 2009;(2):3–8.
2. McPeak LA. Physiatrie history and examination. In: Braddom R, editor. *Physical medicine and rehabilitation*. Philadelphia: WB Saunders Company; 1996. p. 3–42.
3. Boldyreva GN, Sharova EV, Zhavoronkova L, et al. fMRI and EEG response of the brain of a healthy person with active and passive movements of the leading hand. *Zhurnal vysshei nervnoi deyatel'nosti im. I.P. Pavlova*. 2014;64(5):488–99.
4. Boldyreva GN, Sharova EV, Zhavoronkova LA, et al. The fMRI mapping of brain responses of healthy people with active, passive and imagined movements of the hand. *Meditinskaya vizualizatsiya*. 2015;(5):100–12.
5. Sharova EV, Boldyreva GN, Zhavoronkova LA, et al. FMRI analysis of the human brain's neuroplasticity as a basis of movement disorders compensation after traumatic brain injury. *Clinical Neurophysiology*. 2017;128(9):e278–e279.