

# Painless, Reproducible, Reversible Floor of Mouth Herniation: What's in There?

## Opinion

### Description

An 18 year old female presented with a left submandibular swelling (Figure 1) which was diagnosed to be a plunging ranula and surgery was recommended. The patient moved abroad and as the neck lump disappeared spontaneously no active intervention was ever sought. Ten years later, she represented with a right submandibular swelling which fluctuated in size over the preceding 4 months but had failed to disappear completely. The clinical examination revealed a soft, cystic swelling in the right submandibular region (Figure 2). The clinical diagnosis of right plunging ranula was made with no intra-oral component (Figure 3). The Ultrasound scan revealed a complex cyst (measuring 6x4x2cm) sitting in front of a normal right submandibular gland. Two weeks before her planned surgery, she developed acute infection involving her right floor of mouth with significant neck swelling. She responded well to intravenous antibiotics and was discharged home the following day on oral antibiotics.

On the day of surgery, no residual swelling could be seen in her neck at rest (Figure 4). However, the patient volunteered to reproduce her neck swelling (Video 1). To our surprise, there were bilateral submandibular swellings which became apparent on modified valsalva manoeuvre and spontaneously reduced without causing any discomfort to the patient. The patient underwent transoral excision of the right sublingual salivary gland and drainage of the associated cyst under general anaesthetic. The patient made excellent recovery and was discharged home the same day.

Plunging ranula is a pseudocyst formed by extravasation of mucous from the sublingual gland (SLG) into the submandibular space. It is thought to have a congenital aetiology [1]. The plunging ranula is suspected clinically and confirmed; radiologically by Ultrasound [2], CT or MRI scan, and by fine needle aspiration for cytology or biochemical analysis confirming presence of salivary amylase in the aspirate. Based on the published literature, transoral excision of the SLG is favoured to treat the plunging ranula [3].

### Learning points

- Bilateral plunging ranula is a rare phenomenon but is an important differential diagnosis for submandibular region swellings.
- Bilateral reversible sublingual gland herniation through the mylohyoid muscles defect without associated ranula does not always need removal of gland.
- Transoral excision of the sublingual salivary gland with drainage of the cystic contents is the recommended treatment for plunging ranula.

## Opinion

Volume 4 Issue 6 - 2016

**Shakeel M, Nair S and Z Ahmad\***

*Department of Otolaryngology-Head and Neck Surgery, Super Clinic, New Zealand*

**\*Corresponding author:** Zahoor Ahmad, Consultant Otolaryngologist, Department of Otolaryngology- Head and Neck Surgery, Manukau Super Clinic, PO Box 98743, South Auckland Mail Center, Auckland 2241, New Zealand, Tel: +64212137576; Email: Zahoor.Ahmad@middlemore.co.nz

**Received:** August 28, 2016 | **Published:** October 14, 2016




Figure 1: Left submandibular swelling-Left plunging ranula.




Figure 2: Right submandibular swelling-Right plunging ranula.



**Figure 3:** Normal floor of mouth.



**Figure 4:** No visible submandibular regions swelling.



**Figure 5:** Bilateral submandibular swellings secondary to herniation of the sublingual glands.



Please Click here on above video

### References

1. Harrison JD, Kim A, Al-Ali S, Morton RP (2013) Postmortem investigation of mylohyoid hiatus and hernia: aetiological factors of plunging ranula. *Clin Anat* 26(6): 693-699.
2. Jain P, Jain R, Morton RP, Ahmad Z (2010) Plunging ranulas: high-resolution ultrasound for diagnosis and surgical management. *Eur Radiol* 20(6): 1442-1449.
3. Samant S, Morton RP, Ahmad Z (2011) Surgery for plunging ranula: the lesson not yet learned? *Eur Arch Otorhinolaryngol* 268(10): 1513-1518.