

Case Study: How the Use and Abuse of Cannabis is Related with Organic Brain Damage and Psychotic Symptoms

Case Report

Volume 3 Issue 6 - 2015

Victoria Dolia*

University of crete, Spain

***Corresponding author:** Victoria Dolia, University of crete, Carrer pal.las, 12, les rotes, Denia, Spain, Email: psicologa.viktoriadoliala@outlook.com

Received: March 09, 2015 | **Published:** October 15, 2015

Abstract

Background: Many studies have been done in order to show that the use of Cannabis can cause psychotic symptoms and brain damage. The aim of this study is to investigate closely what symptoms are presented in chronic users of cannabis. Also, demonstrate with a clinical investigation that chronic use of cannabis (and withdrawal syndrome too) can cause organic brain damage and psychotic symptoms.

Case presentation: Mr. K is a 31 years old man. He is from Nigeria. He graduated from college at Nigeria. Since 2009 he lives in Greece. He has a story of abuse of cannabis lasting 11 years, since his early 20s. He smokes cannabis daily, two (2) "trifoliolate" cigarettes without adding tobacco with nicotine. The two (2) "trifoliolate" cigarettes correspond to one (1 gr) gram of cannabis approximately.

Methodology and results: During the interview his behavior did not give any information relating to psychiatric/psychological disorder or cerebral dysfunction. He answered openly and honestly all the questions. Through the appropriate questions for detecting clinical symptoms of psychopathology were observed some elements of psychological dysfunction.

Also, he performed a set of neuropsychological tests to investigate, if he presents cerebral dysfunction. The tests showed that there were some elements of organic brain damage. With the help of the interview and tests, that are valid and reliable tools, we were able to identify and find the presence of psychotic symptoms and panic attacks which were directly related to the use of cannabis, as these occur under the influence of cannabis. Moreover, the chronic use of cannabis has caused him brain dysfunction.

Methodology

For this study clinical evaluation within the interview were used, consisting of both closed and open questions [1-4]. Also valid and reliable clinical neuropsychological tests were used, precisely including Bender Visual Motor Gestalt Test and Rey Osterrieth Complex Figure Test.

Clinical evaluation

From the clinical examination was found that Mr. K. presented many positive psychotic symptoms and panic attacks. Also, his mood was characterized as depressed throughout the duration of the interview.

On one hand, when he was under the influence of cannabis, he had delusions and perceptual disorders. He stated that others wanted to harm him (Delusions of Reference and Persecutory Delusions), he also stated that he was capable of [] s ic "speaking with. God, but sometimes also with the Devil" (Religious Delusions) and believed that he was a very important person (Grandiose delusions). Moreover, he believed that he could read the thoughts of others and that others could read his thoughts too (Delusions of mind being read). He had presented Delusion of thought broadcasting, thought insertion and Delusions of catastrophe [5-8].

On the other hand, we observed perceptual disorders. Specifically he stated Auditory Hallucinations. For example, he heard [] s ic "someone shouting his name through the. Closet, listening to curse him and say that "he smokes too much, and will affect the brain". These voices were reported in the third person singular, characteristic of auditory hallucinations. Also, he reported visual hallucinations, e.g. "he sees beautiful women".

The examination of his mental condition has shown that he presents memory lapses and significant impairment of concentration, attention and deductive reasoning.

Neuropsychological assessment

We used the valid and reliable clinical neuropsychological tests Bender Visual Motor Gestalt Test και Rey-Osterrieth Complex Figure Test, in order to investigate if he had brain damage and disturbed cognitive functions [9,10].

Rey-Osterrieth complex figure test: The Rey-Osterrieth Complex Figure Test (ROCF a neuropsychological assessment in) is which examinees are asked to reproduce a complicated line drawing, first by copying it freehand (recognition), and then drawing from memory (recall) [11-14]. This test permits the evaluation of different functions, such as visuospatial abilities, memory, attention, planning and working memory (executive functions).

Three conditions are most commonly used in the ROCF.

i. Copy: In the Copy condition, the examiner is given a piece of paper and a pencil, and the stimulus figure is placed in front of them. They reproduce the figure to the best of their ability. The test is not timed, but the length of time needed to copy the figure is observed. Some administrators use a series of colored pencils, in order to preserve a record of the order in which design elements were reproduced [15]. However, because of concerns that the use of color changes the nature of the test and makes it easier for the subject to remember the figure, the current test manual suggests that this should not be done. Instead, the evaluator should take notes on the process the examinee uses [16]. Once the copy is complete, the stimulus figure and the examinee's copy are removed from view.

ii. Immediate recall: After a short delay, the examinee is asked to reproduce the figure from memory.

iii. Delayed recall: After a longer delay (20–30 minutes), the examinee may again be asked to draw the figure from memory. Examinees are not told beforehand that they will be asked to draw the figure from memory; The Immediate and Delayed Recall conditions are therefore tests of incidental memory [17]. Each copy is scored for the accurate reproduction and placement of 18 specific design elements. Additionally, the test administrator can note their qualitative observations regarding the examinee's approach to the task and the effectiveness of any apparent strategy use.

Results

I. Copy: Rates: 28 points (being excellent a score of 36 points).

The average score on this test is 32 points. The score gathered by the examinee copying the project (28/36) was below the average for the total population, indicating that it was affected the perceptual functions and the ability of the examinee to perform simple tasks, as represented in this case by copying a drawing.

II. Immediate recall: m

Rates: 4 (being excellent a score of 36 points)

The reproduction of a figure by memory was imperfect. Score gathered by the examinee (4/36) has a very important indication that there is some neuropsychological dysfunction, which affects the visual memory of the examinee.

III. Delayed recall: We didn't proceed in this task due to the results of immediate recall.

Bender visual motor gestalt test: The Bender Gestalt Test, or the Bender Visual Motor Gestalt Test, is a psychological assessment instrument used to evaluate visual motor functioning and visual perception skills in both children and adults [18]. Scores on the test are used to identify possible organic brain damage and the degree maturation of the nervous system [19,20].

The Bender Gestalt Test is an individually administered pencil and paper test used to make a diagnosis of brain injury [21-24]. There are nine geometric figures drawn in black. These figures are presented to the examinee one at a time; Then, the examinee is asked to copy the figure on a blank sheet of paper [25,26]. Examinees are allowed to erase, but cannot use any mechanical aids (such as rulers). The popularity of this test among clinicians

is most likely the short amount of time it takes to administer and score. The average amount of time to complete the test is five to ten minutes [27].

The Bender Gestalt Test lends itself to several variations in administration. One method requires that the examinee view each card for five seconds, after which the card is removed [28-31]. The examinee draws the figure from memory. Another variation involves having the examinee draw the figures by following the standard procedure. The examinee is then given a clean sheet of paper and asked to draw as many figures as he or she can recall [32].

Last, the test is given to a group, rather than to an individual (i.e., standard administration). It should be noted that these variations were not part of the original test.

Results

Due to neuropsychological dysfunction of the brain which affect the cognitive functions of the examinee. The results of both ReyOsterrieth Complex Figure Test and Bender Visual Motor Gestalt Test indicated some elements of organic brain damage, which affect his cognitive functions [33-36].

Conclusion

The case study helps to investigate and observe symptoms and signs in which we want to focus on. The neuropsychological tests are tools that help us to get additional information about the state of cognitive functions of the individual and any indication that there is some neuropsychological organic brain dysfunction.

So we observed that in our case patient, the systematic use of cannabis led not only to the appearance of psychotic symptoms, but also was linked to brain dysfunction. Furthermore, were detected perceptual disorders and visual constructive disability? We observed significant problems in short-term and long-term memory of the examinee, characterized mainly by memory lapses and difficulty to recall information that was related with visual memory. Apart from the difficulty that the person had to remember the figure that was copied previously, he had considerable difficulty in reproducing the figures. Reflect this significant evidence relating to organic brain dysfunction.

The psychotic symptoms occurred during the use of cannabis, and deprivation of it. He presented psychotic symptoms triggered at withdrawal. The withdrawal symptoms, he presented included panic attacks. e can hypothesize that when the patient. Experienced W anxiety and excessive stress under withdrawal syndrome of cannabis, it displayed psychotic symptoms, mainly positive psychotic symptoms.

References

1. Arseneaul L, Cannon M, Witton J, Murray RM (2004) Causal association between cannabis and psychosis: examination of the evidence. *Br J Psychiatry* 184: 110-117.
2. Ashtari M, Avants B, Cyckowski L, Cervellione KL, Roofeh D, et al. (2011) Medial temporal structures and memory functions in adolescents with heavy cannabis use. *J Psychiatr Res* 45(8): 1055-1066.
3. Becker B, Wagner D, GouzoulisMayfrank E, Spuentrup E, Daumann J (2010) The impact of early onset cannabis use on functional brain correlates of working memory. *Prog NeuroPsychopharmacology Biol Psychiatry* 34(6): 837-845.

4. Bally N, Zullino D, JeanMichel Aubry (2014) Cannabis use and first manic episode. *Journal of Affective Disorders* 165(20): 103108.
5. Campbell AM, Evans JL, Thomson GM, Williams MJ (1971) Cerebral atrophy in young cannabis smokers. *Lancet* 2(7736): 12191224.
6. Cassidy CM, Lepage M, Harvey PO, Malla A (2012) Cannabis use and anticipatory pressure as reported by subjects with early psychosis and community controls. *Schizophrenia Research* 137(1-3): 3944.
7. Corcoran CM, Kimhy D, Stanford A, Khan S, Walsh J, et al. (2008) Temporal association of cannabis use with symptoms in individuals at clinical high risk for psychosis. *Schizophrenia Research* 106(2-3): 286293.
8. Cunha PJ, Rosa PG, Ayres Ade M, Duran FL, Santos LC, et al. (2013) Cannabis use, cognition and brain structure in first episode psychosis. *Schizophrenia Research* 147(2-3): 209215.
9. Davis GP, Compton MT, Wang S, Levin FR, Blanco C (2013) Association between cannabis use, psychosis and schizotypal personality disorder: Findings from the National Epidemiologic Survey on Alcohol and related conditions. *Schizophrenia Research* 151(1-3): 197202.
10. Degenhardt L, Hall W (2002) Cannabis and psychosis. *Current Psychiatry Reports* 4: 191196.
11. Duley JF, Wilkins J, Hamby S, Hopkins D, Burwell R, et al. (1993) Explicit scoring criteria for the ReyOsterrieth and Taylor complex figures. *Clinical Neuropsychologist* 7(1): 29-38.
12. Hides L, Lubman DI, Buckby J, Yuen HP, Cosgrave E, et al. (2009) The association between early cannabis use and psychoticlike experiences in a community adolescent sample. *Schizophrenia Research* 112(1-3): 130135.
13. Hubley AM, Jassal S (2006). Comparability of the ReyOsterrieth and Modified Taylor Complex Figures using total scores, completion times, and construct validation. *Clin Exp Neuropsychol* 28(8): 1482-1497.
14. Hutt ML (1985) The Hutt Adaptation of the Bender Gestalt Test: Diagnostic and Therapeutic Implications. Grune and Stratton, New York.
15. Jager G, Block RI, Luijten M, Ramsey NF (2010) Cannabis use and memory brain function in adolescent boys: a crosssectional multicenter functional magnetic resonance imaging study. *Journal of the American Academy of Child & Adolescent Psychiatry* 49(6): 561572 (e1-e3).
16. Kalant H (2014) Chapter thirteen effects of cannabis and cannabinoids in the human nervous system. In: Bertha Madras & Michael KI (Eds.), *The Effects of Drug Abuse on the Human Nervous System*. Academic Press, pp. 387422.
17. Kaufman AS, Lichtenberger EO (2001) *Assessing Adolescent and Adult Intelligence*. Allyn and Bacon, USA.
18. Koppitz EM (1975) *The Bender Gestalt Test for Young Children*. Grune and Stratton, US.
19. Leweke FM, Gerth CW, Klosterkötter J (2012) CannabisAssociated Psychosis. *CNS Drugs* 18(13): 895910.
20. Lu PH, Boone KB, Cozolino L, Mitchell C (2003) Effectiveness of the ReyOsterrieth Complex Figure Test and the Meyers and Meyers recognition trial in the detection of suspect effort. *Clin Neuropsychol* 17(3): 426-440.
21. Mata I, PerezIglesias R, RoizSantianez R, TordesillasGutierrez,D, Pazos, A, et al. (2010) Gyrfication brain abnormalities associated with adolescence and earlyadulthood cannabis use. *Brain Research* 1317: 297304.
22. McLaren JA, Silins E, Hutchinson D, Mattick RP, Wayne Hall W (2010) Assessing evidence for causallink between cannabis and psychosis: A review of cohort studies. *International Journal of Drug Policy* 21(1): 1019.
23. Meyers JE, Meyers KR (1995) *Rey complex figure test and recognition trial: Professional manual*. PAR, Inc.
24. Osterrieth PA (1944) Filetest de copie d'une figure complex: Contribution a l'etude de la perception et de la memoire [The test of copying a complex figure: A contribution to the study of perception and memory]". *Archives de Psychologie*: 286-356.
25. Parakh P, Basu D (2013) Cannabis and psychosis: Have we found the missing links? *Asian Journal of Psychiatry* 6(4): 281287.
26. Pascal GR, Suttell BJ (1951) *The Bender Gestalt Test: Quantification and Validation for Adults*. Grune and Stratton, US.
27. Pope HG Jr, Gruber AJ, YurgelunTodd D (1995) The residual neuropsychological effects of cannabis: the current status of research. *Drug and Alcohol Dependence* 38(1): 2534.
28. Radhakrishnan R, Sewell RA, Skosnik PD, Ranganathan M (2014) Chapter Fourteen Cannabis, Cannabinoids and the association with psychosis, In *The Effects of Drug Abuse on the Human Nervous System*. Bertha Madras and Michael Kuhar (Eds.), Academic Press, pp. 423474.
29. Bhattacharyya S, Atakan Z, Martin-Santos R, Crippa JA, Kambeitz J (2015) Impairment of inhibitory control processing related to acute psychotomimetic effects of cannabis. *European Neuropsychopharmacology*. 25(1): 2637.
30. Sattler JM (2002) Assessment of visualmotor perception and motor proficiency. In: Jerome M Sattler (Eds.), *Assessment of Children: Behavioral and Clinical Applications*. (4th edn), San Diego: Publisher, Us.
31. Semple DM, McIntosh MA, Lawrie SM (2005) Cannabis as a risk factor for psychosis: systematic review. *Psychopharmacol* 19(2): 187194.
32. Strauss E, Spreen O (1990) Comparison of the Rey and Taylor figures. *Arch Clin Neuropsychol* 5(4):417-420.
33. Shin MS, Park SY, Park SR, Seol SH, Kwon JS (2006) Clinical and empirical applications of the ReyOsterrieth Complex Figure Test. *Nat Protoc* 1(2): 892-899.
34. van Tricht MJ, Harmsen EC, Koelman JH, Bour LJ, van Amelsvoort TA, et al. (2013) Effects of cannabis use on event related potentials in subjects at ultra high for psychosis and healthy controls. *International Journal of Psychophysiology* 88(2): 149156.
35. Varma VK, Malhotra AK, Dang R, Das K, Nehra R (1988) Cannabis and cognitive functions: A prospective study. *Drug and Alcohol Dependence* 21(2): 147152.
36. Watkins EO (1976) *The Watkins Bender Gestalt Scoring System*. CA: Academic Therapy, Novato, USA.