

Antioxidant and phytochemical analysis of volatile oil and extracts of *Pinus wallichiana*

Abstract

The present study aims to explore the chemical constituents and antioxidant potential of *Pinus wallichiana* essential oil and extracts. Chemical composition of the essential oil was analyzed by GC-MS and result showed that the oil contain three major monoterpene compounds viz. α -pinene (48.6%), β -pinene (45.6%) and limonene (5.6%). Antioxidant potential of *Pinus wallichiana* was analyzed by three basic methods; DPPH radical scavenging method, FRAP assay and Fe²⁺ ion chelating activity. Among leaf and bark extracts, best radical scavenging, activity was determined by aqueous extract (IC₅₀ values 20.83±0.8µg/ml) of bark followed by its methanol extract (IC₅₀ value 25.9±1.6µg/ml). The aqueous extract of bark also exhibited better reducing and chelating activity than leaf extracts. Essential oil showed moderate radical scavenging and chelating activity but negligible reducing activity. Phytochemical analysis of extracts determined that the aqueous and methanol extracts of bark contain rich amount of poly phenol and flavonoids, in comparison to the leaf extracts. A significant correlation between the antioxidant activity and these phytoconstituents of the extracts has been observed. The results of the present study concluded that the bark of *Pinus wallichiana* is a potential source of active antioxidant constituents that could be further explored and exploited for numerous commercial applications.

Keywords: *Pinus wallichiana*, antioxidant, phytochemicals

Volume 4 Issue 2 - 2019

Pooja Sharma, Sahil Gupta, Nazli Bhatt, Shajaat Hussain Ahanger, Divya Gupta, Pramveer Singh, Roshani Lochan, Madhulika Bhagat

School of Biotechnology, University of Jammu, India

Correspondence: Madhulika Bhagat, School of Biotechnology, University of Jammu, Jammu, J&K-180006, India, Tel 011-9419124018, Email madhulikast@gmail.com

Received: November 23, 2018 | **Published:** April 30, 2019

Introduction

Oxidative stress and their adverse effect on human health has become a serious problem. Under stress conditions, there is increase in production of reactive species viz., hydrogen peroxide (H₂O₂), superoxide ion (O₂⁻), and hydroxide radical (OH) during cellular respiration than enzymatic antioxidants (e.g., superoxide dismutase (SOD), glutathione peroxidase (GPx), and catalase) and non-enzymatic antioxidants (e.g., ascorbic acid (vitamin C), α -tocopherol (vitamin E), glutathione, carotenoids, and flavonoids). This imbalance between reactive oxygen species production and the antioxidant defence could cause damage to cellular bio-molecules like carbohydrates, proteins, lipids and DNA, that may leads to metabolic and genetic alteration.¹ A lack of antioxidants could facilitates the development of degenerative diseases,² including cardiovascular diseases, cancers,³ neurodegenerative diseases, Alzheimer's disease⁴ and inflammatory diseases.⁵

Medicine plants are effective in treating various diseases caused by oxidative stress. The medicinal properties of plants are because to their biologically active metabolites like phenolic and flavonoid compounds, which are potential antioxidants.^{6,7} Positive correlation between these phytochemicals in plant extracts and antioxidant activity has been reported b many researchers.^{6,8,9}

Pinus wallichiana commonly called Blue pine or *Kail*, is evergreen coniferous tree native to Himalaya, grows in mountains at altitudes of 1800–4300m. it yields superior timber and especially used as joinery wood. The wood yields oleoresin (by tapping) which by distillation furnishes turpentine and rosin. It is a source of numerous valuable bye products, such as gases, light and heavy tar oils, turpentine

pitch, pyrolygneous acids, etc. The residue of wood is withdrawn as charcoal. *Pinus wallichiana* essential oil produced by distillation is used for medical purposes. Earlier, some efforts were made to use leaves by reducing them to fibre which was used for weaving into medicinal underclothing, for the manufacture of coarse matting, resembling coconut matting, for surgical dressings. During the recent past, in many rural areas of Kashmir, blue pine wood pieces were used to obtain a dark brown, viscous and sticky substance, called *killam*. This substance was traditionally applied by farmers on their arms and legs to protect them from insects (*Khase*) while working in water logged paddy fields. *Killam* would stick firmly; thereby protecting the exposed parts from insect bites (*Khase*), etc.¹⁰ So keeping the immense traditional importance of *Pinus wallichiana* in view, the present study aims to explore the phytochemicals and antioxidant properties of different parts of *Pinus wallichiana*.

Material and methods

Plant material and Preparation of extracts

Spines and bark of *Pinus wallichiana* was collected from China region of Bhaderwah, Jammu and Kashmir. Identification of the plant material was done an expert taxonomist of Department of Botany, University of Jammu, Jammu. Plant material (leaf and bark separately) was shade dried was powdered in an electronic grinder for the preparation of extracts. Three types of extracts were prepared in three different solvents viz., water, methanol and chloroform. 200g of dried powder was extracted thrice for each solvent. Resulting extracts were pooled, filtered and evaporated by using rotary vacuum evaporator. The slurry obtained was finally lyophilized to dried powder.¹¹

Isolation and chemical analysis of essential oil

Fresh spines of *Pinus wallichiana* were subjected to hydro distillation in a Clevenger type apparatus for 3-4h for isolation of essential oil. The extracted oil was dried by adding anhydrous sodium sulphate and stored at low temperature.¹² Chemical composition of the isolated essential oil was analyzed at Indian Institute of Integrative Medicine (CSIR, India), Canal Road, Jammu, India. System used for analysis is GC-MS 4000 (Varian, USA) with a Varian CP-SIL 8CB column (30m×0.32mm i.d., 1µm film thickness). Injector temperature was 230°C. Oven temperature program used was holding at 60°C for 5min, heating to 250°C at 3°C/min. Helium as a carrier gas used at a constant flow of 1.0 ml/min and an injection volume used was 0.20µl. The Mass spectrometer scan parameters included electron impact ionization voltage of 70 eV, a mass range of 40–500m/z. The components present in essential oil were analyzed by comparing their mass spectra with the compounds available in NIST05 (version 2.0) library.¹³

DPPH radical scavenging activity

Antiradical activity of the extracts and essential oil was determined by DPPH method. Essential oil was analyzed by method given by Bozin et al.¹⁴ 1ml of different concentrations of the test sample was mixed with 1ml of a DPPH (90µM, in methanol), and final volume was made to 4ml with methanol. The mixtures were kept at 25°C in the dark for 1hour. The absorbance was measured at 517nm using a UV-VIS spectrophotometer.

Free radical scavenging activity of *Pinus wallichiana* extracts was determined by DPPH method given by Abe et al.¹⁵ with some modifications. 1ml of DPPH (0.5mM in methanol) solution was mixed to 2 ml sample and to this 2ml of 0.1 M sodium acetate buffer (pH 5.5) was added. The mixtures were well shaken and kept at room temperature in the dark for 30min. The absorbance was measured at 517nm. The radical scavenging activity was calculated as a percentage of DPPH radical discoloration, using the equation:

$$\% \text{Radical Scavenging Activity} = \left[\frac{(A_0 - A_s)}{A_0} \right] \times 100$$

Where, A_0 is the absorbance of the control and A_s is the absorbance of the test compound.

FRAP assay

Reducing power of *Pinus wallichiana* extracts was analyzed by FRAP assay according to the method followed by Li et al.¹⁶ FRAP reagent was prepared in acetate buffer (300mM) by adding 10mM 2,4,6-tri (2-pyridyl)-s-triazine (TPTZ) solution in 40mM HCl and 20mM FeCl_3 solution in proportion of 10:1:1 (v/v), respectively. FRAP reagent was prepared fresh at the time of use. 50µl of the sample was added to 1.5ml of the FRAP reagent and after 3-4min, absorbance was measured at 593nm. A standard curve was prepared by using FeSO_4 (100-2000µM) and the result was expressed as µmol Fe (II)/gm dry weight of extract.

Chelation activity

The chelating power of the *Pinus wallichiana* Extracts on ferrous ions was determined by the method given by Dinis et al.¹⁷ Test sample (500µl) and 740µl of methanol were added to 20µl of 2mM FeCl_2 . The reaction was initiated by the addition of 40µl of 5mM ferrozine into

the mixture. The reaction mixture was incubated at room temperature for 10min and finally the absorbance was measured at 562nm. The ratio of inhibition of ferrozine- Fe^{2+} complex formation was calculated using the equation:

$$\% \text{ chelating power} = \left[\frac{(I_0 - I_s)}{I_0} \right] \times 100$$

Where, I_0 is the absorbance of the control and I_s is the absorbance of the test compound.

Total phenols and flavonoids in extracts

Total phenolic content in the extracts was estimated by Folin-Ciocalteu method.¹⁸ 0.5ml of extract was mixed with 0.5ml of 1N Folin-Ciocalteu reagent and 1ml of 20% Na_2CO_3 . After 10 min of incubation, the absorbance was measured at 750nm Gallic acid was used as standard compound to estimate total phenolic content and was expressed in mg GAE/g of the extract.

Total flavonoid content of the extracts was estimated by AlCl_3 colorimetric method.¹⁹ Plant extracts were diluted with distilled water to a volume of 3.5ml and added 150µl of a 5% NaNO_2 solution. After 5min, 300µl of 10% AlCl_3 solution was added. After 6min, 300µl of 1M NaOH and 550µl of distilled water were added. The mixture was well shaken and absorbance was measured at 510nm in UV-VIS spectrophotometer. Quercetin was used as standard compound to estimate total flavonoid content in extracts and was expressed in mg GAE/g of the extract.

Results and discussion

Hydro-distillation of fresh needles of *Pinus wallichiana* yielded 1.2% light yellow coloured essential oil. Chemical composition of the essential oil was analyzed by Gas Chromatography and Mass Spectrometry (GC-MS). Analysis showed the presence of three major components α -pinene (48.6%), β -pinene (45.6%) and limonene (5.6%) accounting for 99.8% of total components of the essential oil. All the three compounds found in essential oil belong to monoterpene class of terpenoids. Dar et al.²⁰ also analyzed the chemical composition of the essential oil of *Pinus wallichiana* growing in high altitudes of Kashmir, India. They found diverse range of compounds in essential oil but still in agreement that β -pinene (46.8%) and α -pinene (25.2%) are the major components of the essential oil. The difference in chemical composition might be due to spatial variation as the plant samples are from two different regions of Jammu and Kashmir.

Antioxidant activity of *Pinus wallichiana* was analyzed by three different methods viz., DPPH radical scavenging activity, metal ion chelating activity and FRAP assay. The results of antioxidant activity have been given in Table 1. The DPPH radical scavenging activity was determined on the basis of concentration providing 50% inhibition. Among the leaf and bark extracts, highest antiradical potential was observed in aqueous and methanol bark extracts with IC_{50} values 20.83±0.8µg/ml and 25.9±1.6µg/ml respectively. Chloroform extract of both leaf and bark have shown negligible radical scavenging activity. Essential oil isolated from leaf part showed moderate radical scavenging activity with IC_{50} values 514.4±5.8µg/ml.

The ability of antioxidants to reduce Fe^{3+} to Fe^{2+} in the presence of TPTZ, was determined in FRAP assay. Reducing potential of *Pinus wallichiana* extracts and essential oil was expressed in µmol Fe (II)/g dry weight of plant material. Strongest antioxidant activity

to reduce the Ferric ions (Fe^{3+}) was observed in bark aqueous extract (3495±11.3µmol Fe (II)/g) followed by bark methanol extract (2530±6.3µmol Fe (II)/g) like in case of radical scavenging activity.

Table 1 Total Phenols, flavonoids and antioxidant potential of *Pinus wallichiana* extracts

Extract	TPC (mg GAE/g dry weight)	TFC (mg QE/g dry weight)	DPPH Assay (IC50 in µg/ml)	FRAP Assay (µmol FeII /g)	%age Chelation capacity at 500µg
(Leaves)	30.8±1.1	-	260.4±3.2	111.6±3.2	57±1.2
	40±1.1	10±0.5	285.3±4.1	86.6±1.3	19±0.6
	Methanolic	10±0.4	-	26.6±2.1	34±0.4
	Aqueous	-	-	514.4±5.8	-
(Bark)	330±6.4	520±12.7	25.9±1.6	2530±6.3	59±1.8
	447±9.6	610±11.2	20.83±0.8	3495±11.3	44±1.4
	Methanolic	30±0.5	-	245±3.3	29±0.9
Aqueous	-	-	-	-	-
Chloroform	-	-	-	-	-

Data presented as mean±standard deviation; TPC, Total phenolic content; TFC, Total flavonoid content; GAE, Gallic acid equivalent; QE, Quercetin equivalent.

Chelating property on metal ions is one of important mechanism of antioxidant activity. Metal ion chelating capacity was determined at 0.5mg/ml concentration. Methanol extract of both leaf and bark of *Pinus wallichiana* showed a better chelating effect on ferrous ions (59±1.8% & 57±1.2%). Aqueous extract of bark also showed moderate chelating capacity with 47% at 0.5mg/ml. Essential oil and chloroform extracts determined poor chelating capacity of Fe^{2+} ions. Biological properties of phenolic and flavonoid compounds have already been known universally. Table 1 also represented total phenol and total flavonoid content in leaf and bark extracts of *Pinus wallichiana*. Highest amount of poly phenols and flavonoids was observed in aqueous extract of bark (447±9.6mg GAE/g; 610±11.2 QE/g dry weight, respectively) followed by methanol extract of bark (330±6.4 mg GAE/g; 520±12.7 QE/g dry weight). Leaf extracts possess comparatively less poly phenol and flavonoid content. Among all the extracts, least content of poly phenol and flavonoid was observed in chloroform extracts of both leaf and bark. Many studies have shown that the antioxidant activities in the plants are associated with their phytochemicals contents. Naeem et al.²¹ reported that the methanol extract of *Pinus wallichiana* (bark) growing in Kuldana, Murree (Pakistan), contained a wide variety of flavanols in considerable amounts. Dar et al.²⁰ analyzed that the essential oil isolated from *Pinus wallichiana* growing in Kashmir (J&K), has strong antiproliferative potential against five human cancer cell lines and good radical scavenging activity on DPPH radicals. Maimoona et al.²² also reported antioxidant potential and phytochemicals in different extracts of *Pinus wallichiana* growing in Pakistan and found a positive correlation between the antioxidant properties and total

All leaf extracts and essential oil exhibited very poor reducing activity in comparison to the bark extracts.

flavonoid content. This may be due to their redox properties of the phenolic and flavonoids that make them good reducing, scavenging and chelating agent.^{18,23} The results also revealed that there is significant correlation between the antioxidant activity (Reducing activity, Radical scavenging activity, and chelating power) and the phenolic/flavonoid content of *Pinus wallichiana* extracts. Therefore, it is possible that these phytochemical compounds are the major contributor for the antioxidant activity of the *Pinus wallichiana*.

Conclusion

In conclusion, the present study explored the phyto constituent and antioxidant properties of *Pinus wallichiana* growing in Baderwah region of Jammu & Kashmir (India). GC-MS analysis showed that its essential oil is a rich source of α -pinene and β -pinene and possess moderate antioxidant activity. Bark of *Pinus wallichiana* determined strong antioxidant activity and can serve as a valuable source of natural antioxidant. A significant correlation between antioxidant activity and TPC/TFC were found indicating phenolic and flavonoids compounds are the major contributor of antioxidant potential of *Pinus wallichiana*.

Acknowledgments

Authors would like to acknowledge the School of Biotechnology and Department of Bio-informatics, University of Jammu, for their support. The authors also wish to acknowledge Instrumentation division, IIM-Jammu, J&K (India) for their kind support.

Conflicts of interest

Author declares that there are no conflicts.

References

- Halliwell B, Gutteridge JMC. Oxygen toxicity, oxygen radicals, transition metals and disease. *Biochem J*. 1984;219(1):1–14.
- Shahidi F, Janitha PK, Wanasundara PD. Phenolic antioxidants. *Crit Rev Food Sci Nutr*. 1992;32(1):67–103.
- Gerber M, Boutron-Ruault MC, Hercberg S, et al. Food and cancer: state of the art about the protective effect of fruits and vegetables. *Bulletin of Cancer*. 2002;89(3):293–312.
- Matteo DV, Esposito E. Biochemical and therapeutic effects of antioxidants in the treatment of Alzheimer's disease, Parkinson's disease, and amyotrophic lateral sclerosis. *Curr Drug Targets CNS Neurol Disord*. 2003;2(2):95–107.
- Sreejayan N, Rao MNA. Free radical scavenging activity of curcuminoids. *Drug Res*. 1996;46(2):169–174.
- Cai Y, Luo Q, Sun M, et al. Antioxidant activity and phenolic compounds of 112 Chinese medicinal plants associated with anticancer. *Life Sci*. 2004;74(17):2157–2184.
- Pietta PG. Flavonoids as antioxidants. *J Nat Prod*. 2000;63(7):1035–1042.
- Duh P. Antioxidant activity of water extract of four Harg Jyur (*Chrysanthemum morifolium* Ramat.) varieties in soybean oil emulsion. *Food Chem*. 1999;66(4):471–476.
- Guleria S, Tiku AK, Rana S. Antioxidant and free radical scavenging activity of acetone extract /fractions of Terminalia bellerica Roxb fruit. *Indian J Biochem Biophys*. 2010;47(2):110–116.
- Dar AR, Dar GH. The Wealth of Himalayas–Gymnosperms. *Asian Journal of Plant Sciences*. 2006;5(2):251–259.
- Thusoo S, Gupta S, Sudan R, et al. Antioxidant Activity of Essential Oil and Extracts of *Valeriana jatamansi* Roots. *BioMed Research International*. 2014. 4 p.
- Guleria S, Tiku AK, Gupta S, et al. Chemical composition, antioxidant activity and inhibitory effects of essential oil of *Eucalyptus teretecornis* grown in north–western Himalaya against *Alternaria alternata*. *J Plant Biochem Biotechnol*. 2011;21(1):44–50.
- Adams R. *Identification of essential oil components by gas chromatography/mass spectrometry*. 5th ed. Texensis Publishing; 2007. 5 p.
- Bozin B, Mimica–Dukic M, Simin N, et al. Characterization of the volatile composition of essential oils of some *lamiaceae* species and the antimicrobial and antioxidant activities of the entire oils. *J Agric Food Chem*. 2006;54(5):1822–1828.
- Abe N, Murata T, Hirota A. Novel DPPH Radical Scavengers, Bisorbicillinol and Demethyltrichodimerol, from a Fungus. *Bioscience Biotechnology and Biochemistry*. 1998;62(4):661–666.
- Li HB, Wong CC, Cheng KW, et al. Antioxidant properties *invitro* and total phenolic contents in methanol extracts from medicinal plants. *LWT–Food science and technology*. 2008;41(3):385–390.
- Dinis TCP, Madeira VMC, Almeida LM. Action of phenolic derivatives (acetoaminophen, salicylate, and 5–aminosalicylate) as inhibitors of membrane lipid peroxidation and as peroxy radical scavengers. *Arch Biochem Biophys*. 1994;315(1):161–169.
- Chang ST, Wu JH, Wang SY, et al. Antioxidant activity of extracts from *Acacia confusa* bark and heart wood. *J Agric Food Chem*. 2001;49(7):3420–324.
- Jia ZS, Tang MC, Wu JM. The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chemistry*. 1999;64(4):555–559.
- Dar MY, Shah WA, Mubashir S, et al. Chromatographic analysis, anti–proliferative and radical scavenging activity of *Pinus wallichiana* essential oil growing in high altitude areas of Kashmir, India. *Phytomedicine*. 2012;19(13):1228–1233.
- Naeem I, Taskeen A, Mubeen H, et al. Characterization of Flavonols Present in Barks and Needles of *Pinus wallichiana* and *Pinus roxburghii*. *Asian Journal of Chemistry*. 2010;22(1):41–44.
- Maimoona A, Naeem I, Shujaat S, et al. Comparison of Radical Scavenging Capacity of Different Extracts of Barks and Needles of *Pinus roxburghii* and *Pinus wallichiana*. *Asian Journal of Chemistry*. 2011;23(2):819–822.
- Guleria S, Tiku AK, Singh G, et al. *In vitro* antioxidant activity and phenolic contents in methanol extracts from medicinal plants. *J Plant Biochem Biotechnol*. 2012;22(1):9–15.