

IRBIT a master regulator of cell physiology

Abstract

Hormones and neurotransmitters regulate cell functions by binding to their receptors, which activate intracellular signaling and produce the physiological response.¹ There are several intracellular pathways, including but not limited to, leading to the activation in protein kinases or phosphatases and increase of intracellular calcium (Ca^{2+}).¹ Calcium signaling controls many cellular responses (e.g. exocytosis, cell spreading, muscle contraction). Therefore it is important to understand the mechanisms regulating intracellular Ca^{2+} . Two major sources exist to raise intracellular Ca^{2+} : extracellular medium and internal stores (e.g., endoplasmic reticulum, mitochondria). While Ca^{2+} channels at the plasma membrane regulate Ca^{2+} entry in the cell, the inositol-1,4,5-trisphosphate receptor (IP_3R) controls the release of Ca^{2+} from the endoplasmic reticulum.^{2,3} Many proteins are known to bind to the IP_3R and regulate Ca^{2+} release.^{3,4} These include IRBIT (inositol-1,4,5-trisphosphate (IP_3) receptors binding protein released with IP_3), a protein discovered by Mikoshiba's group.^{4,5} Over the past decade IRBIT has emerged not only as an IP_3R regulator but also as an important regulator of other cellular functions.

Volume 4 Issue 1 - 2017

Kelli Cook, Natasa Petreska, Patrice G Bouyer

Department of Biology, Valparaíso University, USA

Correspondence: Patrice G Bouyer, Department of Biology, Valparaíso University, USA Tel (219) 4645487, Fax (219) 4645489, Email patrice.bouyer@valpo.edu

Received: February 13, 2017 | **Published:** February 22, 2017

IRBIT Regulates Intracellular Ca^{2+} Release from IP_3R

The IP_3R interacts with specific proteins (e.g. Huntingtin, RACK1) and forms a macrocomplex that is believed to provide the diversity and specificity of the Ca^{2+} signaling.⁴ IRBIT was discovered while searching for novel binding partners of the IP_3R .⁵ IRBIT negatively regulates the IP_3R by competing with IP_3 .⁶ IRBIT interaction with the IP_3R (thus regulation of IP_3 binding to its receptor) is decreased when IRBIT is phosphorylated as protein phosphatase (e.g. protein phosphatase 1) lowers IRBIT affinity to the IP_3R . Furthermore, phosphorylation of IRBIT's serine 68 plays a critical role in IRBIT inhibiting IP_3 from binding to its receptor.^{6,7} In addition to the phosphorylation state of IRBIT, two motifs (PEST and PDZ) were identified by in-silico analysis.⁸ Pull-down assays with IRBIT missing the PEST or PDZ motif demonstrated that these two motifs are important for the binding of IRBIT to the IP_3R .⁸

IRBIT a regulator of membrane transporter

As pointed above, IRBIT is released from the IP_3R after IP_3 binds to its receptor. Thus Mikoshiba's group hypothesized that IRBIT could be a new signaling molecule that interacts with downstream targets. By screening proteins that can bind to IRBIT they found that IRBIT interacts with the electrogenic sodium bicarbonate, NBCe1-B variant (aka pancreatic isoform). Using a heterologous system to co-express IRBIT and NBCe1-B, they showed by electrophysiological measurements that the transport activity of NBCe1-B is substantially increased in the presence of IRBIT.⁹ Another study demonstrated that IRBIT increases NBCe1-B activity by masking in part the auto inhibitory domain of NBCe1-B.¹⁰ These experiments suggest that IRBIT by increasing the activity of NBCe1-B, plays an important role in acid-base homeostasis.

The Na/H exchanger 3 (NHE3) participates in sodium (Na^+) reabsorption in the kidney. Angiotensin II (a hormone elevating Ca^{2+} via IP_3) increases Na^+ reabsorption by stimulating NHE3 in the kidney,¹¹ but the mechanisms by which angiotensin II activates NHE3 are not clear. Using a yeast two-hybrid system He et al. found that IRBIT interacts with NHE3.¹² In addition, knocking down

IRBIT abrogates the stimulatory effect of Ca^{2+} and angiotensin II on the activity of NHE3.^{12,13} These studies provide new ways of better understanding how angiotensin II, via the Ca^{2+} signaling stimulates Na^+ reabsorption in the kidney.

In the pancreatic duct, fluid secretion is driven by an active transcellular secretion of bicarbonate (HCO_3^-). The basolateral NBCe1-B pumps HCO_3^- inside the cell while the apical cystic fibrosis transmembrane regulator (CFTR) and Cl/HCO_3^- exchanger SLC26a6 secrete HCO_3^- in the pancreatic duct lumen. In addition to NBCe1-B, it was found that IRBIT binds to CFTR and SLC26a6 and increases transport activity of both transporters. The authors proposed that IRBIT coordinates the activity of basolateral and apical transporters involved in HCO_3^- -mediated fluid secretion.^{14,15} The results from these studies describe a new process allowing the crosstalk between the apical and basolateral membrane.

Finally, IRBIT has also been reported to bind with the Na-K-2Cl cotransporter 1 (NKCC1) in human T84 cells. Silencing IRBIT in this model decreased the cAMP-stimulated Cl^- secretion presumably by decreasing the activity of NKCC1.¹⁶

IRBIT a synergistic mediator of the cAMP and Ca^{2+} signaling pathway

In secretory epithelia, Ca^{2+} and cAMP signaling pathways control secretion. Both pathways can have an additive or synergistic action.¹⁷⁻¹⁹ For a more detailed discussion on how intracellular Ca^{2+} potentiates the cAMP pathway and vice versa and synergistic effect of Ca^{2+} and cAMP we refer the reader to the review by Ahuja et al.¹⁹ To date, the mechanistic details behind the synergistic action of Ca^{2+} and cAMP remain elusive, but work in the pancreatic duct and the salivary glands has shown that IRBIT can mediate the synergistic activation of the CFTR and SLC26a6 by cAMP and Ca^{2+} .¹⁵ Further work is now needed to test whether IRBIT mediates the synergistic effect of cAMP and Ca^{2+} in other tissues and whether additional proteins participate in the synergistic mechanism.

Other functions of IRBIT

Because of the numerous downstream targets of Ca^{2+} it is not

surprising that IRBIT has additional roles other than the ones discussed above. For instance, IRBIT interacts with the cleavage and polyadenylation specificity factor, which plays a role in mRNA processing. This study suggests that IRBIT could regulate gene expression.²⁰ IRBIT also binds to the phosphatidylinositol phosphate kinase members, who are potent regulators of ion transporters and would provide another level of regulation for electrolyte and solute transport in cells.²¹ IRBIT has been reported to inhibit the calcium calmodulin-dependent kinase II alpha in the brain and thus, would contribute to the regulation of the amount of catecholamine in the brain.²² IRBIT null mice show abnormal social behavior and hyperactivity that can be attributed to the level of catecholamine.²² Most recently IRBIT has been demonstrated to be involved with cell death by blocking the Bcl2l10 protein and facilitating large movements of Ca^{2+} to the mitochondria and thus promoting apoptosis.²³

Conclusion

Although some of IRBIT's cellular functions have started to be uncovered, many unknowns about IRBIT still exist. For instance, there are two isoforms of IRBIT a short and long N-terminus IRBIT.²⁴ Most of the studies have focused on the short N-terminus, but the long N-terminus is also capable of binding to NBCe1-B and decreases the Mg^{2+} -sensitivity of NBCe1-B inhibition during low HCO_3^- .²⁵ The expression and distribution of the two isoforms within tissues is not known. It remains to be determined whether the long and short IRBIT share common target proteins, and if so what is the impact of each isoform on the function of the target protein. Experiments have shown that a phosphorylation state of IRBIT is critical for carrying out its functional role; however, beside this rapid regulation we know very little about the rapid regulation of IRBIT by other proteins (e.g. kinases) or IRBIT gene expression. One study shows that in avian's oviduct, estrogen agonist induced IRBIT expression in luminal and glandular epithelial cells of the oviduct.²⁶ Because of IRBIT involvement in Na^+ and HCO_3^- transport we may expect that acid-base status (e.g. metabolic acidosis, and alkalosis), blood pressure or hormones involved in Na^+ homeostasis (e.g. aldosterone, angiotensin II) influence IRBIT expression. Because of its role as a third messenger released from the IP_3R it is expected that IRBIT will play essential roles in health and diseases.

Acknowledgements

The authors would like to thank Lynn Wellsand for her administrative assistance. PBG is supported by the Start up funds from Valparaiso University Art & Science College and the Wheat Ridge Ministries O.P. Kretzmann Grant for Research in the Healing Arts and Sciences.

Conflict of interest

The author declares no conflict of interest.

References

- Roman LA. Signal transduction. In: *Medical Physiology A cellular and molecular approach*. In: Boron WF, Boulpaep EL, editors. 2003. p. 87–114.
- Rizzuto R, Pinton P, Brini M, et al. Mitochondria as biosensors of calcium micro domains. *Cell Calcium*. 1999;26(5):193–199.
- Mikoshiba K. IP_3 receptor/ Ca^{2+} channel: from discovery to new signaling concepts. *J Neurochem*. 2007;102(5):1426–1446.
- Mikoshiba K. Role of IP_3 receptor signaling in cell functions and diseases. *Adv Biol Regul*. 2015;57:217–227.
- Ando H, Mizutani A, Matsu-ura T, et al. IRBIT, a novel inositol 1,4,5-trisphosphate (IP_3) receptor-binding protein, is released from the IP_3 receptor upon IP_3 binding to the receptor. *J Biol Chem*. 2003;278(12):10602–10612.
- Ando H, Mizutani A, Kiefer H, et al. IRBIT suppresses IP_3 receptor activity by competing with IP_3 for the common binding site on the IP_3 receptor. *Mol Cell*. 2006;22(6):795–806.
- Devogelaere B, Beullens M, Sammels E, et al. Protein phosphatase-1 is a novel regulator of the interaction between IRBIT and the inositol 1,4,5-trisphosphate receptor. *Biochem J*. 2007;407(2):303–311.
- Devogelaere B, Nadif Kasri N, Derua R, et al. Binding of IRBIT to the IP_3 receptor: determinants and functional effects. *Biochem Biophys Res Commun*. 2006;343(1):49–56.
- Shirakabe K, Priori G, Yamada H, et al. IRBIT, an inositol 1,4,5-trisphosphate receptor-binding protein, specifically binds to and activates pancreas-type $\text{Na}^+/\text{HCO}_3^-$ cotransporter 1 (pNBC1). *Proc Natl Acad Sci U S A*. 2006;103(25):9542–9547.
- Lee SK, Boron WF, Parker MD. Relief of auto inhibition of the electrogenic $\text{Na}^+/\text{HCO}_3^-$ cotransporter NBCe1-B: role of IRBIT vs. amino-terminal truncation. *Am J Physiol Cell Physiol*. 2012;302(3):C518–C526.
- Houillier P, Chambrey R, Achard JM, et al. Signaling pathways in the biphasic effect of angiotensin II on apical Na^+/H^+ antiport activity in proximal tubule. *Kidney Int*. 1996;50(5):1496–1505.
- He P, Zhang H, Yun CC. IRBIT, inositol 1,4,5-trisphosphate (IP_3) receptor-binding protein released with IP_3 , binds Na^+/H^+ exchanger NHE3 and activates NHE3 activity in response to calcium. *J Biol Chem*. 2008;283(48):33544–33553.
- He P, Klein J, Yun CC. Activation of Na^+/H^+ exchanger NHE3 by angiotensin II is mediated by IRBIT, IP_3 receptor binding protein released with IP_3 , and CaMKII. *J Biol Chem*. 2010;285(36):27869–27878.
- Yang D, Shcheynikov N, Zeng W, et al. IRBIT coordinates epithelial fluid and HCO_3^- secretion by stimulating the transporters pNBC1 and CFTR in the murine pancreatic duct. *J Clin Invest*. 2008;119(1):193–202.
- Park S, Shcheynikov N, Hong JH, et al. IRBIT mediates synergy between Ca^{2+} and cAMP signaling pathways during epithelial transport in mice. *Gastroenterology*. 2013;145(1):232–241.
- Tang X, Tang J, Mykoniatis A, et al. W1779 Regulation of chloride secretion in the colonic crypt cell line T84 by Inositol 1, 4, 5-trisphosphate (IP_3) Receptor-Binding protein released with IP_3 regulates (IRBIT). *Gastroenterology*. 2010;138(5 Suppl 1):738.
- Cartwright CA, McRoberts JA, Mandel KG, et al. Synergistic action of cyclic adenosine monophosphate- and calcium-mediated chloride secretion in a colonic epithelial cell line. *J Clin Invest*. 1985;76(5):1837–1842.
- Vajanaphanich M, Schultz C, Tsien RY, et al. Cross-talk between calcium and cAMP-dependent intracellular signaling pathways. Implications for synergistic secretion in T84 colonic epithelial cells and rat pancreatic acinar cells. *J Clin Invest*. 1995;96(1):386–393.
- Ahuja M, Jha A, Maléth J, et al. cAMP and Ca^{2+} signaling in secretory epithelia: crosstalk and synergism. *Cell Calcium*. 2014;55(6):385–393.
- Kiefer H, Mizutani A, Iemura S, et al. Inositol 1,4,5-trisphosphate receptor-binding protein released with inositol 1,4,5-trisphosphate (IRBIT) associates with components of the mRNA 3' processing machinery in a phosphorylation-dependent manner and inhibits polyadenylation. *J Biol Chem*. 2009;284(16):10694–10705.

21. Ando H, Hirose M, Gainche L, et al. IRBIT Interacts with the Catalytic Core of Phosphatidyl inositol Phosphate Kinase Type Ialpha and IIalpha through Conserved Catalytic Aspartate Residues. *PLoS One*. 2015;10(10):e0141569.
22. Kawaai K, Mizutani A, Shoji H, et al. IRBIT regulates CaMKII alpha activity and contributes to catecholamine homeostasis through tyrosine hydroxylase phosphorylation. *Proc Natl Acad Sci U S A*. 2015;112(17):5515–5520.
23. Bonneau B, Ando H, Kawaai K, et al. IRBIT controls apoptosis by interacting with the Bcl-2 homolog, Bcl2l10, and by promoting ER-mitochondria contact. *Elife*. 2016;5.
24. Devogelaere B, Sammels E, De Smedt H. The IRBIT domain adds new functions to the AHCY family. *Bioessays*. 2008;30(7):642–652.
25. Yamaguchi S, Ishikawa T. AHCYL2 (long-IRBIT) as a potential regulator of the electrogenic Na⁺-HCO₃⁻ cotransporter NBCe1-B. *FEBS Lett*. 2014;588(5):672–677.
26. Jeong W, Kim J, Ahn SE, et al. AHCYL1 is mediated by estrogen-induced ERK1/2 MAPK cell signaling and microRNA regulation to effect functional aspects of the avian oviduct. *PLoS One*. 2012;7(11):e49204.