

Evaluation of shallow groundwater in Rural Kebbi State, NW Nigeria, using multivariate analysis: implication for groundwater quality management

Abstract

This study assessed shallow groundwater in rural Kebbi State using Pearson's Correlation (r), Factor Analysis (FA), and Hierarchical Cluster Analysis (HCA). One hundred (100) shallow groundwater samples were drawn randomly from hand-dug shallow wells in 10 Local Government Areas (LGAs). Physical parameters (pH, temperature, TDS, EC, salinity) were analysed in situ using hand-held metres. Separate water samples were taken to analyse ions (Fe, Zn, K, Mg, HCO_3^- , Ca). Results revealed that shallow groundwater is lightly acidic, except in Zuru and Fakai LGAs, where an alkaline condition occurred. Correlation analysis revealed that the dissolved solids in shallow aquifers result from natural geological and anthropogenic influences. However, FA had shown that most of the variability in shallow groundwater is influenced by natural geological factors with little evidence from anthropogenic inputs. The HCA categorised shallow groundwater into three groups: those shallow wells having higher concentrations of Zn and Cl with more elevated salinity and temperature; those shallow wells having very low salinity in central Kebbi State, and those shallow aquifers having alkaline waters. Except for higher Fe and Zn concentrations, the shallow aquifers contained water of excellent quality for drinking. Correlation analysis, FA, and HCA present simple statistical tools for assessing the hydrochemistry of groundwater.

Keywords: correlation, factor analysis, hierarchical cluster analysis, shallow aquifers

Volume 7 Issue 3 - 2022

Saadu Umar Wali,¹ Murtala Abubakar Gada,² Isah Hamisu,² Kabiru Jega Umar,³ Isa Garba Abor⁴

¹Department of Geography, Federal University Birnin Kebbi, Nigeria

²Department of Geography, Usmanu Danfodiyo University Sokoto, Nigeria

³Department of Pure and Industrial Chemistry, Federal University Birnin Kebbi, Nigeria

⁴Department of Mathematics, Usmanu Danfodiyo University Sokoto, Nigeria

Correspondence: Saadu Umar Wali, Department of Geography, Federal University Birnin Kebbi, P.M.B. 1057 Kebbi State, Nigeria, Tel +2348065555170, Email saadu.wali@fubk.edu.ng

Received: April 12, 2022 | **Published:** May 05, 2022

Introduction

The assessment of groundwater is an essential activity for rural and urban water management, particularly in semi-arid regions,¹⁻⁴ because the ionic composition of water has a significant influence on human health.⁵⁻⁷ Using low-quality groundwater for drinking and irrigation affects human health and plant growth and contaminates soil, making it less suitable for agriculture and household uses. Shallow groundwater is the primary source of potable water and is commonly used for household and irrigation uses in rural Nigeria,^{8,9} particularly in the semi-arid northern areas of Nigeria.¹⁰⁻¹² The question of portable and adequate water supply, especially in rural areas, has been unanswered for quite a long time,¹³⁻¹⁶ leaving the rural dwellers to exploit the shallow groundwater, which can be accessed using simple technologies.

Consequently, water quality is also of concern since most remote communities cannot conduct water analysis.¹⁷⁻¹⁹ While shallow groundwater quality in rural areas is poorly known, increasing exploitation of these water sources is driven by Malthusian trends and environmental change.²⁰⁻²² The uninterrupted increase in the rural population has led to many social, economic, and environmental problems. These problems include excessive shallow groundwater withdrawal for agricultural and domestic uses in arid and semi-arid environments.²³⁻²⁵ Shallow groundwater is the principal industrial, agricultural and domestic water supply.²⁶⁻²⁸ In some remote communities, shallow groundwater is the principal source of community water supply during the dry season.²⁹⁻³² It has the advantage of being the inexpensive and reliable source of community water supply,³³⁻³⁶ and this attracts even the more coordinated programs on improving rural water supply to resort to groundwater as the best, inexpensive and dependable source of water supply for community water supplies.³⁷⁻⁴⁰

Although shallow groundwater sources are widely used in developing countries,^{37,41,42} their drinking and agricultural use fitness is poorly investigated. However, shallow groundwater quality varies widely between environments and water sources consequence of geological background and land use.⁴³⁻⁴⁶ This erraticism can affect water quality and resultant consequences on human health and wellbeing.⁴⁷⁻⁵⁰ The variability of shallow groundwater composition is primarily influenced by the mineralogy of shallow aquifers⁵¹ and human activities (agriculture, industry, etc.), which alter the natural composition of groundwater and consequently its aptness for drinking and other uses, as they can unite to create varied water types that vary in hydrochemistry.

Shallow groundwater has been widely studied.^{26,36,39,45,50,52-55} Results indicate a need to trace the origin and mechanism of nitrate leakage into the subsurface aquifers of Maiduguri.²⁶ An appraisal of subsurface water quality showed that about half of the shallow groundwater is unsuitable for drinking in Nagpur, India.⁴⁵ Significant temporal and spatial variations in shallow groundwater temperatures have been revealed in the Lower Heihe River in northwestern China.⁵⁴ Assessment of health risks of heavy metal contaminations in shallow aquifers indicated that the heavy metal pollution index (HPI) and hazard quotient (HQ) were higher in more cultivated fields.⁵⁰

Kebbi State is situated in a semi-arid region of West Africa. Apart from human and geological influences on groundwater, the increasing climate vagaries posed a severe challenge to shallow groundwater management. Evaluation of shallow groundwater in the southern of Kebbi State⁵⁶ revealed that the aquifers are characterised by moderate potassium, calcium, chloride, copper, sodium, and bicarbonates concentrations below the World Health Organization (WHO) and Nigerian Standard for Drinking Water Quality (NSDWQ) reference standards. However, iron, zinc, magnesium, and phosphate concentrations are above the WHO and NSDWQ reference

guidelines.⁵⁷ The study also found that TDS correlates strongly with Cu, Fe, Zn, Mg, and PO₄, suggesting that these ions' dissolved solids were derived. However, these findings do not apply to the entire Kebbi State since the geological setting in southern Kebbi State is of a basement complex background. Therefore, this paper aims to assess shallow groundwater across Kebbi State.

The study area

Geographical setting

Kebbi State (Figure 1) is positioned between latitude 10° 8' and 13° 15'N and between longitude 3° 30' E. It occupies a land area of about 36,800 km².⁵⁸ The climate is characterised by a long dry season and a short but intensive wet season. Annual rainfall is highly variable and decreases in volume from southern to northern parts. The mean maximum temperature is highest in April (40°C) and the lowest in December (23°C). The mean relative humidity is maximum in August (90%) and is a minimum in December (10-30%). However, there is an overall increase in relative humidity southward.⁵⁸ The evaporation rate is high, and the state can be classified as having a ustic soil moisture regime.




Figure 1 Map of Kebbi State shows the study area and sampling locations.

Hydrogeological condition

The geology is typical of the geological setting of Northwestern Nigeria. It is mainly sedimentary, with some weathered basement complex rock outcrops in the south.^{59,60} Underlying the sedimentary

formations are crystalline rocks of Pre-cretaceous age.⁶¹ Groundwater in the upland areas of crystalline rocks of the southern part of Kebbi State is ordinarily accessible in small amounts from fissures or other horizontal separations and the regoliths.⁵⁹ River Sokoto (Gulbin-Dukku, locally called), a chief tributary of River Niger, essentially drains the study area.⁶² Despite the potential presented by aquifers in the state (e.g. Gwandu and Illo Formations), many communities are still without improved water supply systems and rely on individual and community hand-dug shallow wells for potable water supply. Figure 2 describes a hydrogeologic cross-section of the Sokoto basin.




Figure 2 Geohydrologic section of Sokoto basin, NW Nigeria, indicating principal aquifers and confining beds.

Although these hand-dug wells are poorly constructed, their water quality remains poorly known because most rural communities lack the capacity for water testing.^{19,63} However, the hydrological condition of northern and southern parts of Kebbi State showed that the underlying shallow aquifers are in hydraulic connectivity with surface water bodies.⁶⁴ Additionally, these aquifers are recharged primarily by infiltrating rainwater, making them prone to contamination. Therefore, proper land use practice is required to effectively manage shallow groundwater quality.⁵⁹

Materials and methods

Groundwater sampling and analysis

The study employed onsite field measurements and laboratory analysis (Table 1). Ten (10) Local Government Areas (LGAs) were purposely studied, and ten (10) communities from each LGA were studied at random. Therefore, one hundred (100) water samples were collected and analysed from Hand-dug shallow wells. Before taking water samples, the plastic containers were rinsed twice. First by deionised water and later with water from experimented shallow wells.

Table 1 Parameters, methods, description, detection limit and reference guidelines

Parameters	Methods	Description	Detection limits	Reference Guidelines	
				WHO (2011)	NSDWQ (2007)
Temperature	Field (DK-MsG01)	Temp/Salinity-meter	Ambient	Ambient	
pH	Field(pHep)	pH meter	-	6.5-8.5	6.5-8.5
Conductivity	Field (DIST)	EC/TDS- meter	uS/cm	-	-
Salinity	Field (DK-MsG01)	Temp/Salinity-meter	mg/l	0.5	0.5
TDS	Field (DIST)	EC/TDS- meter	"	1000	500
Fe	Laboratory	AAS	"	0.3	0.3
Zn	"	"	"	3	4
K	"	"	"	500	1000
Mg	"	"	"	3	4
Mg	"	"	"	200	200
HCO ₃	"	"	"	250	250
Ca	"	"	"	500	500

Conversely, separate water samples were collected for in situ analysis (using water quality Probes) of physical parameters (Temperature, pH, Conductivity/salinity, and TDS). Water quality probes were first rinsed with deionised water and then with water from sampled wells. Separate water samples were taken in 1-litre polyethylene bottles for Laboratory analysis. Water samples were stored in shielded bottles at a controlled temperature (<5°C). Groundwater water samples were filtered through Whatman Polycap GW encapsulated filter for groundwater samples (POLYCARP GW 75, 0.45µm, polyethersulfone). Table 1 summarises analytical methods, the World Health Organization (WHO), and Nigerian Standard for Drinking Water Quality (NSDWQ) reference standards.

Atomic Absorption Spectrometry (AAS) was used for ion analysis.⁶⁵ AAS is a technique in which free vaporous atoms absorb electromagnetic energy at an explicit wavelength to create a measurable indicator. The immersion of those free riveting atoms in the visual route is proportionate to the immersion indicator. Accordingly, for AAS dimensions, the analyte is initially transformed into vaporous traces, usually by directing heat to a cell (i.e., atomiser). The type of atomiser delineates the two major AAS-based analytic procedures: flame atomic absorption spectrometry (FAAS), which steadily proffers analytical indications and electrothermal atomic absorption spectrometry (ETAAS) conveying logical indicators in a sporadic mode (2 to 4/sample).

In both methods, water (or dissolved) samples are mounted into the analyser as an aerosol in FAAS instance for this analysis or as

an apparent low microliter amount in ETAAS. Monochromators founded Ebert, Littrow, and Czerny-Turner schemes are the best standard wavelength pickers applied in AAS. Newly, echelle optics is bonded to marketable AAS appliances. Launching an echelle design connected with a formidable uninterrupted lamp forms momentous capability, for example, the study of the spectral milieu close to the line and simultaneous multi-elemental analysis. The parameters were analysed using AAS (Rs 12.5 Lakh/Piece).

Statistical analysis

The hydrochemical data is summarised and standardised, as presented in Table 2. The data were further analysed using Pearson's correlation (*r*), Factor analysis (FA), and hierarchical cluster analysis (HCA). Pearson's correlations (*r*) were used to study the relationship between hydrochemical variables (Table 3). Factor analysis was used to measure the interrelationships between the studied hydrochemical parameters by lessening the detailed data to a more interpretable form [66-70]. Factor analysis is a multivariate statistical procedure that can be applied to reduce enormous and intricate datasets by identifying a small number of parameters that describe the most significant variation within the original data set. In this paper, FA was carried out on hydrochemical parameters to pinpoint and define the elements that influence the shallow groundwater composition in Kebbi State. The FA was carried out on a subset of 11 chosen hydrochemical parameters (Temperature, EC/Salinity, pH, TDS, K, Ca, Fe, Zn, Mg, Cl, HCO₃), which characterised the general groundwater composition.

Table 3 Person's (*r*) values in bold show a significant correlation between parameters

Parameter	Temp	EC	pH	Sal.	TDS	Fe	Zn	K	Mg	Cl	HCO ₃	Ca
Temp		0.86	0.155	0.382	0.986	0.567	0.028	0.035	0.859	0.311	0.428	0.834
EC	0.064		0.412	0.982	0.035	0.947	0.323	0.638	0.994	0.491	0.668	0.119
pH	-0.485	-0.293		0.408	0.485	0.4	0.709	0.253	0.358	0.311	0.244	0.975
Salinity	-0.311	0.008	0.295		0.831	0.729	0.928	0.472	0.514	0.389	0.755	0.879
TDS	0.006	0.669	0.25	-0.078		0.517	0.396	0.323	0.183	0.717	0.202	0.043
Iron	-0.207	-0.024	-0.3	0.126	-0.233		0.039	0.601	0.132	0.89	0.269	0.379
Zinc	0.686	0.349	-0.135	0.033	0.302	-0.657		0.366	0.207	0.71	0.718	0.498
Potassium	0.667	-0.17	-0.399	-0.258	-0.349	-0.189	0.321		0.46	0.24	0.612	0.746
Magnesium	0.065	-0.003	0.326	-0.234	0.459	-0.51	0.437	-0.264		0.839	0.206	0.668
Chloride	0.357	0.247	-0.357	-0.306	0.132	-0.05	0.135	0.409	-0.074		0.652	0.027
Bicarbonate	-0.283	-0.155	0.406	-0.114	0.441	-0.387	-0.131	-0.183	0.438	0.163		0.121
Calcium	0.076	0.525	-0.011	-0.056	0.647	-0.313	0.243	-0.118	0.155	0.692	0.523	

Hierarchical cluster analysis (HCA) was performed to understand groundwater composition by separating sampling LGAs into various clusters. It benefits from not requiring any prior theory of the number of sets.⁷¹⁻⁷³ Sampling locations with similar groundwater composition or otherwise parameters with typical properties would be grouped into matching clusters [66-70]. In this paper, Ward's-algorithmic clustering method consequent to the Euclidean distance was applied. This is deemed the most potent clustering tool since it can lessen the distorting impact or sum of squared distances of centroids from hypothetical binary clusters formed at every step.

Therefore, applying factor scores to HCA is an excellent method for groundwater analysis by simplifying groups into more unrestrained forms, despite the obliviousness of specific theoretically significant hydrochemical data. However, this method does not account for data validity and disconnected values and neglects credible and possible connections or resemblances (statistical meddling) between variables, which are somewhat clear in most hydrochemical data.⁶⁶⁻⁷⁰ As a result, the output would have been inexorable to manipulation and not be deemed as complete hydrochemical momentous data. Therefore, in this

paper, only raw data were included in the HCA, where full scrutiny of each cluster is possible, and accurate (without modification) condition of groundwater composition could be characterised.

Results and discussion

Physicochemical characteristics of groundwater

Table 2 summarises groundwater parameters (range, mean and standard error). The pH concentration varied in the study area. Mean pH was above 7 in Zuru and Fakai LGAs, indicative of alkaline conditions. The two LGAs are in the basement complex section of southern Kebbi State. Therefore, the observed alkaline condition perhaps results from the region's geology mainly comprises crystalline rocks. High pH in aquifers can be linked to the ion exchange process.^{74,75} It accelerates the evolution from calcium to soda through subsurface water metamorphisation with a successive increase in the carbonate substance and alkalinity of aquifers.⁷⁶

However, a high pH level of 11-12 was recorded in Paju City consequence of shallow groundwater pollution caused by improper

discarding of road constructions litter on the surface at the study location.⁷⁷ Arewa and Jega LGAs have a mean pH of 6.5, indicating a neutral condition. The seventeen LGAs have pH values of less than 5 displaying acidic conditions. Groundwater acidity has been reported from various parts of the world.^{78–80} Aerosol particles and acid gases are incorporated into cloud water and rainwater and deposited by rainfall or by tumultuous mixing and gravitational landing on vegetation, soil, unprotected water, and streams/rivers. The existence of rain tends to be correlated to increased anthropological emission of gaseous pollutants into the atmosphere, mainly sulfur and nitrogen oxides.⁸⁰

The most acidic water has occurred in the eastern section of the Sokoto basin and the south.⁵⁹ Just after Sokoto town, at Dange pH is 5.1 at Borehole GSN 3512, 3.7 at Borehole GSN 3520, and it is 3.7 at 177km at Borehole GSN 3519 along the Sokoto-Gusau road. Groundwater in this vicinity also has low TDS (28–79mg/l). The major anion is SO_4 , which might have been derived from the oxidation of pyrites.⁵⁹ This process perhaps accounts for the acidity of the water in the western Sokoto basin. The temperature varied between the studied LGAs. The mean groundwater temperature is lowest in Zuru and highest in Bunza. Elevated groundwater temperatures are strongly associated with increased chemical reactions in aquifers. A temperature increase by 10°C can lead to the doubling of chemical reactions in aquifers.⁸¹

Mean EC concentration is highest in Gwandu and Augie LGAs, indicating that the two LGAs have high dissolved rock mineral contents in shallow groundwater. Higher groundwater temperatures in the two LGAs could account for higher EC values.^{82,83} There tends to be an upward trend between elevated temperature and EC level in aquifers.⁸¹ Mean salinity values were lowest in Jega and Maiyama LGAs. At the same time, Gwandu LGA has the highest mean salinity, which exceeds WHO and NSDWQ reference guidelines (0.5mg/l). The observed high salinity can be derived from rock minerals and land use (irrigation), which is well pronounced in the Gwandu LGA. High salinity in subsurface water is derived from natural and anthropological processes.^{84–89} Most of the irrigated fields in Gwandu LGA are underlain by a very shallow groundwater table (often less than 2 meters below the ground surface). The continuous application of agrochemicals (especially chemical fertiliser) in these fields alters the chemical composition of shallow aquifers, which is hardly unconnected with high EC/salinity and TDS level in the area. Human activities, especially agriculture, posed a severe threat to groundwater quality.^{72,87,90–92}

Figure 3 shows the relative concentrations of groundwater parameters. Rises in pH accompanied an increase in groundwater temperature while both parameters decreased with an increased salinity level. This was not the expected outcome since EC values in groundwater show an upward trend with rising temperature and TDS levels (EPA, 2001). Similarly, the latter rises with a rise in EC, Zinc, and iron. However, an increase in pH level was also accompanied by an increase in salinity and EC levels. As a result, mean total dissolved solids (TDS) are within WHO and NSDWQ reference guidelines (Table 1).

The TDS is widely used in water quality analysis as it shows the number of dissolved rock materials in groundwater, which linger as a residue after water evaporation from the sample.^{93,94} The general outlook of the TDS concentrations in Kebbi State indicates low concentrations of dissolved solids in shallow groundwater. Results are concurrent with previous investigations on the hydrochemistry of the Sokoto basin.^{59,95,96} Lower TDS intensities (<500mg/l), is

exceptionally essential.⁹⁷ Based on TDS concentrations, shallow aquifers are appropriate for drinking. Iron concentration generally fell outside WHO and NSDWQ reference guidelines (Table 1) (Table 2). This finding concurred with the existing literature relating to the hydrochemistry of the Sokoto basin.^{53,59} Mineral iron is widely distributed on earth and is obtained from different sources.^{98,99} Except for high iron concentrations, groundwater in Kebbi State is generally fit for drinking based on anion chemistry. An elevated concentration of iron is quite harmful to aquatic life.^{100–102} Typically there is no effect on human health; instead, the problems are primarily aesthetic.⁸¹ Except for Birnin kebbi and Zuru LGAs, the mean zinc concentration fell outside the WHO and NSDWQ reference guidelines. Figure 3 shows an upward trend between Zinc and iron concentrations, even though the former showed a downward trend with increased TDS. Therefore, most of the observed TDS values are derived from iron.




Figure 3 The relative concentration of physicochemical parameters. Note: all concentrations are in mg/l except conductivity ($\mu\text{S}/\text{cm}$) and pH (unit).

Zinc concentration was higher in Arewa, Augie, Bagudo, and Maiyama LGAs. Mean zinc values in these LGAs exceed WHO and NSDWQ reference guidelines even though zinc is vital to a man if consumed in significant quantities.⁸¹ The noxiousness of zinc to aquatic life is (as with copper) contingent on the hardness of the water, which declines with increasing hardness.⁸¹ Potassium concentration is within the WHO and NSDWQ reference guidelines. Figure 4 shows the relative potassium, magnesium, chloride, bicarbonates, and calcium concentration. The observed consistency in potassium values results from homogeneity in soil chemistry since potassium is fixed in soils and is not merely seeped out.^{103–105}






Figure 4 The relative concentration of groundwater parameters. Note: all concentrations are in mg/l.

Magnesium is high and exceeds WHO and NSDWQ reference guidelines. Undue intake has not been related to any serious health threat, only that its concentrations are very significant when considered with sulfate.⁸¹ Alkali metals such as Ca joint with Mg are responsible for groundwater hardness. Magnesium is a relatively common reducing element. Temperature changes can impact aquifers by rising oxygen and increasing the reduction process. Also, Mg



can react with water vapour to produce hydrogen gas or magnesium hydroxide, as defined in equation 1. Magnesium constitutes a second major component of hardness (CaCO_3). Dolomite and magnesium carbonate are primary sources of Mg in aquifers.^{106–108}



Mean calcium is high in Bagudo and Bunza LGAs. Typically, Ca concentration is low in Kebbi State. The importance of Ca in the hydrochemical analysis also relates to hardness. Calcium is found naturally in various environmental settings and occurs widely in groundwater aquifers.^{109–111} It is an integral component of coral and is found in high concentrations (400 mg/l) of brine. In lime regions, Ca attains 100 mg/l. Elementary Ca at normal temperature reacts with water, based on the reaction process indicated by equation 2.



Dissolved calcium hydroxide forms soda and hydrogen gas. It typically occurs when CO_2 is freed, resulting in the development of carbonic acid, affecting Ca compounds. The carbon weathering and total reactions are defined in equations 3 and 4. Consequently, calcium hydrogen carbonate is produced.



Mean chloride concentration in Arewa, Bagudo, Bunza, Gwandu, and Maiyama exceeds WHO and NSDWQ reference guidelines (Table 2). As shown in Figure 4, a rise in chloride concentration accompanied rising Mg, HCO_3^- , and Ca levels. However, these ions are geologically unrelated to chloride. Chloride levels can be as high as 70 mg/l under dry season discharges from industrial and municipal sewage. High ingestion presents no health risks. Though, a higher Cl level in irrigation water might render it unsuitable for irrigation.^{112–114}

Combined with Ca and Mg, Bicarbonates develop carbonate hardness.^{115–117} Bicarbonate and carbonate ions merged with Ca, or Mg will precipitate as calcium carbonate (CaCO_3) or magnesium carbonate (MgCO_3) when the soil solution distillates under dry environments.⁴⁵ Consequently, the concentration of Ca and Mg declines relative to sodium, and the SAR index will increase.¹¹⁸ This causes an alkalisating effect and raises the pH level. Thus, when a water analysis shows a high pH level, it can signify high content of carbonate and bicarbonate ions.¹¹⁸ Bicarbonate is another alternative measure of the sodium content concerning Ca and Mg. Bicarbonate ion is formed consequential of bonding positively charged ion with negatively charged O_2 molecules, producing an ionic fused. Bicarbonates usually are soluble in water. Mean bicarbonate in the study area was greater in Bunza and Jega LGAs. The overall bicarbonate concentration in Kebbi State is within WHO and NSDWQ reference guidelines.

Statistical application


Correlation analysis

Table 3 shows the correlations between hydrochemical parameters. Temperature correlates significantly with EC. This was an expected outcome and concurred with most findings in the literature.^{119–121} Conductivity and salinity correlate significantly. This is also an expected outcome since the two parameters are often used interchangeably, as, in most literature, EC is used as an alternative measure of salinity.^{122,123} EC correlates significantly with Fe, K, HCO_3^- , Ca, and Mg. Thus, the dissolved solids are derived from these ions. pH was significantly correlated with Ca. In groundwater, pH and Ca tend to be correlated; the former can be altered by calcium hydroxide

and calcium carbonate. A reduction in pH consequent to precipitation of calcium carbonate has been noticed.¹²⁴

Salinity correlates significantly with all the analysed parameters, except K and Cl, indicative of the lesser contribution of these elements to salinity in the study area. TDS correlates significantly only with Fe and Cl, suggesting that the dissolved solids in shallow aquifers are derived from these ions. Magnesium was associated considerably with Ca and Cl. A significant correlation between Ca and Mg suggests the same source, calcite rocks.¹¹⁸ Chloride correlates significantly with HCO_3^- , meaning the same source, a natural geogenic source. Chloride is inherently found in aquifers since it is increasingly derived from anthropogenic sources (agrochemicals, industrial and urban sewage) and mineral lodes.^{125,126} Though chloride (Cl) is a stable ion, it is naturally combined with sodium and sporadically with Ca, Mg, and K. Natural aquifers rarely contain Cl above 50 mg/l, as any significant rise may lead to suspicion from anthropogenic inputs.

Bicarbonate is a transitional form within the *dehydration* of carbonic acid. It serves an essential role biochemically in the buffering system of pH [127–129]. Bicarbonate is the conjugate acid of CO_3^{2-} , and the conjugate is based on the acid H_2CO_3 (carbonic acid). The carbonate reaction is defined in equations 5 and 6:



The bicarbonate ion is produced by bonding positively charged ions with negatively charged O_2 molecules, creating an ionic composite. Bicarbonates are typically soluble in water.

Factor analysis

Factor analysis (FA) is a robust statistical technique that aids the explanation of groundwater composition.^{130–133} Factor analysis can be applied to reorganise data to describe better the characteristics of the environmental factors that produced the observed groundwater composition. Therefore, a set of factors (Table 4) was extracted to explain the interrelations between groundwater elements. A total of five factors were extracted. The five factors put together explained 88.20% of the total variance. Factor 1 accounts for 27.75% of the variance and is inferred as correlating primarily to the salinisation of the shallow groundwater from the dissolution of rock minerals.^{134,135} This factor has high positive loadings (≥ 0.65) on Zn, Mg, Ca, TDS, and EC. In addition, the major cation exchange elements Ca, and Mg correlate positively, which is also indicative of the effect of the dissolution of rock minerals.

Factor 2 has high positive loading only on pH, indicative of the influence of carbonates.^{124,136,137} Factor 3 had high positive loading of Fe and EC, suggesting salinisation of groundwater sources from rock minerals. Factor 4 had high positive loadings on salinity and Zn, further indicating the dissolution of rock minerals. Igneous rocks, e.g., granite and basalt, might contain 30, 0.6, 160, and 140 mg/kg of Pb, Cd, and Zinc.¹³⁸ Factor 5 had high positive loadings on TDS. It can also be correlated to rock weathering and human activities since the significant correlation between TDS and Fe and Cl was revealed earlier by Pearson's (r) as in Table 2. Anthropological inputs to groundwater have been shown via correlations between TDS and Cl or NO_3^- .^{139,140}

Hierarchical cluster analysis

Hierarchical cluster analysis is a widely used technique in groundwater analysis since it can differentiate groundwater composition.^{71–73,76,141–145} By applying HCA, the sampling LGAs with comparable groundwater hydrochemistry can be merged into

a single cluster. The graphical assemblies of the clustering process are illustrated in Figure 5. Based on the dendrogram, sampling LGAs with matches are collected into three different sets. The first cluster (group 1) sampling LGAs: Arewa, Birnin kebbi, Bunza, Maiyama, Bagudo, and Gwandu. Groundwater composition in these LGAs is characterised by higher chloride, zinc, and higher salinity and temperature levels. The high temperature in aquifers can be related to external influences (climatic factors) or geologic effects.^{146,147}

Table 4 Factor analysis (Varimax rotation), values in bold signify high loadings

Parameter	Factor 1	Factor 2	Factor 3	Factor 4	Factor 5
Temp	0.402	-0.78	-0.26	0.111	-0.044
EC	0.525	-0.053	0.62	0.495	-0.12
pH	-0.004	0.778	-0.355	-0.024	0.237
Sal	-0.277	0.308	0.082	0.505	0.695
TDS	0.74	0.414	0.292	0.203	-0.177
Fe	-0.641	-0.102	0.594	-0.015	-0.214
Zn	0.675	-0.336	-0.391	0.505	0.119
K	0.094	-0.798	-0.288	-0.244	0.255
Mg	0.56	0.386	-0.479	0.039	-0.387
Cl	0.513	-0.426	0.393	-0.481	0.227
HCO3	0.448	0.57	-0.09	-0.573	0.115
Ca	0.785	0.136	0.461	-0.206	0.271
Eigenvalues	3.33	2.917	1.864	1.473	1.001
% of Variance	27.747	24.306	15.532	12.275	8.34
Cumulative %	27.747	52.052	67.584	79.859	88.199




Figure 5 A dendrogram produced from cluster analysis based on the sampling LGAs to identify the significant groundwater characteristics in Kebbi State.

Group 2 comprises sampling LGAs: Augie and Jega. Jega LGA had the lowest salinity. Several factors such as vegetation, quality of recharge and groundwater level depths, and aridity account for salinity variability in groundwater.^{148–150} Sampling LGAs Fakai and Zuru encompass the last group. Groundwater composition in these LGAs is alkaline. The salinisation/alkalinisation of aquifers is controlled by many factors.^{151–154} Overall, three types of shallow aquifers are noticeable in Kebbi State: those shallow aquifers with higher concentrations of Zn and Cl, salinity and temperature in northern Kebbi State; those shallow aquifers with very low salinity in central Kebbi State; and those shallow aquifers with alkaline waters in southern Kebbi State.

The implication for groundwater management

This study showed that shallow groundwater in Kebbi State is fit for human consumption. Rock weathering appeared to be the primary mechanism controlling the hydrochemistry of groundwater. A multivariate analysis approach can be applied to explain the variability of hydrochemistry of shallow groundwater. It can reveal natural geogenic and anthropogenic alteration of groundwater chemistry that may have severe consequences on humans and the entire ecosystem services that rely on shallow aquifers.¹⁵⁵ Over the last 3-4 decades,

many countries have employed direct and indirect approaches to groundwater management.¹⁵⁶

Increasing human activities with corresponding changes in land use has made groundwater quality management an arduous task.^{157–159} Consequently, multivariate analysis can classify groundwater sources and detect possible pollution from anthropogenic activities.^{160–162} Although accumulating groundwater contaminants are rising in shallow aquifers consequence of human activities, increased demand for shallow groundwater for irrigated farming is making groundwater quality management increasingly tricky.^{163,164} It also requires innovative water quality management approaches lacking in most developing countries. Several pollutant modelling approaches exist for assessing pollution in aquifers. However, most of these approaches are complex and require a large amount of data which may be lacking in places like Kebbi State. While multivariate analysis (MA) can be used to study groundwater pollution, MA cannot quantify the actual risk exposure while using polluted aquifers for domestic, industrial, or agricultural purposes. Therefore, many recommendations for enhancing the quality of groundwater aquifers are needed. These include the application of water quality modelling approaches, the formation of pollution monitoring agencies, policies for ensuring sustainable land use and water resources protection.

Conclusion

The literature is undivided on the significance of water quality and the need to understand the physical and chemical composition of any water meant for drinking, especially in developing countries like Nigeria. This study has reported our recent efforts to represent better the status of rural water quality from remote communities of Kebbi State. Understanding the hydrochemistry of groundwater from shallow aquifers is essential for the success of improved water supply programs which are in progress in many parts of the state. Findings from this study lead to the following remark:

- There is water of good quality from rural Kebbi State. Thus, any program designed to improve the living conditions of rural communities in the state must include improved sanitation and proper land use practice to prevent contamination of shallow groundwater;
- Correlation and Factor analyses indicated that the shallow aquifers are primarily influenced by rock weathering; there is little evidence of anthropogenic input;
- The HCA categorised water sources into three groups, all of which are controlled by natural geogenic processes;

The shallow aquifers contained water of good quality based on the analysed parameters. This finding is helpful since it revealed the status of superficial groundwater quality in Kebbi State. Results can be used as future reference material for water quality management. Since there is insignificant proof of shallow groundwater pollution, this finding can be used as a baseline reference material for future analysis of the hydrochemistry of shallow aquifers. It can also be used as a guiding principle for future policy on rural water supply. Correlation analysis, FA and HCA provide simple tools for assessing groundwater. We hope that this paper will stimulate other scholars to use a comparable method in a future water quality study.

Acknowledgments

This study was financed by the Tertiary Education Trust Fund (TETFund). We also thank all the anonymous contributors.

Funding

None.

Conflicts of interest

Authors declare that there is no conflict of interest.

References

1. Everard M. Community-based groundwater and ecosystem restoration in semi-arid north Rajasthan (1): Socio-economic progress and lessons for groundwater-dependent areas. *Ecosystem Services*. 2015;16:125–135.
2. Machiwal D, Jha MK, Mal BC. Assessment of groundwater potential in a semi-arid region of India using remote sensing, GIS and MCDM techniques. *Water Resources Management*. 2010;25(5):1359–1386.
3. Page D, Bekele E, Vanderzalm V, et al. Managed Aquifer Recharge (MAR) in sustainable urban water management. *Water*. 2018;10(3):239.
4. Parisi A, Monno V, Fidelibus MD. Cascading vulnerability scenarios in the management of groundwater depletion and salinisation in semi-arid areas. *International Journal of Disaster Risk Reduction*. 2018;30:292–305.
5. Wali SU, Ibrahim MD, Danjuma SA, et al. Reassessing groundwater potentials and subsurface water hydrochemistry in a Tropical Anambra Basin, Southeastern Nigeria. *Journal of Geological Research*. 2020;2(3):1–24.
6. Wali SU, Ibrahim MD, Danjuma SA, et al. Re-examination of hydrochemistry and groundwater potentials of Cross River and Imo-Kwa-Ibo intersecting Tropical Basins of South-South Nigeria. *Journal of Geological Research*. 2020;2(3):25–42.
7. Wali SU, Ibrahim MD, Danjuma SA, et al., Review of groundwater potentials and groundwater hydrochemistry of semi-arid Hadejia-Yobe Basin, North-eastern Nigeria. *Journal of Geological Research*. 2020;2(2):20–33.
8. Carter RC, Morgulis ED, Dottridge J, et al. Groundwater modelling with limited data: a case study in a semi-arid dunefield of northeast Nigeria. *Quarterly Journal of Engineering Geology and Hydrogeology*. 1994;27(Supplement):S85–S94.
9. Talabi AO, Tijani MN. Hydrochemical and stable isotopic characterisation of shallow groundwater system in the crystalline basement terrain of Ekiti area, southwestern Nigeria. *Applied Water Science*. 2013;3(1):229–245.
10. Tarhule A, Woo M. Characteristics and use of shallow wells in a stream fadama: A case study in northern Nigeria. *Applied Geography*. 1997;17(1):29–42.
11. Abubakar IT, Yusuf MA, Olofin EA, et al. Shallow groundwater condition for irrigation along dryland river basin, northwestern Nigeria. *Journal of Dryland Agriculture*. 2018;4(1):1–11.
12. Nyong AO, Kanaroglou PS. A survey of household domestic water-use patterns in rural semi-arid Nigeria. *Journal of Arid Environments*. 2001;49(2):387–400.
13. Peter AA, Adesiji RA, Hassana MI. Appraisal of rural water supply: case study of Kwara State, North Central Nigeria. *International Journal of Basic and Applied Science*. 2013;1(4):816–826.
14. Arwenyo B, Wasswa J, Nyeko M, et al. The impact of septic systems density and nearness to spring water points, on water quality. *African Journal of Environmental Science and Technology*. 2017;11(1):11–18.
15. Edet A. Groundwater chemistry and quality of Nigeria: A status review. *African Journal of Environmental Science and Technology*. 2012;5(13).
16. Olayiwola LM, Adeleye OA. Rural infrastructural development in Nigeria: between 1960 and 1990 – problems and challenges. *Journal of Social Sciences*. 2017;11(2):91–96.
17. Wilbers GJ, Sebesvari Z, Renaud F. Piped-water supplies in rural areas of the Mekong Delta, Vietnam: water quality and household perceptions. *Water*. 2014;6(8):2175–2194.
18. Ajibade LT. Assessment of water quality along with River Asa, Ilorin, Nigeria. *The Environmentalist*. 2004;24:11–18.
19. Onabolu B, Jimoh OD, Igboro SB, et al. Source to point of use drinking water changes and knowledge, attitude and practices in Katsina State, Northern Nigeria. *Physics and Chemistry of the Earth, Parts A/B/C*. 2011;36(14–15):1189–1196.
20. Khair SM, Shahbaz M, Kate RS, et al. Diverse drivers of unsustainable groundwater extraction behaviour operate in an unregulated water-scarce region. *J Environ Manage*. 2019;236:340–350.
21. Minderhoud PSJ, Erkens G, Pham VH, et al. Impacts of 25 years of groundwater extraction on subsidence in the Mekong delta, Vietnam. *Environ Res Lett*. 2017;12(6):064006.
22. Urfels A, Andrew JM, Timothy JK, et al. Drivers of groundwater utilisation in water-limited rice production systems in Nepal. *Water Int*. 2020;45(1):39–59.
23. Alfarrak N, Walraevens K. Groundwater overexploitation and seawater intrusion in coastal areas of arid and semi-arid regions. *Water*. 2018;10(2):143.
24. Thomas R, Duraisamy V. Hydrogeological delineation of groundwater vulnerability to droughts in semi-arid areas of western Ahmednagar district. *The Egyptian Journal of Remote Sensing and Space Science*. 2018;21(2):21–137.
25. Tian Y, Zheng Y, Wu B, et al. Modeling surface water-groundwater interaction in arid and semi-arid regions with intensive agriculture. *Environmental Modelling & Software*. 2015;63:170–184.
26. Goni IB, Sheriff BM, Alhaji MK, et al. Assessment of nitrate concentrations in the shallow groundwater aquifer of Maiduguri and environs, Northeastern Nigeria. *Scientific African*. 2019;4:1–8.
27. Li X, Tang C, Cao Y, et al. A multiple isotope (H, O, N, C and S) approach to elucidate the hydrochemical evolution of shallow groundwater in a rapidly urbanised area of the Pearl River Delta, China. *Sci Total Environ*. 2020;724:137930.
28. Singh G, Rishi MS, Herojeet R, et al. Evaluation of groundwater quality and human health risks from fluoride and nitrate in semi-arid region of northern India. *Environ Geochem Health*. 2019.
29. Jalali M. Salinization of groundwater in arid and semi-arid zones: an example from Tajarak, western Iran. *Environmental Geology*. 2006;52(6):1133–1149.
30. MacDonald AM, Roger CC, David MJM, et al. What impact will climate change have on rural groundwater supplies in Africa? *Hydrological Sciences Journal*. 2009;54(4):690–703.
31. Mkandawire T. Quality of groundwater from shallow wells of selected villages in Blantyre District, Malawi. *Physics and Chemistry of the Earth, Parts A/B/C*. 2008;33(8–13):807–811.
32. Pritchard M, Mkandawire T, O'Neill JG. Assessment of groundwater quality in shallow wells within the southern districts of Malawi. *Physics and Chemistry of the Earth, Parts A/B/C*. 2008;33(8–13):812–823.
33. Biswas A, Nath B, Bhattacharya P, et al. Hydrogeochemical contrast between brown and grey sand aquifers in shallow depth of Bengal Basin: consequences for sustainable drinking water supply. *Sci Total Environ*. 2012;431:402–412.
34. Foster SD, Hirata R, Howard KWF. Groundwater use in developing cities: policy issues arising from current trends. *Hydrogeology Journal*. 2010;19(2):271–274.
35. Okotto L, Okotto-Okotto J, Price H, et al. Socio-economic aspects of domestic groundwater consumption, vending and use in Kisumu, Kenya. *Applied Geography*. 2015;58:189–197.
36. Wu J, Sun Z. Evaluation of shallow groundwater contamination and associated human health risk in an alluvial plain impacted by agricultural and industrial activities, mid-west China. *Exposure and Health*. 2015;8(3):311–329.

37. Bucci A, Barbero D, Lasagna M, et al. Shallow groundwater temperature in the Turin area (NW Italy): vertical distribution and anthropogenic effects. *Environmental Earth Sciences*. 2017;76(5).
38. Chen S, Jiao XC, Gai N, et al. Perfluorinated compounds in soil, surface water, and groundwater from rural areas in eastern China. *Environ Pollut*. 2016;211:124–131.
39. Gowing J, Parkin G, Forsythe N, et al. Shallow groundwater in sub-Saharan Africa: neglected opportunity for sustainable intensification of small-scale agriculture? *Hydrology and Earth System Sciences Discussions*. 2016:1–33.
40. Nag SK, Ghosh P. Delineation of groundwater potential zone in Chhatna Block, Bankura District, West Bengal, India using remote sensing and GIS techniques. *Environmental Earth Sciences*. 2012;70(5):2115–2127.
41. Moyo NAG. An analysis of the chemical and microbiological quality of groundwater from boreholes and shallow wells in Zimbabwe. *Physics and Chemistry of the Earth, Parts A/B/C*. 2013;66:27–32.
42. Oyelami AC, Aladejana JA, Agbede OO. Assessment of the impact of open waste dumpsites on groundwater quality: a case study of the Onibu–Eja Dumpsite, Southwestern Nigeria. *Procedia Earth and Planetary Science*. 2013;7:648–651.
43. Guo H, Zhang B, Wang G, et al. Geochemical controls on arsenic and rare earth elements approximately along a groundwater flow path in the shallow aquifer of the Hetao Basin, Inner Mongolia. *Chemical Geology*. 2010;270(1–4):117–125.
44. Long DT, Voice TC, Niagolova ND, et al. Effects of human activities on karst groundwater geochemistry in a rural area in the Balkans. *Applied Geochemistry*. 2012;27(10):1920–1931.
45. Marghade D, Malpe DB, Zade AB. Major ion chemistry of shallow groundwater of a fast-growing city of central India. *Environ Monit Assess*. 2012;184(4):2405–2418.
46. Pang Z, Yuan L, Huang T, et al. Impacts of human activities on the occurrence of groundwater nitrate in an alluvial plain: A multiple isotopic tracers approach. *Journal of Earth Science*. 2013;24(1):111–124.
47. Chen S, Wu W, Hu K, et al. The effects of land-use change and irrigation water resource on nitrate contamination in shallow groundwater at county scale. *Ecological Complexity*. 2010;7(2):131–138.
48. Gao Y, Luo C, Zhou P, et al. Groundwater nitrogen pollution and assessment of its health risks: a case study of a typical village in rural-urban continuum, China. *PLoS One*. 2012;7(4):e33982.
49. Rajasooriyar LD, Boelee E, Hiscock KM, et al. Mapping the potential human health implications of groundwater pollution in southern Sri Lanka. *Water Resources and Rural Development*. 2013;1–2:27–42.
50. Wongsasulak P, Chotpanarat S, Siri Wong W, et al. Heavy metal contamination and human health risk assessment in drinking water from shallow groundwater wells in an agricultural area in Ubon Ratchathani province, Thailand. *Environ Geochem Health*. 2014;36(1):169–182.
51. Kumar P, Mahajan AK, Kumar A. Groundwater geochemical facie: implications of rock-water interaction at the Chamba city (HP), northwest Himalaya, India. *Environ Sci Pollut Res Int*. 2020;27(9):9012–9026.
52. Hu K, Huang Y, Li H, et al., Spatial variability of shallow groundwater level, electrical conductivity and nitrate concentration, and risk assessment of nitrate contamination in North China Plain. *Environment International*. 2005;31(6):896–903.
53. Wali SU, Kabiru JU, Danjuma SA, et al. Hydrochemical characterisation of shallow and deep groundwater in Basement Complex areas of southern Kebbi State, Sokoto Basin, Nigeria. *Applied Water Science*. 2019;9(169):1–36.
54. Wang P, Pozdniakov SP, Grinevsky SO, et al., Shallow groundwater dynamics and its driving forces in extremely arid areas: a case study of the lower Heihe River in northwestern China. *Hydrological Processes*. 2014;28(3):1539–1553.
55. Wu GH, Cao SS. Mercury and cadmium contamination of irrigation water, sediment, soil and shallow groundwater in a wastewater-irrigated field in Tianjin, China. *Bull Environ Contam Toxicol*. 2010;84(3):336–341.
56. Wali SU, Adetunji MT, Gbadebo AM, et al. An examination of rural groundwater quality in southwestern Zuru Local Government Area, Kebbi State Nigeria. *The Nigerian Geographical Journal*. 2018;12(1):122–142.
57. NSDWQ. Nigerian standard for drinking water quality: Nigerian Industrial Standard NIS 554. Standards Organisation of Nigeria, 2007. 30 p.
58. Wali SU, Danjuma SA, Ibrahim MD, et al. Distribution of public water supply: Analysis of population density and water supply in Kebbi State, Northwestern Nigeria. *European Journal of Social Sciences Studies*. 2018;3(4):36–55.
59. Anderson HR, Ogilbee W. Aquifers in the Sokoto Basin, Northwestern Nigeria, With a description of the general hydrogeology of the region: contributions to the hydrology of Africa and the Mediterranean Region. *Geological Survey Water-Supply Paper*. 1973:1–88.
60. Emujakorue G, Ofoha CC, Kiani I. Investigation into the basement morphology and tectonic lineament using aeromagnetic anomalies of Parts of Sokoto Basin, North Western, Nigeria. *Egyptian Journal of Petroleum*. 2018;27(4):671–681.
61. Kogbe CA. Cretaceous and tertiary of the Iullemeden basin in Nigeria (West Africa). *Cretaceous Research*. 1981;2:129–186.
62. Paul II, Bayode EN. Watershed characteristics and their implication for hydrologic response in the Upper Sokoto Basin, Nigeria. *Journal of Geography and Geology*. 2012;4(2):147–155.
63. Otti VI. Developing a sustainable water supply and sanitation programme management in the rural areas of Nigeria. *International Journal of Engineering and Technology*. 2012;2(6):1046–1051.
64. Offodile ME. Groundwater study and development in Nigeria. Mecon Geological and Engineering, Ltd Ehinder O, 2nd edn. Jos, Nigeria; 2002.
65. Wali SU, Jega UK, Danjuma AS, et al. Hydrochemical characterisation of shallow and deep groundwater in Basement Complex areas of southern Kebbi State, Sokoto Basin, Nigeria. *Applied Water Science*. 2019;9(169):1–36.
66. Lin CY, Abdullah MH, Praveena SM, et al. Delineation of temporal variability and governing factors influencing the spatial variability of shallow groundwater chemistry in a tropical sedimentary island. *Journal of Hydrology*. 2012;432–433:26–42.
67. Azhar SC, Aris AH, Yusoff MK, et al. Classification of river water quality using multivariate analysis. *Procedia Environmental Sciences*. 2015;30:79–84.
68. Blake S, Henry T, Murray J, et al., Compositional multivariate statistical analysis of thermal groundwater provenance: A hydrogeochemical case study from Ireland. *Applied Geochemistry*. 2016;75:171–188.
69. Rakotondrabe F, Ngoupayou JRN, Mfonka Z, et al. Water quality assessment in the Betare-Oya gold mining area (East-Cameroon): Multivariate Statistical Analysis approach. *Science of the Total Environment*. 2018;610–611:831–844.
70. Yidana SM. Evolutionary analysis of groundwater flow: Application of multivariate statistical analysis to hydrochemical data in the Densu Basin, Ghana. *Journal of African Earth Sciences*. 2018;138:167–176.
71. Adomako D, Gibrilla A, Akiti TT, et al. Hydrogeochemical Evolution and Groundwater Flow in the Densu River Basin, Ghana. *Journal of Water Resource and Protection*. 2011;03(07):548–561.
72. Devic G, Djordjevic D, Sakan S. Natural and anthropogenic factors affecting the groundwater quality in Serbia. *Sci Total Environ*. 2014;468–469:933–942.

73. Tran DA, Tsujimura M, Vo LP, et al. Hydrogeochemical characteristics of a multi-layered coastal aquifer system in the Mekong Delta, Vietnam. *Environmental Geochemistry and Health*. 2020;42(2):661–680.
74. Appelo CAJ. Cation and proton exchange, pH variations, and carbonate reactions in a freshening aquifer. *Water Resources Research*. 1994;30(10):2793–280.
75. Nagaraju A, Sunil Kumar K, Thejaswi A. Assessment of groundwater quality for irrigation: a case study from Bandalamottu lead mining area, Guntur District, Andhra Pradesh, South India. *Applied Water Science*. 2014;4(4):385–396.
76. Krainov SR, Belousova AP, Ryzhenko BN. Formation of high-alkali groundwater in upper aquifers. *Water Resources and Industry*. 2001;28(5):491–501.
77. Lee JY, Moon SH, Yi MJ, et al. Groundwater contamination with petroleum hydrocarbons, chlorinated solvents and high pH: implications for multiple sources. *Quarterly Journal of Engineering Geology and Hydrogeology*. 2008;41:35–47.
78. Clohessy S, Appleyard S, Vogwill R. Groundwater acidification near the water table of the Superficial aquifer, Gngangara Mound, Swan Coastal Plain, Western Australia. *Applied Geochemistry*. 2013;36:140–152.
79. Shi WM, Yao J, Yan F. Vegetable cultivation under greenhouse conditions leads to rapid accumulation of nutrients, acidification and salinity of soils and groundwater contamination in South-Eastern China. *Nutrient Cycling in Agroecosystems*. 2008;83(1):73–84.
80. Takem GE, Kuitcha D, Ako AA, et al. Acidification of shallow groundwater in the unconfined sandy aquifer of the city of Douala, Cameroon, Western Africa: implications for groundwater quality and use. *Environmental Earth Sciences*. 2015;74(9):6831–6846.
81. EPA. Parameters of water quality: Interpretation and Standards. An Ghniomhaireacht um Chaomhnu Comhshaoil. Ireland; 2001.
82. Hayashi M. Temperature–electrical conductivity relation of water for environmental monitoring and geophysical data inversion. *Environmental Monitoring and Assessment*. 2004;96:119–128.
83. Oyem HH, Oyem IM, Ezeweali D. Temperature, pH, electrical conductivity, total dissolved solids and chemical oxygen demand of groundwater in Boji–Boji/Owa area and immediate environs suburbs. *Research Journal of Environmental Sciences*. 2014;8(8):444–450.
84. Ayers JC, Goodbred S, George G, et al. Sources of salinity and arsenic in groundwater in southwest Bangladesh. *Geochem Trans*. 2016;17:4.
85. Chien NP, Lautz LK. Discriminant analysis as a decision–making tool for geochemically fingerprinting sources of groundwater salinity. *Sci Total Environ*. 2018;618:379–387.
86. Ghabayen SMS, McKee M, Kembrowski M. Ionic and isotopic ratios for identifying salinity sources and missing data in the Gaza aquifer. *Journal of Hydrology*. 2006;318(1–4):360–373.
87. Gopinath S, Srinivasamoorthy K, Vasanthavigar M, et al. Hydrochemical characteristics and groundwater salinity in parts of Nagapattinam district of Tamil Nadu and the Union Territory of Puduchery, India. *Carbonates and Evaporites*. 2016;33(1):1–13.
88. Jia Y, Guo H, Xi B, et al. Sources of groundwater salinity and potential impact on arsenic mobility in the western Hetao Basin, Inner Mongolia. *Sci Total Environ*. 2017;601–602.
89. Vengosh A, Spivack AJ, Artzi Y, et al. Geochemical and boron, strontium, and oxygen isotopic constraints on the origin of the salinity in groundwater from the Mediterranean Coast of Israel. *Water Resources Research*. 1999;35(6):1877–1894.
90. Douglas SH, Dixon B, Griffin D. Assessing intrinsic and specific vulnerability models ability to indicate groundwater vulnerability to groups of similar pesticides: a comparative study. *Physical Geography*. 2018:1–19.
91. Li XD, Liu CQ, Harue M, et al. The use of environmental isotopic (C, Sr, S) and hydrochemical tracers to characterise anthropogenic effects on karst groundwater quality: A case study of the Shuicheng Basin, SW China. *Applied Geochemistry*. 2010;25(12):1924–1936.
92. Savci S. Investigation of effect of chemical fertilizers on environment. *APCBEE Procedia*. 2012;1:287–292.
93. Oyem HH, Oyem M, Ezeweali D. Temperature, pH, electrical conductivity, total dissolved solids and biological oxygen demand of groundwater in Boji–Boji/Agbor Owa area and immediate suburb. *Research Journal of Environmental Sciences*. 2014;8(8):444–450.
94. Akhtar MM, Tang Z. Identification of contamination sources and TDS concentration in groundwater of second biggest city of Pakistan. *International Journal of Environmental Science and Development*. 2013:341–345.
95. Adelana SMA, Olasehinde PI, Vrbka P. Groundwater recharge in the cretaceous and tertiary sediment aquifers of northwestern Nigeria, using hydrochemical and isotopic techniques. *Groundwater and human development*. 2002:907–915.
96. Adelana SMA, Olasehinde PI, Vrbka P. Isotope and geochemical characterisation of surface and subsurface waters in the Semi-arid Sokoto basin, Nigeria. *African Journal of Science and Technology (AJST) Science and Engineering Series*. 2008;4(2):80–89.
97. WHO. Guidelines for drinking-water quality: Fourth edition incorporating the first addendum. WHO Library Cataloguing–in–Publication Data. Geneva: World Health Organization; 2018. 631 p.
98. Haddad–Martim PM, Souza Filho CRd, Carranza EJM. Spatial analysis of mineral deposit distribution: A review of methods and implications for structural controls on iron oxide–copper–gold mineralisation in Carajás, Brazil. *Ore Geology Reviews*. 2017;81:230–244.
99. Kanazawa Y, Kamitani M. Rare earth minerals and resources in the world. *Journal of Alloys and Compounds*. 2006:408–412.
100. Nordstrom DK, Alpers CN, Negative pH, efflorescent mineralogy, and consequences for environmental restoration at the Iron Mountain Superfund site, California. *Proceedings of National Academy of Science*. 1999;96:3455–3462.
101. Pourhosseini SEM, Norouzi O, Salimi P, et al. Synthesis of a novel interconnected 3D pore network algal biochar constituting iron nanoparticles derived from a harmful marine biomass as high–performance asymmetric supercapacitor electrodes. *ACS Sustainable Chemistry & Engineering*. 2018;6(4):4746–4758.
102. Zhu X, Tian S, Cai Z. Toxicity assessment of iron oxide nanoparticles in zebrafish (*Danio rerio*) early life stages. *PLoS One*. 2012;7(9):e46286.
103. Yoder HS. Potassium–rich rocks: Phase analysis and heteromorphic relations. *Journal of Petrology*. 1986;27(5):1215–1228.
104. Sun Y, Teng FZ, Ying JF, et al. Magnesium isotopic evidence for ancient subducted oceanic crust in LOMU–like potassium–rich volcanic rocks. *Journal of Geophysical Research: Solid Earth*. 2017;122(10):7562–7572.
105. Zhang Y, Zeng Z, Li X, et al. High–potassium volcanic rocks from the Okinawa Trough: Implications for a cryptic potassium–rich and DUPAL–like source. *Geological Journal*. 2018;53(5):1755–1766.
106. Chapman PJ, Reynolds B, Wheatler HS. Sources and controls of calcium and magnesium in storm runoff: the role of groundwater and ion exchange reactions along water flowpaths. *Hydrology and Earth System Sciences*. 1997;1(3):671–685.
107. Magaritz M, Kafri U. Concentration of magnesium in carbonate nodules of soils: an indication of fresh groundwater contamination by intruding seawater. *Chemical Geology*. 1979;27:143–155.
108. Rapant S, Cvečková V, Fajčíková K, et al. Impact of calcium and magnesium in groundwater and drinking water on the health of inhabitants of the Slovak Republic. *International Journal of Environmental Research and Public Health*. 2017;14(278):1–22.

109. Barnes I. Geochemistry of highly basic calcium hydroxide groundwater in Jordan. *Chemical Geology*. 1982;35:147–154.
110. Gu B, Brooks, Scott C, Roh Y, et al. Geochemical reactions and dynamics during titration of a contaminated groundwater with high uranium, aluminum, and calcium. *Geochimica et Cosmochimica Acta*. 2003;67(15):2749–2761.
111. Schot PP, Wassen MJ. Calcium concentrations in wetland groundwater in relation to water sources and soil conditions in the recharge area. *Journal of hydrology*. 1993;141:197–217.
112. Sharma CM, Kang S, Tripathee L, et al. Major ions and irrigation water quality assessment of the Nepalese Himalayan rivers. *Environment, Development and Sustainability*. 2020:1–13.
113. Singh S, Raju NJ, Ramakrishna C. Evaluation of groundwater quality and its suitability for domestic and irrigation use in parts of the Chandauli–Varanasi Region, Uttar Pradesh, India. *Journal of Water Resource and Protection*. 2015;07(07):572–587.
114. Varlagas H, Savvas D, Mouzakis G, et al., Modelling uptake of Na⁺ and Cl⁻ by tomato in closed-cycle cultivation systems as influenced by irrigation water salinity. *Agricultural Water Management*. 2010;97(9):1242–1250.
115. Rokicki CA, Boyer TH. Bicarbonate–form anion exchange: affinity, regeneration, and stoichiometry. *Water Res*. 2011;45(3):1329–1337.
116. Alfa–Sika Mande SL, Mingzhu L, Ibrahim T, et al. Water–rock interaction effect on evolution of total hardness in groundwater in urban. *American Journal of Water Resources*. 2018;6(2):48–52.
117. Lo IM, Lam CS, Lai KC. Hardness and carbonate effects on the reactivity of zero–valent iron for Cr(VI) removal. *Water Research*. 2006;40(3):595–605.
118. Marghade D, Malpe DB, Zade AB. Geochemical characterisation of groundwater from northeastern part of Nagpur urban, Central India. *Environmental Earth Sciences*. 2011;62(7):1419–1430.
119. Debernardi L, De Luca DA, Lasagna M. Correlation between nitrate concentration in groundwater and parameters affecting aquifer intrinsic vulnerability. *Environmental Geology*. 2007;55(3):539–558.
120. Katsoyiannis IA, Stephan JH, Adrian A, et al. Arsenic speciation and uranium concentrations in drinking water supply wells in Northern Greece: correlations with redox indicative parameters and implications for groundwater treatment. *Sci Total Environ*. 2007;383(1–3):128–140.
121. Mrazovac S, Vojinović–Miloradov M. Correlation of main physicochemical parameters of some groundwater in northern Serbia. *Journal of Geochemical Exploration*. 2011;108(3):176–182.
122. Koc C. The environmental effects of salinity load in Great Menderes Basin irrigation schemes. *Environ Monit Assess*. 2008;146(1–3): 479–489.
123. Rhoades JD. Electrical conductivity methods for measuring and mapping soil salinity. 1993;49:201–251.
124. Dupraz S. Experimental and numerical modelling of bacterially induced pH increase and calcite precipitation in saline aquifers. *Chemical Geology*. 2009;265(1–2):44–53.
125. Fisher RS, Mullican F. Hydrochemical evolution of sodium–sulfate and sodium–chloride groundwater beneath the Northern Chihuahuan Desert, Trans–Pecos, Texas, USA. *Hydrogeology Journal*. 1997;5(2):4–16.
126. Roy JW, McInnis R, Bickerton G, et al., Assessing potential toxicity of chloride–affected groundwater discharging to an urban stream using juvenile freshwater mussels (*Lampsilis siliquoidea*). *Sci Total Environ*. 2015;532:309–315.
127. Trainer FW, Heath RC. Bicarbonate content of groundwater in carbonate rock in Eastern North America. *Journal of Hydrology*. 1976;31:37–55.
128. Anawar HM, Akai J, Sakugawa H. Mobilisation of arsenic from subsurface sediments by effect of bicarbonate ions in groundwater. *Chemosphere*. 2004;54(6):753–762.
129. Chae GT, Yun ST, Kim K, et al. Hydrogeochemistry of sodium–bicarbonate type bedrock groundwater in the Pocheon spa area, South Korea: water–rock interaction and hydrologic mixing. *Journal of Hydrology*. 2006;321(1–4):326–343.
130. Bencer S, Boudoukha A, Mouni L. Multivariate statistical analysis of the groundwater of Ain Djacer area (Eastern of Algeria). *Arabian Journal of Geosciences*. 2016;9(248):1–10.
131. Belkhirli N, Narany TS. Using multivariate statistical analysis, geostatistical techniques and structural equation modelling to identify spatial variability of groundwater quality. *Water Resources Management*. 2015;29:2073–2089.
132. Egbueri JC. Water quality appraisal of selected farm provinces using integrated hydrogeochemical, multivariate statistical, and microbiological technique. *Modeling Earth Systems and Environment*. 2019;5(3):997–1013.
133. Patil GR, Basu R, Rashidi TH. Mode choice modeling using adaptive data collection for different trip purposes in Mumbai Metropolitan Region. *Transportation in Developing Economies*. 2020;6(1).
134. Adams S, Titus R, Pietersen K, et al. Hydrochemical characteristics of aquifers near Sutherland in the Western Karoo, South Africa. *Journal of Hydrology*. 2001;241:91–103.
135. Kazakis N, Mattas C, Pavlou A, et al. Multivariate statistical analysis for the assessment of groundwater quality under different hydrogeological regimes. *Environmental Earth Sciences*. 2017;76(9):1–13.
136. Keul N, Langer G, Thoms S, et al. Exploring foraminiferal Sr/Ca as a new carbonate system proxy. *Geochimica et Cosmochimica Acta*. 2017;202:374–386.
137. Lambkin DC, Gwilliam KH, Layton C, et al. Soil pH governs production rate of calcium carbonate secreted by the earthworm *Lumbricus Terrestris*. *Applied Geochemistry*. 2011;26:S64–S66.
138. Tabelin CB, Igarashi T, Mylah VT, et al. Arsenic, selenium, boron, lead, cadmium, copper, and zinc in naturally contaminated rocks: A review of their sources, modes of enrichment, mechanisms of release, and mitigation strategies. *Sci Total Environ*. 2018;645:1522–1553.
139. Mukate S, Panaskar D, Wagh V, et al. Impact of anthropogenic inputs on water quality in Chincholi industrial area of Solapur, Maharashtra, India. *Groundwater for Sustainable Development*. 2018;7:359–371.
140. Subba Rao N. Seasonal variation of groundwater quality in a part of Guntur District, Andhra Pradesh, India. *Environmental Geology*. 2005;49(3):413–429.
141. Haritash AK, Kaushik CP, Kaushik A, et al. Suitability assessment of groundwater for drinking, irrigation and industrial use in some North Indian villages. *Environmental Monitoring Assessment*. 2008;145(1–3):397–406.
142. Missi C, Atekwana EA. Physical, chemical and isotopic characteristics of groundwater and surface water in the Lake Chilwa Basin, Malawi. *Journal of African Earth Sciences*. 2020;162:103737.
143. Scheelbeek PF, Aneire EK, Sontosh M, et al. Drinking water sodium and elevated blood pressure of healthy pregnant women in salinity–affected coastal areas. *Hypertension*. 2016;68(2):464–470.
144. Selvakumar S, Chandrasekar N, Kumar G. Hydrogeochemical characteristics and groundwater contamination in the rapid urban development areas of Coimbatore, India. *Water Resources and Industry*. 2017;17:26–33.
145. Smith LED, Siciliano G. A comprehensive review of constraints to improved management of fertilisers in China and mitigation of diffuse water pollution from agriculture. *Agriculture, Ecosystems & Environment*. 2015;209:15–25.

146. Calvache ML, Duque C, Gomez Fontalva JM, et al. Processes affecting groundwater temperature patterns in a coastal aquifer. *International Journal of Environmental Science and Technology*. 2011;8(2):223–236.
147. Morales I, Villanueva-Estrada RE, Rodríguez R, et al. Geological, hydrogeological, and geothermal factors associated to the origin of arsenic, fluoride, and groundwater temperature in a volcanic environment “El Bajío Guanajuatense”, Mexico. *Environmental Earth Sciences*. 2015;74(6):5403–5415.
148. Mcfarlane DJ, George RJ. Factors affecting dryland salinity in two Wheatbelt catchments in Western Australia. *Australian Journal of Science*. 1992;30:85–100.
149. Jolly ID, McEwan KL, Holland KL. A review of groundwater–surface water interactions in arid/semi-arid wetlands and the consequences of salinity for wetland ecology. *Ecohydrology*. 2008;1(1):43–58.
150. Wu J, Li P, Qian H, et al. Assessment of soil salinisation based on a low-cost method and its influencing factors in a semi-arid agricultural area, northwest China. *Environmental Earth Sciences*. 2013;71(8):3465–3475.
151. Argamasilla M, Barbera JA, Andreo B. Factors controlling groundwater salinisation and hydrogeochemical processes in coastal aquifers from southern Spain. *Sci Total Environ*. 2017;580:50–68.
152. Bowen BB, Benison KC. Geochemical characteristics of naturally acid and alkaline saline lakes in southern Western Australia. *Applied Geochemistry*. 2009;24(2):268–284.
153. Xia J, Zhao X, Ren J, et al., Photosynthetic and water physiological characteristics of *Tamarix Chinensis* under different groundwater salinity conditions. *Environmental and Experimental Botany*. 2017;138:173–183.
154. Yu R, Liu T, Xu T, et al. Analysis of salinisation dynamics by remote sensing in Hetao Irrigation District of North China. *Agricultural Water Management*. 2010;97(12):1952–1960.
155. Qureshi AS, McCornick PG, Sarwar A, et al. Challenges and Prospects of Sustainable Groundwater Management in the Indus Basin, Pakistan. *Water Resources Management*. 2009;24(8):1551–1569.
156. Aguilera H, Luis M, Jiménez-Hernández ME, et al. Management implications inferred from the multivariate analysis of vadose zone chemical variables in Las Tablas de Daimiel National Park (Spain). *Geoderma*. 2011;162(3–4):365–377.
157. Butler JRA, Wong GY, Metcalfe DJ, et al. An analysis of trade-offs between multiple ecosystem services and stakeholders linked to land use and water quality management in the Great Barrier Reef, Australia. *Agriculture, Ecosystems & Environment*. 2013;180:176–191.
158. Khan HH, Khan A, Ahmed S, et al. GIS-based impact assessment of land-use changes on groundwater quality: study from a rapidly urbanising region of South India. *Environmental Earth Sciences*. 2010;63(6):1289–1302.
159. Vrščaj B, Poggio L, Marsan FA. A method for soil environmental quality evaluation for management and planning in urban areas. *Landscape and Urban Planning*. 2008;88(2–4):81–94.
160. Güler C, Mehmet AK, Musa A, et al. Assessment of the impact of anthropogenic activities on the groundwater hydrology and chemistry in Tarsus coastal plain (Mersin, SE Turkey) using fuzzy clustering, multivariate statistics and GIS techniques. *Journal of Hydrology*. 2012:414–415.
161. Kazi TG, Arain MB, Jamali MK, et al., Assessment of water quality of polluted lake using multivariate statistical techniques: a case study. *Ecotoxicol Environ Saf*. 2009;72(2):301–309.
162. Mico C, Recatalá L, Peris M, et al. Assessing heavy metal sources in agricultural soils of an European Mediterranean area by multivariate analysis. *Chemosphere*. 2006;65(5):863–872.
163. Wichelns D, Qadir M. Achieving sustainable irrigation requires effective management of salts, soil salinity, and shallow groundwater. *Agricultural Water Management*. 2015;157:31–38.
164. Li P, Tian R, Xue C, et al. Progress, opportunities, and key fields for groundwater quality research under the impacts of human activities in China with a special focus on western China. *Environ Sci Pollut Res Int*. 2017;24(15):13224–13234.