

Massive Hemothorax after Posterior Correction Surgery of Adolescent Idiopathic Scoliosis: A Case Report

Abstract

Hemothorax is a rare complication of postoperative correction of adolescent idiopathic scoliosis (AIS), with an incidence of 0,1%. It is mostly related to thoracoplasty or misplaced pedicle screws. The authors describe the clinical case of a 15-year-old girl who developed a massive hemothorax after AIS surgical correction and fusion.

Fifteen-year-old girl with a Lenke type 6BN scoliosis who underwent posterior correction and fusion surgery with selective pedicle screw construct between D5 and L4 was carried without complications. Twenty-four hours after the procedure, the patient developed pleuritic chest pain and type 1 respiratory failure with hypoxemia (69 mmHg). The thoracic roentgenogram showed a left-sided pleural effusion, confirmed by the CT scan. Neither CT nor angio-CT imaging showed an evident haemorrhagic source or misplacement of pedicle screws.

A chest tube was inserted with immediate drainage of 800 mL of blood and 100 mL more during the following 12 hours. In this case, the complication seems to have been due to the use of a Gelpi retractor with invasion of the intercostal space. Alternatively, an intercostal artery might have been injured during the procedure, going unnoticed by the surgical team. Even though this is a very rare complication, it is important to be aware of its existence, not only to perform a quick diagnosis, but also to avoid it.

Keywords: Scoliosis; Hemothorax; Surgery; Roentgenogram; Thoracic; CT; Pleuritic

Case Report

Volume 5 Issue 6 - 2016

Rita Grazina^{1*}, Renato Andrade², José Marinhos¹, Filipe Santos¹, André Costa¹, Rolando Freitas¹ and Rui Rocha^{1,2}

¹Department of Orthopaedic Surgery, University of Porto, Portugal

²Clínica do Dragão, Espregueira-Mendes Sports Centre - FIFA Medical Centre of Excellence, Portugal

***Corresponding author:** Rita Grazina, Department of Orthopaedic Surgery at Centro Hospitalar de Vila Nova de Gaia/Espinho E.P.E. Portugal; Rua Francisco Sá Carneiro, s/n, 4400-129 Vila Nova de Gaia, Portugal, Tel: 00351 910900425; Email: rita.grazina@gmail.com

Received: September 16, 2016 | **Published:** October 03, 2016

Abbreviations: AIS: Adolescent Idiopathic Scoliosis; CT: Computer Tomography

Introduction

Posterior spine correction and fusion for AIS is considered a safe procedure and is currently the gold-standard in the treatment of progressive curves. Nevertheless, both neurologic and non-neurologic complications may occur [1,2]. Lung complications are rare, being reported in 0.9% of cases. In this sense, hemothorax complications occur in only 0.1% of surgical procedures [3-5]. Hemothorax typically results from medical procedures, such as central venous line insertion and thoracentesis. When occurring in a postoperative context, it is usually related to thoracoplasty and misplaced pedicle screws [1,5].

Case Presentation

A 15-year-old girl was referred to our orthopaedic department due to spine deformity and lumbar pain. Physical examination showed a right rib hump and shoulder asymmetry. The radiographic study showed a curve type 6BN of Lenke, with a left lumbar curve centred in L2 and a Cobb angle of 43° and a right thoracic curve centred in T8 and a Cobb angle of 25°. The patient was proposed for surgery, undergoing a posterior correction and fusion surgery with a pedicle screw selective construct between

D5 and L4. During the procedure there was no evidence of pedicle screw misplacement.

A rod was placed on the concave side of the main curve and, along with a reduction maneuver allowed for partial curve correction. The same procedure was carried with the contralateral rod with subsequent derotation cluster maneuver [1,3]. The patient remained hemodynamically stable during the whole procedure and the postoperative thoracic roentgenogram lacked pleural or parenchymal alterations. Nevertheless, due to intraoperative blood loss, two units of red blood cells were transfused, achieving postoperative haemoglobin of 12.5 g/dL.

Twenty-four hours after the surgery the patient started to complain of thoracic pain with pleuritic characteristics. A blood gas analysis without oxygen supply was carried showing a hypoxemia with an arterial partial pressure of oxygen of 69 mmHg and a two-unit drop on haemoglobin. A new thoracic roentgenogram was carried and showed a massive hemothorax, which was confirmed by CT imaging. Angio-CT imaging allowed excluding pulmonary thromboembolism. Misplacement of pedicle screws was also excluded by the original CT scan (Figure 1&2).

A chest tube was inserted with immediate drainage of 800 mL of blood. During the following 12 hours, 100 mL more were drained. Another CT scan was performed in order to confirm

the position of the chest tube. All the clinical and analytical manifestations were solved after the chest tube insertion (Figure 3&4).




Figure 1: AIS: coronal and axial radiographs.



Figure 2: On the left: thoracic roentgenogram showing a left white lung. On the right: CT image showing massive hemothorax.




Figure 3: CT scan imaging obtained after thoracic drainage. On the right, it is possible to observe the chest tube.

Discussion

Pulmonary complications after AIS correction surgery are rare. A study with 702 AIS patients reported only 10 respiratory complications. These complications do not seem to be related to factors like Lenke curve type, region of the major curve or even the number of instrumented levels. Among those complications,

hemothorax is a quite rare one. The most commonly described causes of hemothorax after posterior correction and fusion surgery are thoracoplasty and misplaced pedicle screws [1,5]. Pedicle screw fixation is considered as gold standard procedure in which concerns the spinal deformity correction [2,6] (Figure 1).




Figure 4: Post-operative radiographs.

According to Hicks et al. [2], the risk of complications is higher when approaching the thoracic spine, as the pedicles are smaller. Moreover, thoracolumbar vertebral bodies lie very closely to visceral organs. The most commonly reported complication is, actually, screw misplacement, whose real incidence might be around 15%, confirmed by CT scan. Studies that use radiographies alone seem to underestimate this incidence [2,7].

Not only the malposition but also an inadequate length of the screw may cause damage to the structures nearby, as a long screw might cause damage to the oesophagus, pleura or lung [5]. However, the need for reoperations due misplacement is only 0.6%, allowing to conclude that, most of the times, screw malposition does not warrant damage to the surrounding structures. As a matter of fact, according to Gautschi et al. [5]. Review, only two visceral complications were reported after the placement of 35 630 pedicle screws in 5654 patients. In this specific case, thoracoplasty has not been performed and imaging studies failed to confirm any misplacement of the screws. A plausible source for the haemorrhage might be the use of a Gelpi retractor.

The Gelpi retractor is a helpful tool that allows a better surgical exposure at the same time that limits the retraction-associated damage to the paravertebral muscles, but its sharp tips may be associated with pleural damage in this case [1]. Another hypothesis to take into account is the possible injury to an intercostal artery during instrumentation that might have gone unnoticed during the procedure. Intercostal arteries supply the posterior part of the pleurae and they run posteriorly along the ribs. As imaging excludes a conflict with the screws, a lesion to one of those vessels might have happened during probing [3].

Surgical correction of scoliosis is a major procedure, so every attempt should be done in order to minimize post-operative complications. The knowledge of the possible complications, even the rarest ones, allows an earlier diagnosis and treatment with a consequent better outcome for the patient. This specific case alerts for the possibility of a rare but potentially lethal complication, whose early recognition is of vital importance.

Conclusion

Although rare, hemothorax is an important and severe complication that can arise from posterior spine instrumentation. Surgeons must be aware of its existence in order to recognize and treat it in time and, above all, to avoid it. This case report shows the need for a multidisciplinary approach in the treatment of such complications.

References

1. Pang L, Watanabe K, Toyama Y, Matsumoto M (2014) Massive hemothorax caused by Gelpi retractor during posterior correction surgery for adolescent idiopathic scoliosis: a case report. *Scoliosis* 9: 17.
2. Hicks JM, Singla A, Shen FH, Arlet V (2010) Complications of Pedicle Screw Fixation in Scoliosis Surgery: A Systematic Review. *Spine* 35(11): E465-E470.
3. Ogura Y, Watanabe K, Hosogane N, Toyama Y, Matsumoto M (2013) Acute respiratory failure due to hemothorax after posterior correction surgery for adolescent idiopathic scoliosis: a case report. *BMC Musculoskelet Disord* 14: 132.
4. Fu KM, Smith JS, Polly DW, Ames CP, Berven SH, et al. (2011) Morbidity and mortality associated with spinal surgery in children: a review of the scoliosis research society morbidity and mortality database. *J Neurosurg Pediatr* 7(1): 37-41.
5. Gautschi OP, Schatlo B, Schaller K, Tessitore EE (2011) Clinically relevant complications related to pedicle screw placement in thoracolumbar surgery and their management: a literature review of 35,630 pedicle screws. *Neurosurg Focus* 31(4): E8.s
6. Lenke LG, Kuklo TR, Ondra S, Poly D (2008) Rationale behind the current state-of-the-art treatment of scoliosis. *Spine* 33(10): 1051-1054.
7. Ruf M, Harms J (2002) Pedicle screws in 1- and 2-year-old children: technique, complications, and effect on further growth. *Spine* 27(21): E460-E466.