

Comparison of Clinical Outcomes between Different Femoral Tunnel Positions after Anterior Cruciate Ligament Reconstruction Surgery

Research Article

Abstract

There is no robust consensus on whether the more oblique femoral tunnel position offers better results than standard surgical technique in term of operative outcomes. Thus, it is important to determine the thorough position of the femoral tunnels. This study investigated whether a change in the femoral tunnel position in both axial and coronal planes can significantly alter the postoperative functional and clinical outcomes of the patients. Method: This comparative, retrospective, single-center study was performed on 44 patients who had underwent single-bundle Anterior Cruciate Ligament Reconstruction (ACLR). To evaluate the tunnel position in coronal and axial planes, radiographic assessments were done. Based on radiographic data, the patients were classified into 4 groups. The time interval between surgery and last visit averaged 23.6 ± 2.2 months (18-30 mos.). Lysholm knee score, and Cincinnati score were completed for all of the patients. Furthermore, the Lachman, anterior drawer and pivot-shift tests were performed. Results: Of the 44 patients included in the study, 9 patients (20.4%) were classified as the low-anterior group, 17(38.6%) were classified as the low-posterior group and 18(40.9%) were classified as the high-posterior group. None of the patients were included in high-anterior group. A greater mean Lysholm score (96 ± 3) in low-posterior group was the only significant difference between the three groups. Conclusion: Findings of the current study show that with the low-posterior placement of the femoral tunnel, based on both anteroposterior (AP) and tunnel-view x-rays, the better clinical outcomes can be achieved in short-term than the routine tunnel placements

Volume 7 Issue 2 - 2017

Seyed Morteza Kazemi¹, Ali Akbar Esmailijah², Mohammad Reza Abbasian³, Amir Hossein Keshavarz^{4*}, Ali Zafari⁵, Zahra Salehi Shahrabaki⁵ and Farshad Safdari⁶

¹Associate Professor of Orthopaedic Surgery, Bone, Joint and Related Tissues Research Center, Akhtar Hospital, Iran

²Assistance Professor of Orthopaedic Surgery, Bone, Joint and Related Tissues Research Center, Akhtar Hospital, Iran

³Professor of Orthopaedic Surgery, Bone, Joint and Related Tissues Research Center, Akhtar Hospital, Iran

⁴Orthopaedic Surgery Resident, Bone, Joint and Related Tissues Research Center, Akhtar Hospital, Iran

⁵Medical Student, Faculty of Medicine, Shahid Beheshti University of Medical Sciences, Iran

⁶Orthotist and Prosthetist, Bone, Joint and Related Tissues Research Center, Akhtar Hospital, Iran

*Corresponding author: Amir Hossein Keshavarz, MD, Department of orthopaedics, Akhtar Hospital, SharifiManesh St., Elahiye, Tehran, Iran, Tel: +98-912-2486544; Email: amirhosseinkeshavarz@aol.com

Received: September 20, 2016 | Published: January 17, 2017

Introduction

Arthroscopic techniques in knee surgeries have been introduced in recent decades and consequently our views on anterior cruciate ligament (ACL) reconstructions has been changed a lot over the years.

The favorable placement of the ACL graft is now under investigation. There are some factors that may affect achieving acceptable ACL reconstruction (ACLR) results. Incorrect placement of femoral and tibial tunnels seems to be a reasonable cause of failure in ACL reconstruction outcomes and it has been reported to be 4%-63% by several recent studies [1,2].

The evidences show that anatomic ACL graft positioning can restore rotational stability, resulting in better functional outcomes [3-6].

Recently, the more important role of femoral tunnel than the tibial tunnel has been emphasized [7]. Technically it is difficult to assess femoral tunnel position especially when the double-bundle ACLR is considered [8-10]. Femoral tunnel misplacement may result in a loss of flexion and an elongated graft and knee joint instability as a result of the substantial forces applied on the reconstructed tissue [11-16].

There is no robust consensus on whether the more oblique femoral tunnel position offers better results than standard surgical technique in term of postoperative knee laxity. Moreover, there are few studies concerning the impact of femoral tunnel position in both coronal and axial planes.

In this regard, it is important to determine the correct position of the tibial and femoral tunnels. The purpose of this study was to attempt to investigate whether a change in the femoral tunnel position in both axial and coronal planes could change the postoperative knee joint laxity include anterior, lateral or rotational instability in addition to functional outcomes of the patients.

Methods

This comparative, retrospective, single-center study was performed on 60 patients who had underwent single-bundle ACLR using semitendinosus autografts in 2013 by a expert surgeon. The trans-portal technique was used for the all cases.

All of the 60 patients were recalled to further evaluations. The patients had to be aged over 18 years, who did not have a history of multi-ligament injury, inflammatory arthritis and osteoarthritis. Also patients with non-anatomic femoral tunnel

position which was recognized by postoperative CT-Scanning and lateral view plain radiographs of the knee were excluded from study. The criteria for acceptable placement of the femoral tunnel based on the CT images were as follows:

The correct position was defined as placing posteromedial surface of the lateral condyle on axial plane. Also, the origin of femoral tunnel should be at 10 o'clock on the right and 2 o'clock on the left knee. The insertion of the tunnel should be on anterolateral, lateral or posterolateral of femur with 3-4 cm distant from lateral condyle [17,18]. The thickness of posterior cortex should be 1-2mm in axial slice.

Also using quadrant method on lateral view plain radiographs, we defined anatomic tunnel positions in sagittal plane. Thus, 44 patients (37male - 7male) aged 27.2 ± 5.6 years were eligible to be included in the study and 16 patients were excluded from the study.

To evaluate the femoral tunnel position in both coronal and axial planes, anteroposterior (AP) - and tunnel-view plain radiographs of the knee were taken. For tunnel view, the knee was placed in 60° flexion.

The tunnels were divided into two groups regarding their location on coronal plane according the AP x-rays.

- A. Low-position group: In this group femoral tunnel locates at 10o'clock for the right knee or 2o'clock for the left knee (30° from vertical line over the anatomic axis of femur).
- B. High-position group: Femoral tunnel locates at 11o'clock for the right knee or 1o'clock for the left knee (was oriented 60° from vertical line over the anatomic axis of femur) in this group of the patients.

However, in many of the cases, the tunnel was located between these values. Thus the low-position and high position were considered between 30° to 45° and 45° to 60° , respectively.

Furthermore, we determined the tunnels position based on the tunnel-view radiographs and patients were assigned to two groups:

- A. High-position group (Anterior group): In this group femoral tunnel located at 11 o'clock for the right knee or 1 o'clock for the left knee (60° from a line parallel to femur condyles) (Figure 1).
- B. Low-position group (Posterior group): in this group femoral tunnel located at 10 o'clock for the right knee or 2 o'clock for the left knee (30° from a line parallel to femur condyles) (Figure 2).(In fact, tunnels with this position will located more deeper in sagittal view of the notch than the anterior group.)

Similar to AP x-rays, in many of the cases, the tunnel was located between these values in tunnel-view x-rays. Thus the low-position and high position were considered between 30° to 45° and 45° to 60° , respectively.

Finally, the patients were classified into 4 groups included: low-anterior, low-posterior, high-anterior and high-posterior.




Figure 1: Correlation between the clock-face reference and the tunnel position in plain radiographs. (a) One o'clock position (high-position tunnel) in tunnel-view x-ray of the left knee. (b) The more anterior placement of the high-position femoral tunnel in comparison with low-position tunnel. (Note the endobutton insertion site.)




Figure 2: Correlation between the clock-face reference and the tunnel position in plain radiographs. (a) Ten o'clock position (low-position tunnel) in tunnel-view x-ray of the right knee. (b) The more posterior placement of the low-position femoral tunnel in comparison with high-position tunnel. (Note the endobutton insertion site.)

The time interval between surgery and last visit averaged 23.6 ± 2.2 months (18-30 mos.). Lysholm knee score [19], and Cincinnati score were completed for all of the patients. Furthermore, the Lachman, anterior drawer and pivot-shift tests were performed. Results of the Lachman and anterior drawer tests were considered positive if there was an anterior tibial translation $>5\text{mm}$ comparing the normal knee. Pivot-shift test results were graded as follows: 0 (absent), grade I (gentle slide), grade II (definite subluxation), and grade III (subluxation and momentary locking) [20]. All tests were performed by an expert orthopedist who was not a part of the investigation team and was blind to the group assignment. Also the intraobserver reliability of the examiner, based on a pilot study, was 0.9.

In addition, anterior tibial translation was assessed using the KT-1000 knee arthrometer for both operated and normal knees [21-23]. The maximum score for Lysholm knee score was 100 points, while higher scores indicated the better outcomes.

The Cincinnati score was categorized in 4 groups: excellent (80-100 points), good (55-79), fair (30-54), and poor (fewer than 30).

Statistical analysis was performed using SPSS statistical software (version 15.0; SPSS, Chicago, IL). One-way ANOVA and post hoc tests were utilized to compare quantitative data. Besides for comparing qualitative data, the chi-square test was employed. P value < 0.05 was considered significant.

Results

Of the 44 patients, 9 patients (20.4%) were classified as the low-anterior group, 17 patients (38.6%) as the low-posterior

group and 18 patients (40.9%) as the high-posterior group. None of the patients were included in high-anterior group. Table 1 show that there was no significant difference between 3 groups in term of age and gender. The mean Lysholm score was significantly higher in the low-posterior group ($p < 0.001$) (Table 2). However, the mean of Cincinnati score was the same in three groups. (Table 2) Of interest, only one patient in high posterior categorized as fair based on Cincinnati score, while all of the other patients were classified as good or excellent (Table 2). Additionally, anterior tibial translation did not differed significantly between the three groups (Table 3). Lachman test was negative in all of the patients. Anterior drawer test was negative in the low-posterior group. However, anterior drawer test was positive in 1 patient in the low-anterior (11.2%) and 1 patient in the high-posterior group (5.5%). Pivot shift test was graded IV in none of the patients (Table 3).

Table 1: Age-Sex distribution of the patients.

| | LA Group | LP Group | HP Group | p value |
|-----------|----------|----------|----------|---------|
| Age, yr | 25.7±4.2 | 28.4±4.4 | 27.5±8.3 | 0.411 |
| Gender, n | | | | 0.353 |
| Male | 7 | 16 | 14 | |
| | -77.80% | -94.10% | -77.80% | |
| Female | 2 | 1 | 4 | |
| | -22.20% | -5.90% | -22.20% | |

Table 2: Comparison of Lysholm and Cincinnati scores in different tunnel positions.

| | | LA Group | LP Group | HP Group | p value |
|--------------------------|-----------|------------------|------------------|-----------------|---------|
| | | n=9 | n=17 | n=18 | |
| Mean Lysholm Score | | 89±5 (82-100) | 96±3 (88-100) | 87±4 (84-94) | <0.001 |
| Lysholm Score Grading | Excellent | 7(77.8%) | 9(53%) | 14(77.8%) | 0.296 |
| | Good | 1(11.1%) | 8(47%) | 4(22.2%) | |
| | Fair | 1(11.1%) | 0 | 0 | |
| | Poor | 0 | 0 | 0 | |
| Mean Cincinnati Score | | 87±4 | 91±7 | 90±8 | 0.859 |
| Cincinnati Score Grading | Excellent | 8(88.8%) | 16(94.2%) | 12(66.7%) | 0.275 |
| | Good | 1(11.2%) | 1(5.8%) | 5(27.7%) | |
| | Fair | 0 | 0 | 1(5.6%) | |
| | Poor | 0 | 0 | 0 | |

Table 3: Comparison of clinical tests in different tunnel positions.

| | | LA Group | LP Group | HP Group | p value |
|-----------------------------------|----------|----------|----------|-----------|---------|
| | | n=9 | n=17 | n=18 | |
| Lachman test | Positive | 0 | 0 | 0 | - |
| | Negative | 9(100%) | 17(100%) | 18(100%) | |
| Anterior drawer test | Positive | 1(11.2%) | 0 | 1(5.5%) | 0.418 |
| | Negative | 8(88.8%) | 17(100%) | 17(94.5%) | |
| Pivot shift test | I | 5(55.5%) | 8(47.5%) | 7(38.8%) | 0.939 |
| | II | 3(33.3%) | 7(41.2%) | 9(50%) | |
| | III | 1(11.2%) | 2(11.8%) | 2(11.2%) | |
| | IV | 0 | 0 | 0 | |
| Anterior tibial translation* (mm) | | 2.2±0.4 | 2.5±0.7 | 2.4±0.5 | 0.444 |

Discussion

The main goal of the study was to see whether an alteration in the femoral tunnel position in both axial and coronal planes could change the postoperative joint laxity include anterior, lateral or rotational instability in addition to functional outcomes of the patients.

As stated by literature it is possible that patients reconstructed with a higher tunnel positions have an increased laxity as a result of misplaced femoral graft that does not mimic the positioning of the intact or normal ACL. In contrast, graft placement can restore normal knee motion if we perform it in the anatomic fashion [24-28].

Our findings demonstrated a greater mean Lysholm score in the low-posterior group in term of functional outcomes comparing the other groups of the patients. However, we did not find a significant difference in the remaining clinical evaluations include Cincinnati score, Lachman test, pivot shift test and anterior drawer test. A study by Lee et al. [29] showed a lower Lysholm score and higher femoral tunnel positioning in the knees with positive pivot shift test than in the knees without pivot shift.

This is not in accordance with the results of Tsuda et al. [30] who found that the difference between the low- and high-positions is not enough to convince them that different tunnel positions can be associated with clinical and functional outcomes. Beside, Markolf reported on a method for the impact of linear regression slopes for the femoral tunnels on postoperative results. He concluded that the slope difference between the above-mentioned positions was not so significant as to be a reason for any advantage of the oblique (or low) tunnel over standard (or high) tunnel positions [31]. It seems that femoral tunnel obliquity may result in marked clinical outcomes if there be a great difference between tunnels linear slope.

Practically the other findings in our study are not clarifying in favor of which tunnel position is preferable. In this regard, the harvested data from anterior drawer and Lachman tests support that all the 3 tunnel positions are quite enough for stopping the anterior tibial translation. This maybe because of the ACLR surgery and its outcome, in which almost all anatomic reconstructed ACLs can control anterior tibial translation [31].

It has been common to place the femoral graft at 11 o'clock position to recover the function of the AM bundle of the ACL [14,32]. Once an ACL reconstructed knee is subjected to rotatory loads, the high-position tunnel for graft placement will not avoid rotational instability anymore [6,33,34]. Moreover, It has been revealed that the 10 o'clock position resembles the PL bundle attachment and can be more sufficient at rotatory loads and limiting anterior tibial translation [35] which was confirmed by other biomechanical studies [14,36]. So it can answer this question that why the mean Lysholm score in the low-posterior group could be greater comparing two other groups and consequently justify the fact that how our remaining non-biomechanical evaluations have the same results.

We did not found a detectable difference in the pivot test between three groups. This test is the most widely used dynamic test, which correlates with instability symptoms [37]. However, our results could be due to the low sensitivity of the pivot shift test [38]. This is in accordance with the result of Jepsen et al. who found no difference between the high- and low-position tunnels regarding anterior laxity and pivot shift test [39].

There are a number of potential limitations that warrant consideration. The first is that in our view the postoperative radiologic assessment of tunnel positions can be somehow challenging with routine radiography, as this is 2-Dimensional illustration of a 3-Dimensional situation [40]. This is a reason why we should investigate tunnel positions in both axial and

coronal planes concurrently. Furthermore, it is found that the tunnel-view radiograph is not satisfactory to assess the femoral tunnel placement, because there are variations in radiographic projection at the different phases of postoperative evaluation of the same patient [39]. The second limitation concerns that the mean follow-up time was about 2 years, and thus we cannot debate about the long-term surgical outcomes associated with clinical and radiologic developments. According to the literature Lysholm score was not sensitive to detect changes over time, and then it cannot be a precise scoring scale for long-term postoperative follow-up [41].

The final limitation was that like most related studies [39,42,43] we did not consider the tibial tunnel position which can be one of the important factors on clinical results.

The methodological pitfalls that we encountered with them include: first, the number of patients in this study should be more due to the study type. The second is the fact that this retrospective study investigates the tunnel positions and ACL grafts in patients with previous ACLR surgeries and consequently we had no role in their tunnel positioning.

Conclusion

Although the anatomic ACLR can sufficiently restore the knee stability and be associated with considerable functional improvement, the current study showed that low-posterior tunnel placement resulted in significantly higher knee scores. It is important to consider the femoral tunnel position in different planes and further investigations are required.

Disclosure

The authors report no conflict of interest concerning the materials or methods used in this study or the findings specified in this paper.

References

1. Howell Msm, Clark Ja (1992) Tibial tunnel placement in anterior cruciate ligament reconstructions and graft impingement. *Clinical orthopaedics and related research* 283: 187-195.
2. Williams RJ, Hyman J, Petrigliano F, Rozental T, Wickiewicz TL (2004) Anterior cruciate ligament reconstruction with a four-strand hamstring tendon autograft. *J Bone Joint Surg Am* 86(2): 225-232.
3. Bedi A, Raphael B, Maderazo A, Pavlov H, Williams RJ (2010) Transtibial versus anteromedial portal drilling for anterior cruciate ligament reconstruction: a cadaveric study of femoral tunnel length and obliquity. *Arthroscopy* 26(3): 342-350.
4. Heming JF, Rand J, Steiner ME (2007) Anatomical limitations of transtibial drilling in anterior cruciate ligament reconstruction. *Am J Sports Med* 35(10): 1708-1715.
5. Lubowitz JH (2009) Anteromedial portal technique for the anterior cruciate ligament femoral socket: pitfalls and solutions. *Arthroscopy* 25(1): 95-101.
6. Yagi M, Wong EK, Kanamori A, Debski RE, Fu FH, et al. (2002) Biomechanical analysis of an anatomic anterior cruciate ligament reconstruction. *Am J Sports Med* 30(5): 660-666.
7. Ahn JH, Kim JD, Kang HW (2015) Anatomic Placement of the Femoral Tunnels in Double-Bundle Anterior Cruciate Ligament Reconstruction Correlates With Improved Graft Maturation and Clinical Outcomes. *Arthroscopy* 31(11): 2152-2161.
8. Kawakami Y, Hiranaka T, Matsumoto T, Hida Y, Fukui T, et al. (2012) The accuracy of bone tunnel position using fluoroscopic-based navigation system in anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 20(8): 1503-1510.
9. Lertwanich P, Martins CA, Asai S, Ingham SJ, Smolinski P, et al. (2011) Anterior cruciate ligament tunnel position measurement reliability on 3-dimensional reconstructed computed tomography. *Arthroscopy* 27(3): 391-398.
10. Tsukada H, Ishibashi Y, Tsuda E, Fukuda A, Toh S (2008) Anatomical analysis of the anterior cruciate ligament femoral and tibial footprints. *J Orthop Sci* 13(2): 122-129.
11. Amis A, Zavras T (1995) Isometricity and graft placement during anterior cruciate ligament reconstruction. *The Knee* 2(1): 5-17.
12. Good L, Odensten M, Gillquist J (1994) Sagittal Knee Stability After Anterior Cruciate Ligament Reconstruction with a Patellar Tendon Strip A Two-year Follow-up Study. *Am J Sports Med* 22(4): 518-523.
13. Howell SM, Taylor M (1993) Failure of reconstruction of the anterior cruciate ligament due to impingement by the intercondylar roof. *J Bone Joint Surg Am* 75(7): 1044-1055.
14. Loh JC, Fukuda Y, Tsuda E, Steadman RJ, Fu FH, et al. (2003) Knee stability and graft function following anterior cruciate ligament reconstruction: Comparison between 11 o'clock and 10 o'clock femoral tunnel placement. *Arthroscopy* 19(3): 297-304.
15. Musahl V, Burkart A, Debski RE, Van Scyoc A, Fu FH, et al. (2003) Anterior cruciate ligament tunnel placement: comparison of insertion site anatomy with the guidelines of a computer-assisted surgical system. *Arthroscopy* 19(2): 154-160.
16. Sommer C, Friederich NF, Müller W (2000) Improperly placed anterior cruciate ligament grafts: correlation between radiological parameters and clinical results. *Knee Surg Sports Traumatol Arthrosc* 8(4): 207-213.
17. Canale ST, Beaty JH (2012) *Campbell's Operative Orthopaedics*. Elsevier Health Sciences.
18. Griffin LY, Agel J, Albohm MJ, Arendt EA, Dick RW, et al. (2000) Noncontact anterior cruciate ligament injuries: risk factors and prevention strategies. *J Am Acad Orthop Surg* 8(3): 141-150.
19. Lysholm J, Gillquist J (1982) Evaluation of knee ligament surgery results with special emphasis on use of a scoring scale. *The Am J Sports Med* 10(3): 150-154.
20. Seon JK, Park SJ, Lee KB, Seo HY, Kim MS, et al. (2011) In vivo stability and clinical comparison of anterior cruciate ligament reconstruction using low or high femoral tunnel positions. *Am J Sports Med* 39(1): 127-133.
21. Daniel DM, Malcom LL, Losse G, Stone ML, Sachs R, et al. (1985) Instrumented measurement of anterior laxity of the knee. *The Journal of Bone & Joint Surgery* 67(5): 720-726.
22. Steiner ME, Brown C, Zarins B, Brownstein B, Koval P, et al. (1990) Measurement of anterior-posterior displacement of the knee. A comparison of the results with instrumented devices and with clinical examination. *J Bone Joint Surg Am* 72(9): 1307-1315.

23. Wroble RR, Van Ginkel LA, Grood ES, Noyes FR, Shaffer BL (1990) Repeatability of the KT-1000 arthrometer in a normal population. *The American Journal of Sports Medicine* 18(4): 396-399.
24. Abebe ES, Utturkar G, Taylor D, Spritzer C, Kim J, et al. (2011) The effects of femoral graft placement on in vivo knee kinematics after anterior cruciate ligament reconstruction. *J biomech* 44(5): 924-929.
25. Allen CR, Giffin JR, Harner CD (2003) Revision anterior cruciate ligament reconstruction. *Orthop Clin North Am* 34(1): 79-98.
26. Harner CD, Giffin JR, Duntzman RC, Annunziata CC, Friedman MJ (2000) Evaluation and Treatment of Recurrent Instability After Anterior Cruciate Ligament Reconstruction. *J Bone Joint Surg Am* 82(11):1652-1652.
27. Jaureguito JW, Paulos LE (1996) Why grafts fail. *Clinical orthopaedics and related research* 325: 25-41.
28. Johnson DL, Swenson TM, Irrgang JJ, Fu FH, Harner CD (1996) Revision anterior cruciate ligament surgery: experience from Pittsburgh. *Clinical orthopaedics and related research* 325: 100-109.
29. Lee MC, Seong SC, Lee S, Chang CB, Park YK, Jo H, et al. (2007) Vertical femoral tunnel placement results in rotational knee laxity after anterior cruciate ligament reconstruction. *Arthroscopy* 23(7): 771-778.
30. Tsuda E, Ishibashi Y, Fukuda A, Yamamoto Y, Tsukada H, et al. (2010) Tunnel position and relationship to postoperative knee laxity after double-bundle anterior cruciate ligament reconstruction with a transtibial technique. *Am J Sports Med* 38(4): 698-706.
31. Markolf KL, Jackson SR, McAllister DR (2010) A Comparison of 11 O'clock Versus Oblique Femoral Tunnels in the Anterior Cruciate Ligament-Reconstructed Knee Knee Kinematics During a Simulated Pivot Test. *Am J Sports Med* 38(5): 912-917.
32. Sapega AA, Moyer R, Schneck C, Komalahiranya N (1990) Testing for isometry during reconstruction of the anterior cruciate ligament. Anatomical and biomechanical considerations. *J Bone Joint Surg Am* 72(2): 259-267.
33. Kanamori A, Woo SL, Ma CB, Zeminski J, Rudy TW, Li G, et al. (2000) The forces in the anterior cruciate ligament and knee kinematics during a simulated pivot shift test: a human cadaveric study using robotic technology. *Arthroscopy* 16(6): 633-639.
34. Woo SL, Kanamori A, Zeminski J, Yagi M, Papageorgiou C, et al. (2002) The effectiveness of reconstruction of the anterior cruciate ligament with hamstrings and patellar tendon. *J Bone Joint Surg Am* 84(6): 907-914.
35. Sakane M, Fox RJ, Glen SLYW, Livesay A, Li G, Fu FH (1997) In situ forces in the anterior cruciate ligament and its bundles in response to anterior tibial loads. *J Orthop Res* 15(2): 285-293.
36. Scopp JM, Jasper LE, Belkoff SM, Moorman CT (2004) The effect of oblique femoral tunnel placement on rotational constraint of the knee reconstructed using patellar tendon autografts. *Arthroscopy* 20(3): 294-299.
37. Kujala U, Nelimarkka O, Koskinen S (1992) Relationship between the pivot shift and the configuration of the lateral tibial plateau. *Archives of orthopaedic and trauma surgery* 111(4): 228-229.
38. Scholten R, Opstelten W, van der Plas CG, Bijl D, Deville W, et al. (2003) Accuracy of physical diagnostic tests for assessing ruptures of the anterior cruciate ligament: a meta-analysis. *Journal of family practice* 52(9): 689-694.
39. Jepsen CF, Lundberg-Jensen AK, Faunoe P (2007) Does the position of the femoral tunnel affect the laxity or clinical outcome of the anterior cruciate ligament-reconstructed knee? A clinical, prospective, randomized, double-blind study. *Arthroscopy* 23(12): 1326-1333.
40. Martins CA, Kropf EJ, Shen W, van Eck CF, Fu FH (2012) The concept of anatomic anterior cruciate ligament reconstruction. *Operative Techniques in Sports Medicine* 16(3): 104-115.
41. Risberg M, Holm I, Steen H, Beynnon BD (1997) Sensitivity to changes over time for the IKDC form, the Lysholm score, and the Cincinnati knee score A prospective study of 120 ACL reconstructed patients with a 2-year follow-up. *Knee Surg Sports Traumatol Arthrosc* 7(3): 152-159.
42. Fernandes TL, Fregni F, Weaver K, Pedrinelli A, Camanho GL, et al. (2014) The influence of femoral tunnel position in single-bundle ACL reconstruction on functional outcomes and return to sports. *Knee Surg Sports Traumatol Arthrosc* 22(1): 97-103.
43. Luites JW, Verdonshot PN (2015) Radiographic positions of femoral ACL, AM and PL centres: accuracy of guidelines based on the lateral quadrant method. *Knee Surg Sports Traumatol Arthrosc* 1-9.