

Review Article





Leveraging peer-network interventions to increase the uptake and continuation of pre-exposure prophylaxis by adolescent girls and young women at high risk of HIV in low and middle-income countries

Abstract

Background: Adolescent girls and young women (AGYW) in low- and middle-income countries (LMICs) continue to bear a disproportionate burden of new HIV infections, driven by structural, social, and behavioral vulnerabilities. Although oral pre-exposure prophylaxis (PrEP) is an effective biomedical prevention method, its uptake and continuation among AGYW remain suboptimal. This narrative review explores the effectiveness of peernetwork interventions in improving PrEP uptake and continuation among AGYW at high risk of HIV in LMICs.

Methods: A comprehensive search of PubMed, Embase and Google Scholar databases was conducted to identify peer-reviewed articles published between January 2010 and October 2024. Eligible studies involved AGYW aged 15–25, the studies were conducted in LMICs, and evaluated peer-based interventions with outcomes related to PrEP uptake, adherence, or continuation.

Results: Eleven studies which met the inclusion criteria, spanning sub-Saharan African countries were selected from 1003 references. Peer-network strategies ranged from peer outreach and education to adherence clubs and behavior-centered social marketing. Initiation rates ranged from 28.1% to 95%, with significantly higher uptake reported in peer-supported interventions compared to standard approaches. However, continuation rates remained modest (<30%). Peer strategies were found to be acceptable and feasible, especially when integrated into youth-friendly settings, though challenges with sustainability and system integration persist.

Conclusion: Peer-network interventions offer a promising approach to enhance PrEP uptake among AGYW in LMICs. However, sustained continuation will require multi-layered support, digital innovations, and integration into broader adolescent health platforms. Strategic investment in peer-led models could strengthen HIV prevention efforts and reduce incidence in this high-risk population.

Keywords: pre-exposure prophylaxis, HIV prevention, adolescent girls, young women, peer-network interventions, PrEP uptake, low- and middle-income countries, implementation strategies

Volume 14 Issue 3 - 2025

Omenoba Theodora C,¹ Ogbonna Brian O,^{2,3,4} Ibeneme Georgian C,^{3,5} Ogbonna Chigozie A,³ Folaranmi Nkeiruka⁵

Department of Clinical Pharmacy and Pharmacy Management, Faculty of Pharmacy, Veritas University, Abuja, Nigeria Department of Clinical Pharmacy and Pharmacy Practice, Faculty of Pharmacy, David Umahi Federal University of Health Sciences, Uburu, Nigeria

³International Institute for Pharmaceutical Research and Innovations (IIPRI), David Umahi Federal University of Health Sciences, Uburu, Ebonyi State, Nigeria

International Institute for Health Policy Research and Translation (IIHPRT), David Umahi Federal University of Health Sciences, Uburu, Ebonyi State, Nigeria

⁵Faculty of Dentistry, College of Medicine, David Umahi Federal University of Health Sciences, Uburu, Nigeria

Correspondence: Ogbonna BO, Department of Clinical Pharmacy and Pharmacy Practice, Faculty of Pharmacy, David Umahi Federal University of Health Sciences, Uburu, Nigeria, Tel +2348037794206

Received: August 25, 2025 | Published: September 15, 2025

Introduction

Globally, adolescent girls and young women (AGYW) between the ages of 15 and 24 face a disproportionately high risk of HIV infection, particularly in sub-Saharan Africa. This vulnerability is influenced by biological, social, structural, and behavioral factors including engagement in sex work or transactional sex, unequal gender power dynamics, and lack of youth-friendly sexual and reproductive health services. While transactional sex differs from commercial sex work, both increase susceptibility to HIV acquisition due to overlapping risk factors such as inconsistent condom use, multiple sexual partners, and limited negotiating power. 3.6-8

PrEP is a scientifically proven biomedical intervention involving the use of daily oral antiretrovirals by individuals at high risk of HIV, offering up to 99% protection when adherence is optimal. 9–13 However, real-world implementation has revealed significant gaps in uptake and adherence, especially among AGYW in LMICs. Societal and systemic

barriers such as stigma, clinic-level factors, legal restrictions, and fear of identification as high-risk individuals deter young women from initiating and continuing PrEP.¹⁴⁻²² The intersection of adolescent risk behaviors, peer influence, and emerging autonomy adds another layer of complexity in delivering sustained prevention efforts to this population.²³⁻²⁵

Peer-network interventions are gaining recognition as culturally acceptable and potentially scalable strategies to increase PrEP engagement. These approaches harness the power of peer influence, leveraging trust and relatability to bridge information and access gaps.

Methods

This narrative review synthesized existing evidence on peernetwork interventions aimed at increasing the uptake and continuation of pre-exposure prophylaxis (PrEP) among adolescent girls and young women (AGYW) at high risk of HIV infection in low- and





middle-income countries (LMICs). The review primarily draws from studies identified in a comprehensive literature search using structured methodologies to ensure the relevance and comprehensiveness of the evidence included. A systematic search was conducted using three databases: PubMed, Embase, and Google Scholar for studies published between January 2010 and October 2024. This time-frame was chosen to reflect data generated after the:

U.S. Food and Drug Administration (FDA) approved daily oral PrEP in 2012 and to capture insights from the global expansion of PrEP programs. The search strategy included both Medical Subject Headings (MeSH) and free-text keywords such as "PrEP," "peer support," "peer network," "adolescent girls," "young women," "HIV prevention," and "low-income countries. "Inclusion criteria were: (1) studies conducted in LMICs as defined by the WoBank, (2) participants aged 15–25 years, (3) studies evaluating interventions that involved a peer component (e.g., peer educators, peer support groups, peer marketing), and (4) outcomes related to PrEP uptake, continuation, or adherence. Eligible studies comprised randomized controlled trials, observational studies, implementation research, and mixed-methods evaluations. Studies such as commentaries, editorials, and those that did not focus on adolescent girls and young women (AGYW) or did not involve peer-led interventions were excluded.

Key data were extracted from each study, including country, study design, target population, type of intervention, primary outcomes, and implementation outcomes. The review applied Proctor's Implementation Outcomes Framework to organize the findings, evaluating each intervention across dimensions such as acceptability, adoption, feasibility, penetration, and sustainability. Findings were synthesized thematically, emphasizing patterns, gaps, and lessons relevant to the design and implementation of peer-led or peer-supported PrEP interventions in real-world settings.

Results

Characteristics of included studies

A total of 1003 references were retrieved by the database searches on 13 March 2025, resulting in 405 unique results for the title and abstract screening. Fifty-five of the retrieved references were judged to be appropriate after screening their title/abstract against the eligibility criteria. A total of 11 studies were included in the review after the screening of the full manuscripts against the eligibility criteria (Figure 1 & Table 1).

Table I Evidence-based table of selected studies

Study / Project	Country	Design	Target population	Intervention	Key findings	Implementation outcomes
HPTN 082 Celum, ²⁸ Velloza, ²²	South Africa, Zimbabwe	RCT + Qualitative	AGYW 16–25 yrs	SMS + Peer Clubs + Feedback	95% initiation; ~20% continuation	High adoption; Low sustainability
POWER Cohort study, ^{32,33}	Kenya, South Africa	Cohort + Qualitative	AGYW 16–25 yrs	Youth- friendly clinics	94% initiation; 20% continuation	High adoption; Feasible; Low penetration
DREAMS Chabata, ³⁴ Hensen, ³⁵	Zimbabwe	Non- randomized	Young women who sell sex	Peer outreach + support	28.1% vs 0.6% uptake (intervention vs control)	High adoption
DREAMS (SA) Chimbindi, ³¹	South Africa	Mixed- methods	Young women who sell sex	Hospital + Peer outreach	0% uptake; awareness ↑ 2% → 9%	Low adoption; awareness improved
MPYA Trial, ^{29,30}	Kenya	RCT	AGYW (VOICE score ≥5)	Daily SMS	No significant effect on continuation	Low sustainability
3Ps for Prevention, ^{26,27}	South Africa	RCT + Mixed	AGYW sexually active	Marketing + Incentives	56% interested; 56% vs. 41% adherence (p=0.067)	Moderate adoption; borderline sustainability

Peer-network interventions and demand creation

Effective demand creation is pivotal in encouraging AGYW to consider and initiate PrEP. In high HIV-incidence settings such as South Africa, peer-driven social marketing has shown promise in increasing awareness and intent. A culturally resonant campaign, which incorporated messages like "PrEP enhances the power you have" and "PrEP protects those you love," was disseminated via peer influencers and youth social networks. 26-27 Following the intervention, over 56% of AGYW expressed a definitive interest in initiating PrEP, highlighting the potential of targeted communication through trusted peer channels. 27 Such demand-generation strategies are grounded in behavior-centered design, where peers act not only as conduits of information but also as social proof that reinforces PrEP as a normative and empowering health choice. 27

Peer support for PrEP initiation

Beyond generating interest, peer support structures can influence the transition from intent to actual PrEP initiation. In some studies, peer educators embedded within community-based HIV programs identified eligible AGYW and linked them to PrEP services through youth-friendly entry points such as schools, mobile clinics, or adolescent health centers.²⁶⁻²⁷

Evidence from Zimbabwe demonstrated significantly higher PrEP uptake in sites where peer outreach was integrated into DREAMS programming (28.1% vs. 0.6% in control sites; adjusted OR 63.82, 95% CI 19.78 to 205.90). Similarly, peer mobilization within existing social structures improved the identification and referral of AGYW who otherwise might not have accessed prevention services.

Citation: Theodora OC, Brian OO, Georgian IC, et al. Leveraging peer-network interventions to increase the uptake and continuation of pre-exposure prophylaxis by adolescent girls and young women at high risk of HIV in low and middle-income countries. MOJ Public Health. 2025;14(3):224–227. DOI: 10.15406/mojph.2025.14.00494




Figure 1 Study articles selection process flowchart.

Peer Strategies for PrEP continuation and adherence support

While uptake is crucial, maintaining adherence to PrEP over time presents an even greater challenge. Peer adherence clubs and digital peer-led support mechanisms have been tested to address this gap.

In the HPTN 082 trial, AGYW was offered standard adherence support plus two-way SMS messaging and the option to join peerled adherence clubs. Although initiation rates were high (95%), only 20.9% of participants remained on PrEP at six months, with no statistically significant difference between enhanced support and control groups. Similarly, in the MPYA trial, SMS reminders alone did not significantly improve continuation rates (27.0% vs. 26.7%; adjusted IR 1.16, 95% CI 0.93 to 1.45). These findings suggest that peer engagement, while helpful, must be embedded in broader structural and emotional support systems to impact sustained use. Nevertheless, AGYW reported that peer adherence clubs reduced feelings of isolation, enhanced privacy, and mitigated stigma—factors known to influence long-term engagement.

Peer influence in combination interventions

The DREAMS initiatives in Zimbabwe and South Africa employed multi-layered strategies that included peer-led services alongside economic empowerment, social protection, and gender- based violence prevention. ^{26,27,31} In South Africa, although no participants initiated PrEP during the evaluation period, awareness increased over time, suggesting a cumulative effect of peer engagement and education. ³¹

Notably, in Zimbabwe, the involvement of young women who sell sex as peer educators contributed significantly to higher PrEP initiation, emphasizing the role of relatable messengers in reaching hidden or stigmatized populations.²⁷

Implementation outcomes and real-world considerations

Assessing peer-network interventions through an implementation science lens reveals insights into their potential for scale-up:

Acceptability: Peer-led interventions are generally well-received, particularly when they mirror the age, gender, and lived experiences of AGYW.³²

Adoption: Studies with peer components report initiation rates between 56%–95%, far above national averages in many LMICs. ^{26,28,33}

Feasibility: Peer-based delivery was more effective in youth-friendly settings, such as private clinics than in traditional family planning settings.³²

Sustainability: Continuation rates remain low, often under 30%, indicating that peer influence must be continuously nurtured and supplemented with system-level support.^{28,29}

Penetration: Integration into routine services remains limited. For example, in some hospital settings, non-study staff viewed PrEP as outside their scope, limiting program reach.³²

Challenges and future directions

Despite their promise, peer interventions face several challenges:

Resource intensity: Sustaining trained peer educators and ensuring quality delivery requires consistent investment.

Structural barriers: Legal and policy restrictions around adolescent consent, stigma, and criminalization of sex work continue to undermine peer strategies.

Fragmented service delivery: Without integration into existing health systems, peer programs may remain siloed and unsustainable.

Future directions include: Digital peer models, such as WhatsApp groups or virtual adherence buddies; Peer-facilitated differentiated service delivery, including PrEP delivery through pharmacies and mobile units; Evaluation frameworks that capture cost-effectiveness and real- world fidelity.^{33–35}

Conclusion

Peer-network interventions hold substantial potential to improve PrEP uptake and continuation among AGYW at high risk of HIV in LMICs. When designed with youth-centric principles and embedded in broader structural support, peer strategies can drive demand, foster trust, and support adherence. However, their success depends on integration into sustainable, scalable health systems and consistent evaluation of impact. Future programming should aim at strengthen peer networks as an essential pillar in adolescent HIV prevention.

Acknowledgments

None.

Conflicts of interest

The authors declare there is no conflict of interest.

Funding

None.

References

- 1. World Health Organization. Adolescent health. 2021.
- 2. UNAIDS. In danger: UNAIDS global AIDS update. Geneva: Joint United Nations Programme on HIV/AIDS, 2022.
- 3. Stoebenau K, Heise L, Wamoyi J, et al. Revisiting the understanding of "transactional sex" in sub-Saharan Africa: A review and synthesis of the literature. *Soc Sci Med.* 2016;168:186–197.

Citation: Theodora OC, Brian OO, Georgian IC, et al. Leveraging peer-network interventions to increase the uptake and continuation of pre-exposure prophylaxis by adolescent girls and young women at high risk of HIV in low and middle-income countries. MOJ Public Health. 2025;14(3):224–227. DOI: 10.15406/mojph.2025.14.00494

- 4. Jewkes R, Dunkle K, Nduna M, et al. Transactional sex and HIV incidence in a cohort of young women in the stepping stones trial. J AIDS Clin Res. 2012;3:5.
- 5. Juma M, Alaii J, Bartholomew LK, et al. Risky sexual behavior among orphan and non-orphan adolescents in Nyanza Province, Western Kenya. AIDS Behav. 2013;17:951-960.
- 6. Konstant TL, Rangasami J, Stacey MJ, et al. Estimating the number of sex workers in South Africa: rapid population size estimation. AIDS Behav. 2015;19 Suppl 1:3-15.
- 7. Wamoyi J, Stobeanau K, Bobrova N, et al. Transactional sex and risk for HIV infection in sub- Saharan Africa: a systematic review and meta-analysis. J Int AIDS Soc. 2016;19(1):20992.
- 8. Kerrigan D, Wirtz A, Semini I. The global HIV epidemic among sex workers. The World Bank, 2012.
- 9. Baeten JM, Donnell D, Ndase P, et al. Antiretroviral prophylaxis for HIV prevention in heterosexual men and women. N Engl J Med. 2012;367(5):399-410.
- 10. Choopanya K, Martin M, Suntharasamai P, et al. Antiretroviral prophylaxis for HIV infection in injecting drug users in Bangkok, Thailand (the Bangkok Tenofovir study): a randomized, double-blind, placebo--controlled phase 3 trial. Lancet. 2013;381(9883):2083-2090.
- 11. Grant RM, Lama JR, Anderson PL, et al. Pre-exposure chemoprophylaxis for HIV prevention in men who have sex with men. N Engl J Med. 2010:363(27):2587-2599.
- 12. McCormack S, Dunn DT, Desai M, et al. Pre-exposure prophylaxis to prevent the acquisition of HIV-1 infection (PROUD): effectiveness results from the pilot phase of a pragmatic open-label randomized trial. Lancet. 2016;387(10013):53-60.
- 13. CDC. Prep effectiveness | PreP | HIV basics | HIV/AIDS | CDC, 2021.
- 14. Irungu EM, Baeten JM. Prep rollout in Africa: status and opportunity. Nat Med. 2020;26(5):655-664.
- 15. Koss CA, Charlebois ED, Ayieko J, et al. Uptake, engagement, and adherence to pre-exposure prophylaxis offered after population HIV testing in rural Kenya and Uganda: 72-week interim analysis of observational data from the search study. Lancet HIV. 2020;7(4):e249-e261.
- 16. Atkins K, Kan L, Musau A, et al. Adaptation and psychometric evaluation of a scale to measure oral pre-exposure prophylaxis-related stigma among key and vulnerable populations in Kenya. J Int AIDS Soc. 2022;25(Suppl 1):e25929.
- 17. Yun K, Xu J-J, Zhang J, et al. Female and younger subjects have lower adherence in PreP trials: a meta-analysis with implications for the uptake of PreP service to prevent HIV. Sex Transm Infect. 2018;94(3):163–168.
- 18. UNAIDS. UNAIDS Data 2000. In: Geneva: UNAIDS joint United Nations program on HIV/AIDS; 2020. Joint United Nations Programme on HIV/AIDS, 2020.
- 19. Phoenix J. Regulating sex for sale: prostitution policy reform in the UK. 1st ed. Bristol University Press, 2009.
- 20. Luecke EH, Cheng H, Woeber K, et al. Stated product formulation preferences for HIV pre- exposure prophylaxis among women in the VOICE-D (MTN-003D) study. J Int AIDS Soc. 2016;19(1):20875.

- 21. Rousseau E, Katz AWK, O'Rourke S, et al. Adolescent girls and young women's PrEP-user journey during an implementation science study in South Africa and Kenya. PLoS One. 2021;16(10):e0258542.
- 22. Velloza J, Khoza N, Scorgie F, et al. The influence of HIV-related stigma on PreP disclosure and adherence among adolescent girls and young women in HPTN 082: a qualitative study. J Int AIDS Soc. 2020;23(3):e25463.
- 23. Albert D, Chein J, Steinberg L. The teenage brain: peer influences on adolescent decision making. Curr Dir Psychol Sci. 2013;22(2):114-120.
- 24. Chein J, Albert D, O'Brien L, et al. Peers increase adolescent risk--taking by enhancing activity in the brain's reward circuitry. Dev Sci. 2011:14(2):F1-10.
- 25. Gardner M, Steinberg L. Peer influence on risk-taking, risk preference, and risky decision making in adolescence and adulthood: an experimental study. Dev Psychol. 2005;41(4):625-635.
- 26. Celum CL, Gill K, Morton JF, et al. Incentives conditioned on tenofovir levels to support PreP adherence among young South African women: a randomized trial. J Int AIDS Soc. 2020;23(11):e25636.
- 27. Morton JF, Myers L, Gill K, et al. Evaluation of a behavior-centered design strategy for creating demand for oral PreP among young women in Cape Town, South Africa. Gates Open Res. 2020;4:29.
- 28. Celum C, Hosek S, Tsholwana M, et al. Prep uptake, persistence, adherence, and effect of retrospective drug level feedback on PreP adherence among young women in southern Africa: results from HPTN 082, a randomized controlled trial. PLoS Med. 2021;18(6):e1003670.
- 29. Haberer JE, Mugo N, Bukusi EA, et al. Understanding pre-exposure prophylaxis adherence in young women in Kenya. J Acquir Immune Defic Syndr. 2022;89(3):251-260.
- 30. Haberer JE, Bukusi EA, Mugo NR, et al. Effect of SMS reminders on PreP adherence in young Kenyan women (MPYA study): a randomised controlled trial. Lancet HIV. 2021;8(3):e130-e137.
- 31. Chimbindi N, Mthiyane N, Zuma T, et al. Antiretroviral therapy based HIV prevention targeting young women who sell sex: a mixed method approach to understand the implementation of PreP in a rural area of KwaZulu-Natal, South Africa. AIDS Care. 2022;34(2):232-240.
- 32. Roche SD, Barnabee G, Omollo V, et al. Implementation strategies for integrating pre- exposure prophylaxis for HIV prevention and family planning services for adolescent girls and young women in Kenya: a qualitative study. BMC Health Serv Res. 2022;22(1):422.
- 33. Celum CL, Bukusi EA, Bekker LG, et al. Prep use and HIV seroconversion rates in adolescent girls and young women from Kenya and South Africa: the power demonstration project. J Int AIDS Soc. 2022;25(7):e25962.
- 34. Chabata ST, Hensen B, Chiyaka T, et al. The impact of the Dreams partnership on HIVincidence among young women who sell sex in two Zimbabwean cities: results of a non- randomized study. BMJ Glob Health. 2021;6(4):e003892.
- 35. Hensen B, Machingura F, Busza J, et al. How can we support the use of oral PreP among young women who sell sex? A PreP cascade analysis. J Acquir Immune Defic Syndr. 2021;88(1):45-56.