

Assessment of food frequency for diabetic patients after coronary artery bypass graft at ibn al-bitar specialized center for cardiac surgery in Baghdad city

Abstract

Diabetes is a serious public health problem with serious secondary complications, one of these problems are coronary artery disease. The global burden of diabetes is large and growing, affecting populations in every region of the world.

Objective: To assess the diabetic patients dietary habits and find out the relationship between these patients food frequency and their socio-demographic characteristics of age, gender, marital status, and economic status.

Methodology: A descriptive analytical study was conducted on Non-probability (purposive sample) of (100) a diabetic patients at (Ibn Al-Bitar specialized center for cardiac surgery) in Baghdad City. A questionnaire was used as a tool of data collection for the period of 12th May to 12th July 2016. Descriptive statistical analyses were used to analyze the data.

Results: The results of the study shows that the majority of the age group were (60-69) years old (41%). Most of the study samples (73%) were male. Most of them (69%) were married and (62%) the test of (HbA1c) indicates more than seven. More than study samples (91%) do not follow the diet program.

Conclusion: The study results indicate highly significant differences association between age group with knowledge of the dietary habits. There is indicated that high significant association between marital status and dietary habits high significant differences association between marital statuses food frequency which are more frequent among married patients.

Recommendation: Diabetes self-management education (DSME) and support (DSMS) programs are appropriate for education and support to develop and maintain behaviors that can prevent diabetes, perform the A1C test at least two times a year in patients who are meeting treatment goals. In adults, a screening lipid profile is reasonable at the time of first diagnosis, at the initial medical evaluation twice yearly.

Keywords: assessment, diabetic patients, food frequency, coronary artery bypass graft

Volume 5 Issue 1 - 2018

Saad M Abdulhameed, Khalida M Khudur

Department of Adults Nursing, College of Nursing, University of Baghdad, Iraq

Correspondence: Khalida M Khudur; Department of Adults Nursing, College of Nursing, University of Baghdad, Iraq, Email: kdalida_mohammed@yahoo.com

Received: April 21, 2017 | **Published:** January 09, 2018

Introduction

All forms of diabetes increase the risk of long-term complications. These typically develop after many years (10-20), but may be the first symptom in those who have otherwise not received a diagnosis before that time.¹ The major long-term complications relate to damage to blood vessels. Diabetes doubles the risk of cardiovascular disease and about 75% of deaths in diabetics are due to coronary artery disease.²

The role of nutrition in maintaining health and reducing the damage caused by diabetes and atherosclerosis helping keep your blood glucose, also called blood sugar, in the target range. Physical activity and, if needed, diabetes medicines also help.³ One of the risk factor for the development of coronary artery disease is diabetes mellitus is approximately 20% to 30% of the patients undergoing Coronary Artery Bypass Graft (CABG) have diabetes mellitus. There is not enough evidence to determine the impact of DM on short-term mortality and morbidity in the patients undergoing CABG.⁴ Cardiovascular disease

is the most common cause of mortality. Each year it accounts for 30% of deaths worldwide and 38.5% in the United States and Europe. Although the incidence of cardiovascular disease is decreasing in high-income countries due to education and advancement in medical therapy, it is rapidly rising in middle- and low-income countries as they become increasingly industrialized and urbanized.⁵

Methodology

A descriptive study used to achieve the early stated objectives. The study was done from 23rd February up to 23rd July 2016. In order to obtain an accurate data and a representative sample, non-probability a purposive sample of (100) diabetic patients in Ibn Al-Bitar specialized center for cardiac surgery were chosen. The data were collected in the period from 12th May to 12th July 2016. The researcher collected the samples by interview with patients through a special designed questionnaire. This interview took a period of about 10-15 Minutes for each sample. The answers for the third and fourth

part of the questionnaire rated as (3) for always, (2) for sometimes and (1) for never. The questionnaires was constructed and composed of four parts.

Part I: demographic and socio-economic status

Consisted of (6) items which included: age, gender, occupation, marital status, (level of education, and Housing expenses such as family type, home type and have a car) to calculate the economic status.

Part medical information

Second part consists of (11) items. The first item is for Body mass index The second item about smoker, the third item about drinking alcohol, the fourth item about the cumulative rate of sugar HbA1c, the fifth item about the date of diagnosed the diabetes mellitus, the sixth item about the diet without taking treatment, the seventh item about the take tablets for diabetes mellitus, the eighth item about the inject insulin, the ninth item about the take some herbs therapeutic, the tenth item about the history of diagnosis of a blockage in the coronary artery, and the eleven item about the date of coronary artery bypass graft surgery.

Part III: (FFQ) FOOD FREQUENCY QUESTIONERS

This part includes the food frequency Questioners which Consisted of different categories of food that contain food nutrients, the part consists of (9) domains which include (71) items.

Results

(Table 1) The analysis of socio-demographic characteristics shows that the more frequent age group is 60-69 years old among the patients (41%); two third of the sample are male (71%). Regarding socioeconomic status, 78% of them are having moderate level. More than half of the patients are living in a normal marital status, in which the finding shows that 69% of them are married (Table 2).

Table 1 Socio-demographic characteristics of the diabetic patients

No.	Characteristics	F	%
1	Age group	30-39 year	1
		40-49 year	17
		50-59 year	30
		60-69 year	41
		70≤year	11
		Total	100
2	Gender	Male	73
		Female	27
		Total	100
3	Socioeconomic status	Low	5
		Moderate	78
		High	17
		Total	100
4	Marital status	Unmarried	0
		Married	69
		Separated	4
		Divorced	9
		Widowed	18
		Total	100

Table 2 Medical characteristics of the diabetic patients

No.	Characteristics	F	%
1	Body mass index	Underweight	0
		Normal weight	0
		Overweight	28
		Obesity I	54
		Obesity II	17
		Obesity III	1
2	Smoking	Total	100
		Yes	76
		No	24
3	Consuming alcohol	Total	100
		Yes	18
		No	82
4	Cumulative average of blood sugar (HbA1c)	Total	100
		Normal rate	10
		diabetes	28
		High rate	62
5	Diagnosis date of diabetes mellitus	Total	100
		Low	31
		Moderate	32
		High	37
6	Following diet without treatment	Total	100
		Yes	9
		No	91
7	On tablets	Total	100
		Yes	18
		No	82
8	On insulin	Total	100
		Yes	33
		No	67
9	On herbs therapy	Total	100
		Low	54
		Moderate	37
		High	9
10	Diagnosis date of coronary heart occlusion	Total	100
		Before 3 or 8 months	40
		Before 9 or 12 months	32
		More than 12 months	28
11	Date of coronary artery bypass graft surgery	Total	100
		Before 3 or 8 months	40
		Before 9 or 12 months	32

This table reveals that 54% of the patients have obesity class (I) according to body mass index; 76% of them are smoking cigarettes; and only 18% of them are consuming alcohol. The cumulative average of blood sugar test indicates more than seven (62%) among the patients. The date of diabetes mellitus is more than ten years (37%). 91% of the patients do not following a diet without treatment,

73% of them are taking anti-diabetic tablets and 82% of them didn't take an insulin and herbal treatment (67%) and only 33% take herbs as treatment. The diagnosis date of coronary heart occlusion is 1-5 years (54%) and the date of bypass graft surgery is before three or eight months ago (40%) Table 3.

Table 3 Food frequency questionnaire by diabetic patients (N=100)

Food frequency questionnaire		Never	Sometimes	Always	M.S	Severity
Protein	Sheep meat 210g boneless	5	46	49	2.44	H
	Beef meat 210g Boneless	9	71	20	2.11	M
	Chicken 210g Boneless	5	19	76	2.71	H
	Fish 210g	4	43	53	2.49	H
	Eggs 50gm	0	75	25	2.25	M
	Milk all kinds 100ml	47	36	17	1.7	M
	Cream 50gm	69	23	8	1.39	L
	Cooked cheese 250g	4	79	17	2.13	M
Legume	Lentils 90g amount (a small cup)	16	70	14	1.98	M
	Phaseolus 90grams (a small cup)	26	65	9	1.83	M
	Pisum sativum 90g (a small cup)	31	65	4	1.73	M
	Cicer arietinum 90g (a small cup)	8	59	33	2.25	M
	Vigna 90g (a small cup)	39	61	0	1.61	L
	Mung bean 2-6 90g (a small cup)	82	18	0	1.18	L
Grains	Rice 120g	9	56	35	2.26	M
	Bread or loaf (wheat 120g)	5	30	65	2.6	H
	Bread or loaf (hordeum 120g)	5	42	53	2.48	H
	Bulgur 120g	0	75	25	2.25	M
	Macaroni 120g	32	41	27	1.95	M
Nuts	Almonds 90g	0	70	30	2.3	M
	Walnuts 90gm	30	54	16	1.86	M
	Pistacia vera 90g	10	60	30	2.2	M
	Cashew 9 gm	5	65	30	2.25	M
	Corylus 90g	44	40	16	1.72	M
Fruits	Apples 120g medium-sized	4	34	62	2.58	H
	Banana 120g medium-sized	37	51	12	1.75	M
	Orange 120g medium-sized	19	32	49	2.3	M
	Pomegranate 120g medium-sized	10	48	42	2.32	M
	Lemon 120g medium-sized	5	34	61	2.56	H
	Peaches 120g medium-sized	30	53	17	1.87	M
	Citrus reticulate 120 g medium-sized	18	52	30	2.12	M
	Grapefruit 120 medium-sized gm	46	27	27	1.81	M
	Berry 120 grams	0	46	54	2.54	H

Table Continued....

Food frequency questionnaire		Never	Sometimes	Always	M.S	Severity
Vegetables	Tomato (One serving is 250g)	0	26	74	2.74	H
	Potato (250g per meal)	0	40	60	2.6	H
	Cucumber (One serving is 250g)	0	9	91	2.91	H
	Pepper (250g per meal)	5	32	63	2.58	H
	Brassica (One serving is 250g)	19	47	34	2.15	M
	Broccoli (meal per 250g)	70	17	13	1.43	L
	Turnip (One serving is 250 g)	38	49	13	1.75	M
	Carrot (One serving is 250 g)	15	68	17	2.02	M
	Eggplant (One serving is 250g)	4	76	20	2.16	M
	Okra (One serving is 250g)	0	70	30	2.3	M
	Garlic (20 grams per meal)	13	35	52	2.39	H
	Onions (50 grams per meal)	4	29	67	2.63	H
	Marrow (One serving is 250g)	14	56	30	2.16	M
	Cabbage (One serving is 250g)	5	69	26	2.21	M
	Leafy vegetables (basil, mint, , etc) 50 g	0	14	86	2.86	H
Fat	Animal fat (fat free) spoon cup 5ml	66	30	4	1.38	L
	Cups vegetable fat spoon 5ml	83	17	0	1.17	L
	Olive oil cup 5ml	18	58	24	2.06	M
	Palm oil cup 5ml	0	23	77	2.77	H
Fluids	Water 500ml	0	27	73	2.73	H
	Lemon juice 330ml	48	26	26	1.78	M
	Pomegranate juice 330ml	64	31	5	1.41	L
	Orange juice 330ml	51	39	10	1.59	L
	Cocktail juice 330ml	65	26	9	1.44	L
	Cola 330ml	52	35	13	1.61	L
	Seven up 330ml	23	54	23	2	M
	Miranda 330 ml	70	25	5	1.35	L
	Soda330ml	46	37	17	1.71	M
	Diet 330 ml	10	42	48	2.38	H
	Carbonate with caffeine	74	22	4	1.3	L
	Tea 30ml	0	5	95	2.95	H
	Coffee 30ml	10	51	39	2.29	M
	Cocoa 30ml	52	44	4	1.52	L
	Botanical Herbs					
Botanical Herbs	Cinnamon (students) 10g	67	9	24	1.57	L
	Oliban 10g	77	5	18	1.41	L
	Cyperus 10g	62	0	38	1.76	M
	Ginger (race warm), 10g	67	10	23	1.56	L
	Hawthorn 10g	82	4	14	1.32	L
Total					2.04	M

M.S, mean of score; Sig, significance; L, low; Moderate, H, high;

(Low=1-1.66, Moderate=1.67-2.33, High=2.34-3)

This table shows the food frequent intakes by diabetic patients; up the protein nutrients, sheep meat, chicken, and fish are highly significant (M.S=2.44, 2.71, & 2.49) while beef meat, eggs, milk, and cooked cheese are moderately significant except cream which has low significant among them. The legumes are moderately significant in all nutrients except vigna and mung bean which are low significant (M.S=1.61 & 1.18). Regarding grains, most nutrient show moderate significant except bread of wheat and hordeum which are show high significant (M.S=2.60 & 2.48). Nuts' nutrient items are show moderate significant in its entire item. The fruits show moderate significant n most of fruit's type except apples, lemons, and berry are highly significant (M.S=2.58, 2.56, and 2.54). Vegetables reveal that are undertaking moderate to high; tomato, potato, cucumber, pepper, garlic, onions, and leafy vegetables are high significant intake by the patients while turnip, carrot, eggplant, okra, marrow, and cabbage are moderate significant except broccoli is show low significant among them. Regarding fat intake, palm oil is highly intake by the patients (M.S=2.77) and olive oil is moderately intake while animal fat and cups vegetables are low intake. Among fluid, water, diet drink, and tea are highly significant (M.S=2.73, 2.38, & 2.95), lemon juice, seven up, soda, and coffee are moderate significant but the other fluid are low significant. The botanical herbs are rarely intake by the patients which are revealed low significant except Cyperus which is moderately significant among them (M.S=1.76) Table 4.

Table 4 Association between dietary habits, food frequency questionnaire and age group of diabetic patients (N=100)

Age \ FFQ	Moderate	High	Total
30 -39 year	1	0	1
40 -49 year	17	0	17
50 -59 year	26	4	30
60-69 year	41	0	41
70≤year	11	0	11
Total	96	4	100

$\chi^2_{obs}=9.722$ $df=4$ $\chi^2_{crit}=9.49$ $P\leq 0.05$ Sig.= S

This table shows that highly significant differences association between age group with knowledge of the dietary habits and food frequency questionnaire Table 5. This table shows that no significant association between socioeconomic status and dietary habits and food frequency questionnaire Table 6. The table above shows that high significant differences association between marital statuses dietary habits, food frequency which are more frequent among married patients Table 7.

Table 5 Association between dietary habits food frequency questionnaire and socioeconomic status among diabetic patients (N=100)

Socioeconomic \ FFQ	Moderate	High	Total
Low socioeconomic	5	0	5
Moderate socioeconomic	75	3	78
High socioeconomic	16	1	17
Total	96	4	100

$\chi^2_{obs}=0.370$ $df=2$ $\chi^2_{crit}=5.99$ $P\leq 0.05$ Sig.= N.S

Table 6 Association between dietary habits and food frequency questionnaire with marital status of diabetic patients (N=100)

Marital status \ FFQ	Moderate	High	Total
Married	69	0	69
Separated	4	0	4
Divorced	5	4	9
Widowed	18	0	18
Total	96	4	100

$\chi^2_{obs}=42.130$ $df=3$ $\chi^2_{crit}=7.81$ $P\leq 0.05$ Sig.=H.S

Discussion

Part I: Discussion of the socio-demographic characteristics of studied sample

Through the data analysis of distribution of the socio-demographic shows in Table 1 that the more frequent age groups are (60–69) years old among the patients (41%). This results were similar a study done by Pertti et al.⁶ who indicate that (46.9%) of the sample were less than 65 years old. The variables reveal that most of the samples were males (73%). This results were similar to a study done by Pertti et al.⁷ who showed that (79.1%) of patients were males. But these results disagreed with Smeltzer et al.⁸ who stated that women have higher incidence of coronary artery disease and more exposed for CABG surgery. Regarding socioeconomic status, 78% of them are having moderate level (71-86). This results disagrees with a study done by Abbas⁹ who found that (87.8%) of patients were in a low economic status. This also disagrees with Nateghian¹⁰ who reported that patients with low economic status are at risk of coronary artery disease. More than half of the patients are living in a normal marital status, in which the finding shows that 69% of them are married, this result is similar to the study, by Noha et al.¹¹ who found that (84.2%) of the sample were married.

Part II: discussion of the medical characteristics of studied sample.

The medical characteristics of the patients in part two reveal that Table 2 54% of the patients have obesity class one; (30.0-34.9) according to the body mass index this results were similar to a study done by Yap¹² who reported that the age group (60-69) years with the highest prevalence of obesity. The search for smokers is 76% of the total sample result. This findings agrees with Abbas⁹ who found that (80%) of patients were smokers and Wang et al.¹³ who showed that (59.1%) of the sample are smoked. But this result disagrees with Helen where findings uncovered that approximately (88%) of sample were not smokers. Regarding of alcohol only 18% of them consumed alcohol this ratio is acceptable in the Muslim countries compared to other communities in which to disagree from drinking alcohol (the researcher). The cumulative average of blood sugar test (HbA1c) indicates more than seven (62%) among the patients because the glucose level is affected by several factors, such as the psychological situation, nutritional status, and physical factors (the researcher).

Part III: discussion of the food frequent questionnaire characteristics of studied sample.

This part show the food frequent questionnaire by diabetic patients

Table 3 the protein nutrients, sheep meat, chicken, and fish are highly significant with diabetes appears because they are food items to help patients with diabetes and do not affect the muscular effort of the heart and are easily metabolic in the body while beef, eggs, milk, and cheese are relatively acceptable except cream which has low significant among them because it affects directly on the cardiac arteries, as well as diabetes. Reveal that the legumes are moderately significant (acceptable) because it is available it easier for everyone dealt with it does not directly affect the diabetic patients who Coronary Artery bypasses Graft the grains, most nutrient show moderate significant except bread of wheat and barley which are show high significant because the most of the patients who rely on these foods in our daily lives reveal that the nuts' nutrient items are show moderate significant in its entire item because they contain omega-3 ingredients that help improve circulation and give energy to people with diabetes also the fruits show moderate significant in most of fruit's type except apples, lemons, and berry are highly significant most available, and prices suitable for everyone as well as its benefits for patients reveal that Vegetables reveal that are undertaking moderate to high; tomato, potato, cucumber, pepper, garlic, onions, and leafy vegetables are high significant intake by the patients because it is cheap and beneficial for patients this vegetable is unfavorable when people, especially cardiac disease patients while turnip, carrot, eggplant, okra, marrow, and cabbage are moderate significant except broccoli is show low significant among them unknown to most people, and a little availability the fluid, water, diet drink, and tea are highly significant, it does not harm the health of the body and the habits of the people in these areas to allow taking it at any time lemon juice, seven up, soda, and coffee are moderate significant but the other fluid are low significant being hurt directly to patients the fat intake, palm oil is highly intake by the patients and olive oil is moderately intake while animal fat and cups vegetables are low intake, this result is identical to the instructions of the physician of patients with diabetes and patients with C.A.B.G. the botanical herbs are rarely intake by the patients which are revealed low significant except cyperus which is moderately significant among them, Herbs in general is slow and preventive treatments and patients generally need quick fixes hits in addition it is weak in the culture of our society (the researcher).

Regarding Table 4 the dietary habits and food frequency questionnaire are more frequently held by patients with age group (60-69) years; the indicates high significant association between age group and dietary habits, food frequency questionnaire because the old age usually need most to the quality of food will help them to overcome the deficits they face as well as concern for the health after C.A.B.G. (the researcher).

Table 5 shows that dietary habits and food frequency questionnaire are more frequent among those who are moderate level of socioeconomic status; but the table indicates that there is no significant association between socioeconomic status with both dietary habits and food frequency questionnaire (the researcher).¹⁴

The Table 6 indicates that dietary habits and food frequency questionnaire are associated with married patients; the table indicates high significant association between marital status with dietary habits and food frequency questionnaire because the married constitute the majority from among (69 %), the study similar to the Kari et al.¹⁵ who showed that (68.4%) of study sample were married.

Conclusion

Reveals the study sample to those (eating three meals, eating salad

vegetables with every meal, and drink more water) as positive habits moral high food held by diabetic patients the results indicate highly significant differences association between age group with knowledge of the dietary habits. There is indicated that high significant association between marital status and dietary habits high significant differences association between marital statuses food frequency which are more frequent among married patients.

Recommendation

Diabetes self-management education (DSME) and support (DSMS) programs are appropriate for education and support to develop and maintain behaviors that can prevent or delay the onset of diabetes, Diabetes self-management education (DSME) and support (DSMS) programs are appropriate for education and support to develop and maintain behaviors that can prevent or delay the onset of diabetes, Screening for and treatment of modifiable risk factors for cardiovascular disease is suggested follow-up for counseling may be important for health, Perform the A1C test at least two times a year in patients who are meeting treatment goals (and who have stable glycemic control), Activating the role of nutrition Nutrition Research Institute at the National Centers for diabetes about diet programs for patients with diabetes, In adults, a screening lipid profile is reasonable at the time of first diagnosis, at the initial medical evaluation twice yearly.

Acknowledgements

None.

Conflict of interest

The author declares no conflict of interest.

References

1. Kushner FG, O'Gara PT, Ascheim DD, et al. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: a report of the American college of cardiology foundation/American heart association task force on practice guidelines. *Circulation*. 2013;127(4):e362–425.
2. Sarwar N, Gao P, Seshasai SR, Gobin R, et al. Diabetes mellitus, fasting blood glucose concentration, and risk of vascular disease: A collaborative meta-analysis of 102 prospective studies. *Lancet*. 2010;375(9733):2215–2222.
3. Abraham G Thomas. *Food and Nutrition for Nurses Christian Medical College & Hospital*. Ludhiana, Punjab, India: Jaypee Brothers Medical Publishers; 2012. p. 8–12]
4. Koochemeshki V, Salmanzadeh HR, Sayyadi H, et al. The effect of diabetes mellitus on short term mortality and morbidity after isolated coronary artery bypass grafting surgery. *Int Cardiovasc Res J*. 2013;7(2):41–45.
5. John Betteridge and Stephen Nicholls. *Managing Cardiovascular Complications in Diabetes*. USA: John Wiley, Sons; 2014.
6. Pertti Lopenon, Micael Luther, Jan-Ola Wistbaka, et al. Quality of life during 18 months after coronary artery bypass grafting. *Eur J Cardiothorac Surg*. 2007;32(1):77–82.
7. Pertti Lopenon, Micheal Luther, Juha Nissinen, et al. EuroSCORE predicts health-related quality of life after coronary artery bypass grafting. *Interact Cardiovasc Thorac Surg*. 2008;7(1):564–568.
8. Smeltzer SC, Bare B, Hinkle JL. *Brunner and Suddarth's Textbook of Medical - Surgical Nursing*. 12th ed. Philadelphia: Lippincott Williams and Wilkins Company; 2010;28(687):756–786.

9. Abbas B Mohammed. *Assessment of Stressful Life Events of Adult Patients with Ischemic Heart Disease in Baghdad City*. College of Nursing/University of Baghdad, Iraq: Springer; 2012. p. 55–57]
10. Nateghian S. Mental health and stressful life events in coronary heart disease patients and non- patients. *Iranian Journal of Psychiatry*. 2008;3(2):71–74.
11. Noha El Baz, Berrie M, Jitse P, et al. Coronary artery bypass graft (CABG) surgery patients in a clinical pathway gained less in health-related quality of life as compared with patients who undergo CABG in a conventional-care plan. *Journal of Evaluation in Clinical*. 2009;15(3):498–505.
12. Yap C, Mohajeri M, Yii M. Obesity and early complications after cardiac surgery. *Med J Aust*. 2007;186(7):350–354.
13. Wang J, Xiao F, Li Y, et al. Risk factors for operative mortality in 1098 patients with coronary artery bypass grafting surgery: a single center report. *Beijing Da Xue Xue Bao*. 2011;43(1):134–139.
14. Kinley Mc. *Macronutrients: the Importance of Carbohydrate, Protein, and Fat*. Health Center; 2014.
15. Kari Hanne Gjeiilo, Alexander Wahba, Pal Klepstad, et al. Survival and quality of life in an elderly cardiac surgery population: 5-year follow-up. *Eur J Cardiothorac Surg*. 2013;44(1):182–188.