

Insecticidal and mosquito repellency property of essential oil from *hyptis suaveolens* leaves

Abstract

Essential oil was extracted from the leaves of *Hyptis suaveolens* by hydro-distillation by placing 1kg of the washed fresh leaves in round bottom flasks set up with 1liter distilled water. The Mosquitocidal activity, mosquito larvae (larvicidal) and mosquito repellency effect were carried-out using standard modified methods. Percentage yield of the essential oils was gotten as 0.05%. The essential oil showed very strong larvicidal properties against mosquito larvae. There was low mortality rate at lower dose, at a concentration of 6.25ppm the mortality rate was 4% over a period of 6 hours, same concentration at 12 and 24 hours gave mortality rate of 11%. The LC₅₀ and LC₉₀ at this concentration were 25.21 and 302.67 respectively. At a concentration of 12.50ppm a mortality rate of 4% was also recorded at 6 hours, at 12 and 24 hours 13% mortality were recorded, and same LC₅₀ and LC₉₀ with 6.25ppm. At concentration of 25.00ppm, 20%, mortality rate was recorded for 6 hours and 30% for both 12 and 24 hours, the LC₅₀ and LC₉₀ are 9.575 and 352.458 respectively. There were no significant differences at 50, 100 and 1000ppm, the mortality rate was 100% for the period of 6, 12 and 24 hours with LC₅₀ and LC₉₀ of 0.549 and 0.853 respectively. There was no record of death in the control even after 24 hours. It was observed that the amount of air that enters and leaves the cage affect the rate at which mosquitoes were repelled. In a group of mosquitoes kept in a cage with all sides open (ASO) 65% of the mosquitoes were repelled within 30 minutes while for one side open (OSO) and all sides closed (ASC), 73% and 85% of the mosquitoes were repelled within the same time frame. In a second experiment, 100% of mosquitoes were repelled from the surface of rats with shaven skin where essential oil was applied. The essential oil also demonstrated dose dependent mosquitocidal activity with LC₅₀ and LC₉₀ values of 6 and 21ppm respectively.

Keywords: *hyptis suaveolens*, essential oils, mosquitocidal, larvicidal

Volume 5 Issue 1 - 2022

Sheneni Victor Duniya,¹ Bala Christiana Joseph,² Momoh Isaac Sokoato³

¹Department of Biochemistry, Federal University Lokoja, Nigeria

²Department of Biochemistry, Ahmadu Bello University, Nigeria

³Department of Biochemistry, University of Ibadan, Nigeria

Correspondence: Sheneni Victor Duniya, Department of Biochemistry, Federal University Lokoja, Nigeria, Tel +234-80-3351-9009, Email shenenivicto@gmail.com

Received: October 25, 2022 | **Published:** November 04, 2022

Introduction

A synthetic insecticide is a poisonous chemical or mixture of chemicals that is intended to prevent, repel, or kill any insect or pest. However, synthetic insecticides present hazardous impacts far beyond their intended targets. Insecticides have inherent toxicity because they are designed to kill living organisms that are considered "pests", that is any unwanted insect. Many insecticides are known to pose significant, acknowledged health risks to people—including birth defects, damage to the nervous system, disruption of hormones and endocrine systems, respiratory disorders, skin and eye irritations and various types of cancers (Agency for Toxic Substances and Diseases Registry, 1994). Repellents are substances applied to the skin, which prevent insects from biting such surface.¹ An insect repellent is a substance that causes an organism to move away from the odour source, insects perceive the volatile repellents by smell (Luts et al., 2014). Chemical repellents are important in protecting people from blood-feeding insects such as mosquito, ticks, mites, and other arthropods and may therefore also reduce transmission of arthropod borne diseases. N, N-diethyl-3-methylbenzamide (DEET) is one of the most well-known arthropod repellents and has been in the market for almost half a century. DEET is effective against many different blood-sucking arthropods. The protection efficacy depends on the type of formulation, application pattern, species, and feeding behavior of the arthropod. DEET is generally safe for topical use if applied as recommended, although adverse effects such as serious neurologic effects have been reported. Many people consider that DEET and related compounds as a health and environmental hazard. It does not readily degrade by hydrolysis at environmental pHs and has been identified as a ubiquitous pollutant in aquatic ecosystems.

Concern about the deleterious effects associated with synthetic chemicals has revived interest to explore plants as a source of natural insecticides, acaricides, and repellents for medical, veterinary and crop protection use.² Essential oils are volatile natural complex secondary metabolites characterized by a strong odour and have a generally lower density than that of water.³ They are natural volatile mixtures of hydrocarbons with a diversity of functional groups, and their repellent activity has been linked to the presence of monoterpenes and sesquiterpenes.^{4,1} Essential oils are plant products obtained by hydro- distillation or other methods (Luts et al., 2014). This complex of compounds is produced by plants, giving them their characteristic smell and taste, and are usually composed of 20–80 or more substances. Their main components are monoterpenes (C10) and sesquiterpenes (C15) derived from isoprene (Luts et al., 2014). Several monoterpenes have been reported as insect repellents (Luts et al., 2014). There are 17,500 aromatic plant species among higher plants and approximately 3,000 essential oils are known out of which 300 are commercially important for pharmaceuticals, cosmetics and perfume industries. Apart from insecticidal potential they are lipophilic in nature and interfere with basic metabolic, biochemical, physiological and behavioural functions of insects. They are also used as flavour in food products, odorants in fragrances, pharmaceuticals (antimicrobial) and as insecticides.⁴

Plants are endowed with a potential to produce a wide range of allelochemicals that protect the plants from insect-pests.⁵ Mechanisms were developed to repel phytophagous pests, i.e. protective hairy or waxy surfaces, hardened cuticles or tissues or the cellular production of repellent and toxic chemicals. In most cases these protective devices are sufficient to repel generalists among the

phytophagous insects.^{6,7} In developing countries, Lamiaceae have traditionally been used for their insecticidal and repellent properties against several insects' species. Most of them belong to the *Hyptis* genus that includes more than 400 species that grow in the tropical regions of the world, mainly in Africa and America and are highly aromatic plants.^{8,9} *Hyptis suaveolens* (L. Poit) is one of the aromatic and odoriferous important traditional plants belonging to the family lamiaceae, an ethno-botanically important medicinal plant. The plant has been considered as an obnoxious weed. It is a brushy erect plant with fragrant hairy cordate.^{10,8,4,11-17} *Hyptis suaveolens* is a medium aromatic shrub found in the tropics and subtropics, and distributed as the aggressive annual weedy species in the northern part of Thailand. The plant is native to tropical America but now distributed throughout the whole world from tropical to subtropical regions and, therefore, the plant is sometimes regarded as pan-tropical, it is a soft ruderal weed that normally grows along the roadsides and the wet margins of ponds.¹⁰⁻¹⁷

Materials and methods

Plant material

Fresh *Hyptis suaveolens* leaves were used in the experiment for the extraction of essential oil via

hydro-distillation.

Other materials

Anopheles' mosquito larvae were identified and gotten from the drainage in Samaru, Zaria along Sokoto Road, Nigeria. Some of the larvae were allowed to pupate and transform to the adult mosquito.

Chemicals and reagents

All chemicals and reagents were of analytical grade

Equipments

Round bottom flask, water condenser, Erlenmeyer flasks, separating funnel, glass bottles with tight cover, paper tape, Cardboard papers, Aluminium cage with wire gauge, Cage made of Mosquito net with wooden support, plastic containers and petri dishes.

Methods

Plant Material Collection and Identification

The plants were collected from Samaru area of Zaria, Kaduna-Nigeria. They were then taken to the herbarium, department of Biological Science for identification. Voucher specimen number 2020 was deposited. More of the plant leaves were then plucked and taken to the laboratory for further use.

Extraction of essential oil from *hyptis suaveolens* leaves

Hyptis suaveolens (1kg) leaves were washed and placed in round bottom flasks of the hydro-distillation set up with 1liter distilled water. The round bottom flask was then heated on heating mantle for 2 hrs. The essential oil evaporated due to the heat, it was cooled and collected as distillate (mixture of essential oil and water). This was transferred to a glass separating funnel and the essential oil separated from the water based on density. Percentage yield of the essential oil extracted was then calculated using this formula;

$$\% \text{ yield} = x \ 100$$

Mosquito breeding

Some of the collected mosquito larvae were taken to the laboratory washed three to four times and introduced into a clean transparent glass container; the container was placed in a mosquito cage that was

prepared for the experiment. Cracker biscuit powder was sprinkled into the glass container which serves as food for the larvae. The 3rd and 4th instar larvae were collected and used according to WHO 2005 guidelines for the larvicidal bioassay. For mosquito repellency determination, the larvae were left to pupate, transform into adult stage and remain in that same cage for at least 24 hours before introducing them into another cage for repellency experiment to be carried out. They were left for 24 hours prior to the commencement of the experiment so as to be strong to be sure that whatever behavior that will be observed will not be due to their fragility.

Determination of mosquitocidal capability of the essential oil

In this experiment 25 adult mosquitoes each were introduced into a separate mosquito cages and tagged 1-6. Group 1 had no essential oil and therefore served as control, group 2 had 1 ml essential oil, group 3 had 2 mls essential oil, group 4 had 3 mls essential oil, group 5 had 4 mls essential oil while group 6 had 5 mls essential oil dropped into a container in the cages according to the method described by Bulugahapitiya, et al.,¹⁸ The mosquitocidal effects of the essential oil were observed and records taken at the end of 90 minutes in all the groups. Determination of Mosquito Repellency Effect of Essential oil This was carried out by modified method of Bulugahapitiya, et al.¹⁸. Large cages were prepared by introducing 1 ml of the essential oil into each cage containing 25 adult mosquitoes. The behavior of the mosquitoes was then observed and recorded at 30 minutes, 40 minutes, 60 minute and 90 minutes respectively. The control group was set up with pure acetone and no essential oil. The test groups were three; group one was carried out under 100% aeration, group two was done under 20% aeration and group three under 0% aeration. All the above methods were repeated three times.

Determination of mosquito repellency activity of essential oil of *hyptis suaveolens* leaves on rats

Mosquitoes (25 adults) were introduced into each three separate cages grouped I, II and III. Restrained rats (3) each were also introduced into the cages. The rats had their back hairs scrapped off and essential oil (0.5 ml) applied on the bare skin of group one (test rats). The other two groups served as control. Group two (II) had no essential oil applied to the scrapped skins but glycerine. Group three (III) had no essential oil or glycerine applied to their bare skins. The number of mosquito bites were counted at the end of 90 minutes by observing red spot on the scrapped area due to the effect of bite on the rats' bare skin.

Test for larvicidal effect of the essential oil (WHO, 2005)

Essential oil concentrations of 6.25 ppm, 12.50 ppm, 25 ppm, 50 ppm, 100ppm, and 1000ppm were prepared from 1% stock solution. For each of the above concentrations using 25 larvae in their 3rd and 4th instar, numbers of dead larvae were counted after 6 hours, 12 hours and 24 hours. A control group was set up for the same timing without the essential oil.

Results

Extracted essential oil from *Hyptis suaveolens* leaves

The quantity of essential oil obtained from three (3) independent extractions using 1 kg in each case of leaves of *Hyptis suaveolens* is presented in Table 1. An average quantity of 0.4-0.5 mls was extracted and the average yield was 0.1%. The oil is a clear liquid with physical appearance of pale yellow. Mosquitocidal capability of Essential oil

from *Hyptis suaveolens* leaves Essential oil from *Hyptis suaveolens* leaves showed mosquitocidal activity against adult mosquitoes. At the end of 90-minute observation of mosquitocidal effect, all the mosquitoes in different groups except those in control group were not active. In the first cage, that is, the group with 1 ml essential oil 16 % were knockdown (dead) while the remaining 84% were seen resting on the wall, all far from the petri dish where the essential oil was placed. In the group with 2mls essential oil, 72% were inactive far away from the essential oil while 28% were knockdown (dead). As the concentration increases the percentage of dead mosquitoes increases and the inactive ones in all cases were seen far away from the essential oil. At 3mls 36% were knockdown (dead) and 64% weakened while at 4mls, 40% were knockdown (dead) and 60% weakened. At 5mls, 44% were knockdown (dead) while 56% were weakened (not active) Table 2.

Table 1 Extracted Essential oil from *Hyptis suaveolens* leaves

Weight of leaves used obtained	Quantity of oil
One (1) kilogram	0.5 mls
One (1) kilogram	0.6 mls
One (1) kilogram	0.4 mls
Average quantity extracted	0.5 mls
Percentage yield of essential oil extracted	0.05 mls

Table 3 Repellency property of *Hyptis suaveolens* essential oil at 100% aeration

Amount of essential oil (mls)	Time (mins)	Active Mean/SD	Percentage Active (%)	Inactive Mean/SD	Percentage inactive (%)	Mortality (%)
1	30	8.67+1.25 ^b	34.68	16.33+1.25 ^a	65.32	0
1	40	6.67+1.25 ^{ab}	26.68	18.33+1.25 ^{ab}	73.32	0
1	60	5.67+1.25 ^a	22.68	19.33+1.25 ^b	77.32	0
1	90	7.33+0.94 ^{ab}	22.68	17.66+0.94 ^{ab}	70.64	0

Group one (1), N=25.LC 50 = 0.061 and LC 90 = 183.078ppm. Means with different letters are statistically different (p≤0.05)

Table 4 Repellency property of *Hyptis suaveolens* Essential oil 20% aeration

Amount of essential oil (mls).	Time (mins)	Active Mean/SD	Percentage Active (%)	Inactive Mean/SD	Percentage inactive (%)	Mortality (%)	Percentage Mortality (%)
1	30	6.67+1.57 ^b	26.68	18.33+2.05 ^a	73.32	0	0
1	40	3.00+2.16 ^a	12	22.00+2.16 ^b	88	0	0
1	60	0.33+0.47 ^a	1.32	23.33+0.47 ^b	93.32	1.33+0.47	5.32
1	90	0.00+0.00 ^a	0	22.67+0.94 ^b	90.68	2.33+1.33	9.32

Group two (2), N =25. LC 50 = 0.572 and LC 90 = 2.005ppm. Means with different letters are statistically different (p≤0.05).

A third group recorded in Table 5 was done under 0% aeration; the repellency effect was higher in this group. At 30-minute post application 85.32% were repelled from the essential oil, at 40 minutes 97.32% were repelled at 60 minutes there was 100% (88% repelled and

Table 2 Mosquitocidal potential of the *Hyptis suaveolens* essential oil

Amount of essential oil (mls)	Time (mins)	Percentage dead (%)	Percentage inactive (%)
0	90	0	0
1	90	16	84
2	90	28	72
3	90	36	64
4	90	40	60
5	90	44	56

N = 25. LC₅₀ = 6.20 and LC₉₀ = 21.28ppm

Repellency property of *Hyptis suaveolens* Essential oil

The mosquito repellency property of *Hyptis suaveolens* essential oil under 100% Aeration is presented in Table 3. Essential oil from *Hyptis suaveolens* leaves had strong repellency effect against mosquitoes. In this study 1 ml of the essential oil repelled 65.32% of 25 adult mosquitoes within a period of 30 minutes. At 40 minutes 73.32% of the mosquitoes were repelled, at 1-hour post application, 77.32% of the mosquitoes were completely repelled from the source of the essential oil not only were they repelled but were weakened as they were observed to be resting on the wall of the net far away from the source of the essential oil. Table 4 shows the result of group two (2) conducted under 20% aeration. At 30 minutes after application, 73.32% of 25 adult mosquitoes were repelled from the source of the oil. The increase in percentage repellency within the same time frame may be due to level of aeration that is less in this second group which builds up the concentration of the essential oil considering its volatility.

12% dead) repellency and at the end of 90 minutes 100% repellency (84% repelled and 16% dead) was recorded also. In all groups the repelled mosquitoes were weak, unable to fly and resting on the wall of the cage far away from the essential oil.

Table 5 Repellency effect of *Hyptis suaveolens* Essential oil 0% aeration

Amount of essential oil (mls).	Time (mins)	Active Mean/SD	Percentage Active (%)	Inactive Mean/SD	Percentage inactive (%)	Mortality (%)	Percentage Mortality (%)
1	30	3.67+0.47 ^b	14.68	21.33+0.47 ^{ab}	85.32	0	0
1	40	0.67+0.47 ^a	2.68	24.33+0.47 ^c	97.32	0	0
1	60	0.00+0.00 ^a	0	22.00+0.00 ^b	88	3.00+0.00	12
1	90	0.00+0.00 ^a	0	21.00+0.00 ^a	84	4.00+0.00	16

Group three (3), N =25.LC 50 = 0.51 l and LC 90 = 1.178ppm. Means with different letters are statistically different (p≤0.05)

Control groups

Control was set up for each of the three group without essential oil but conditions remain unchanged (aeration of 100%, 20% and 0%) and in all cases the mosquitoes were actively flying through out, the results are shown in Table 6, 7 and 8. At the end of 90 minutes there was 100% repellency in the first group, this has been proven by zero number of bites on shaven dorsal part of rats. In the second group with 0.5mls glycerine on the bare skin there was 46.67% of bites at the end of 90 minutes. In the third group with nothing applied on the bare skin of the rats, 45.33% bites were recorded at the end of 90 minutes. Table 9 shows the summary of this experiment.

Table 6 Repellency property of *Hyptis suaveolens* Essential oil at 100% aeration (control)

Amount of essential oil (mls).	Time (mins)	Percentage Repellency (%)	Percentage Mortality (%)
0	30	0	0
0	40	0	0
0	60	0	0
0	90	0	0

N= 25

Table 7 Repellency effect of *Hyptis suaveolens* Essential oil at 20% aeration (control) Amount of essential oil (mls)

Amount of essential oil (mls).	Time (mins)	Percentage Repellency (%)	Percentage Mortality (%)
0	30	0	0
0	40	0	0
0	60	0	0
0	90	0	0

N= 25

Table 8 Repellency effect of *Hyptis suaveolens* Essential oil at 0% aeration (control) Amount of essential oil (ml)

Amount of essential oil (mls).	Time (mins)	Percentage Repellency (%)	Percentage Mortality (%)
0	30	0	0

Table 10 Larvicidal property of essential oil from *Hyptis suaveolens* leaves

Concentrations (ppm).	Time (hours)	Percentage Mortality (%)	Lc50	Lc90	Regression Equation
6.25	6	4	25.206	302.669	Y=818.6x-161.36
	12	11			
	24	11			
12.5	6	4	25.206	302.669	Y=818.6x-161.36
	12	13			
	24	13			
25	6	20	9.575	352.458	Y=1695.8x-360.07

Amount of essential oil (mls).	Time (mins)	Percentage Repellency (%)	Percentage Mortality (%)
0	40	0	0
0	60	0	0
0	90	0	0

N= 25

Table 9 Summary of repellency effect of *Hyptis suaveolens* Essential oil

Amount of fluid applied (ml)	Bites	Percentage Bites (%)
0.5 essential oil	0.00+0.00 ^b	0
0.5 glycerine	11.67+1.25 ^a	46.67
0	11.33+1.70 ^a	45.33

Larvicidal Property of Essential oil from *Hyptis suaveolens*

Hyptis suaveolens essential oil has shown very strong larvicidal properties against mosquito larvae. There is low mortality rate at lower dosage and not significantly different from each other. At a concentration of 6.25ppm the mortality rate is 4% over a period of 6 hours, same concentration at 12 and 24 hours gave mortality rate of 11%. The Lc50 and Lc90 at this concentration were 25.206 and 302.669 respectively and a regression equation of Y= 818.6x-161.36. At a concentration of 12.50ppm a mortality rate 4% was recorded at 6 hours, at 12 and 24 hours 13% mortality were recorded, same Lc50, Lc90 and regression equation with 6.25ppm. At concentration of 25.00ppm 20% mortality rate was recorded for 6 hours and 30% for both 12 and 24 hours, the Lc50, Lc90 and regression equation are 9.575, 352.458 and Y=0.7264x+0.2465 respectively. There are no significant differences at 50ppm, 100ppm and 1000ppm, the mortality rate was 100% for the period of 6 hours, 12 hours and 24 hours with Lc50, Lc90 and a regression equation of 0.549, 0.853 and Y=0.7264x+0.2465 respectively. There was no record of death in the control even after 24 hours. The summary of the larvicidal properties are recorded in Table 10.

Table Continued..

Concentrations (ppm).	Time (hours)	Percentage Mortality (%)	Lc50	Lc90	Regression Equation
50	12	30	0.549	0.853	Y=0.7264x+0.2465
	24	30			
	6	100			
	12	100			
100	24	100	0.549	0.853	Y=0.7264x+0.2465
	6	100			
	12	100			
	24	100			
1000	6	100	0.549	0.853	Y=0.7264x+0.2465
	12	100			
	24	100			
	6	100			
Control	24	0			
	6	0			
	12	0			
	24	0			

Larvicidal overall Lc 50 = 3.579

Lc 90 = 13.828

Overall Regression equation = $y = 0.2868e^{0.1402x}$

Overall $R^2 = 0.9959$

Discussion

The clear liquid oil with physical appearance of pale yellow is same with all previous work. Essential oil yield of about 0.4-0.6ml/kg (equivalent to 0.1%) got in this studies is same with the work of Raizada¹⁹ who got a percentage yield of 0.1 from the same plant, never the less it is low compared with the work of the following researchers; Iwalokun et al.²⁰ a percentage yield of 0.31 and Okonogi et al.²¹ percentage yield of 0.21; The difference in percentage yield could be due to the moisture level of the leaves and the chemotypic profile of the *Hyptis suaveolens* strains analysed.²⁰ Latitude, altitude, soil composition, climate and genetic composition are factors that have been implicated for chemotype variations in *Hyptis suaveolens* and other species of *Hyptis* as well as other aromatic herbs belonging to the Lamiaceae family.²⁰ Various bioactive compounds were recovered from the essential oil samples of this chemotypes with variations in yield, composition and pharmacological effects.²⁰ Mandal et al.²² got the following percentage yield from the same plant using several solvents which are Steam distillation (yield: 0.24%), petroleum ether extract (yield: 1.6%) and ethanol extract (yield: 2.64%); Bachheti et al.²³ got a percentage yield of 17.44 from seed oil and not leave oil; Shenoy et al.⁸ reported percentage yield of 4.78% for Petroleum ether, 8.52% for Solvent ether, 3.30% for Chloroform, 5.48% for Alcohol and 15.22% for Chloroform water; Gavani and Paarakh¹⁰ got a percentage yield of 4.86% methanol extract. The difference in these studies and the present work could be due to the different solvents used and in one case different part of the plant, apart from the above mentioned factors. Mosquitocidal capability against adult mosquitoes could be due to the presence of terpenes in the essential oil as reported by Bala et al.²⁴ Terpenes are generally known to have insecticidal capability. Earlier work done by Olotuah²⁵ showed that essential oil from *Hyptis suaveolens* leaves has insecticidal activities against several insects. Adda et al.²⁶ also showed that essential oil from *Hyptis suaveolens* leaves has larvicidal activity against some insects, all these could be as a result of terpenes in the plant as reported by Bala et al.²⁴

The strong repellent activity against mosquitoes as seen in this work is still connected to the presence of terpenes. This agrees with the work of Benelli et al.⁹ where the essential oil repels some insects such as *Sitophilus granarius* which kept distance from source of the essential oil that was applied on white man filter paper. As expected

the repellency effect is concentration dependant, this goes in line with the work of Abagli and Alavo² which showed that essential oil from *Hyptis suaveolens* has strong repellency property towards mosquitoes. The higher the concentration the higher the repellency property. In their work 97% of mosquitoes were repelled within a period of 15 minutes which demonstrated same effectiveness as DEET. The overall lethal concentration at 50% (Lc 50) is 3.336ppm and Lc 90 is 0.878ppm. A significant (≤ 0.05) repellent activity with differences in repellency rates is a function of both concentration and observation time, this is in perfect agreement with the work of Conti et al.²⁷ these authors showed that differences in repellency rates, were as a function of both concentration and observation time. Singh et al.¹¹ also showed that *Hyptis suaveolens* has strong ability to repel mosquitoes, these authors proved that plant placement, smoldering (dry and fresh leaves, which gives 90% repellency), spraying (essential volatile oil) and sticks formation of *Hyptis suaveolens* leaves repels mosquitoes strongly due to the presence of intense pungency of the leaves. In their study they were able to prove that *Hyptis suaveolens* is an effective plant to repel mosquito as well as various insects.

Strong larvicidal properties against mosquito larvae due to the presence of terpenes are concentration dependant. Lower concentrations gave lower mortality rates and not significantly (≤ 0.05) different from each other, at higher concentrations mortality rates was higher and there were no significant (≤ 0.05) differences between higher concentrations. These are further confirmed by giving same Lc 50 and Lc 90 for lower concentration (25.206 and 302.669 respectively for both 6.25ppm and 12.50ppm) and same Lc 50 and Lc 90 for higher concentrations (0.549 and 0.853 respectively for 50-1000ppm). The work of Conti et al.²⁷ also showed that there is no significant difference between lower concentrations, at dosages ranging from 250 to 350 ppm, mortality rates were lower and not significantly different from each other. Arivoli and Samuel²⁸ also showed that essential oil of *Hyptis suaveolens* has larvicidal activities against mosquito larvae, which is concentration dependant as was seen in this research work.²⁹⁻³²

Conclusion

Percentage yield of 0.05. This is averagely sufficient using hydro-distillation. Essential oil from *Hyptis suaveolens* leaves has shown

mosquitocidal activity. It has strong repellency effect against adult mosquitoes. It also has very strong larvicidal properties. Since terpenes have been revealed from previous studies to be present in the essential oil from the leaves of this plant (*Hyptis suaveolens*) and the oil has successfully killed and repelled adult mosquitoes and at the same time eliminated mosquito larvae, it then suggests that the activities of the essential oil is as a result of the terpenes contained in the oil. Considering the insecticidal properties of essential oil from *Hyptis suaveolens* leaves, we therefore recommend that the essential oil be used in subsequent work to make soap, perfume and or mosquito repellent popularly known as mosquito coil (right concentration of the essential oil mixed with saw dust or any other dust that can be used). Further studies could be done by extracting the components of the essential oil and test each one of them to know the particular compound(s) that are responsible for the mosquitocidal, larvicidal and repellency properties. This information could help in determining the possibility of using the oil to make insecticides or use it as a component of fumigant to eliminate all unwanted target insects.

Acknowledgments

None.

Conflicts of interest

The author declares there is no conflict of interest.

References

- Traoré-Coulibaly M, Paré-Toél L, Sorgho H. Antiplasmodial and repellent activity of indigenous plants used against malaria. *Journal of Medicinal Plants Research*. 2013;7(42):3105–3111.
- Abagli AZ, Alavo TBC. Essential Oil from Bush Mint, *Hyptis suaveolens*, is as effective as DEET for Personal Protection against Mosquito Bites. *The Open Entomology Journal*. 2011;2011;5:45–48.
- Arun KT, Upadhyay S, Bhuiyan M. A review on prospects of essential oils as bio-pesticide in insect-pest management. *Journal of Pharmacognosy and Phytotherapy*. 2009;1(5):052–063.
- Moreira ACP, Lima EO, Wanderley PA, et al. Chemical composition and antifungal activity of *Hyptis suaveolens* (L.) Poit leaves essential oil against aspergillus species. *Brazilian Journal of Microbiology*. 2010;41:28–33.
- Ramya S, Rajasekaran C, Kalaivani T, et al. Biopesticidal Effect of Leaf Extracts of *Catharanthus roseus* L (G) Don. On the Larvae of Gram Pod Borer - *Helicoverpa armigera* (Hübner). *Ethnobotanical Leaflets*. 2008;12:1096–1101.
- Kiprop KA, Rajab MS, Wanjala FME. Isolation and Characterization of larvicidal components against mosquito larvae (*Aedes aegypti* linn) from calodendrum capense thunp. *Bulletin Chemical Society Ethiopia*. 2005;19(1):145–148.
- Din S, Akram W, Khan HAA, et al. Citrus Waste-Derived Essential Oils: Alternative Larvicides for Dengue Fever Mosquito, *Aedes albopictus* (Skuse) (Culicidae: Diptera). *Pakistan Journal of Zoology*. 2011;43(2):367–372.
- Shenoy C, Patil MB, Kumar R. Wound Healing Activity of *Hyptis suaveolens* (L.) Poit (*Lamiaceae*). *International Journal of Pharm Tech Research*. 2009;1(3):737–744.
- Benelli G, Flamin G, Canale A, et al. Repellence of *Hyptis suaveolens* whole essential oil and major constituents against adults of the granary weevil *Sitophilus granarius*. *Bulletin of Insectology*. 2012;65(2):177–183.
- Gavani U, Paarakh PM. Antioxidant activity of *Hyptis suaveolens* Poit. *International Journal Pharmacology*. 2008;4(3):227–229.
- Singh V, Shrivastava G, Shukla S. Mosquito repellent activity of essential oils of *Hyptis suaveolens*. *Journal of Pharmacy Research*. 2011;4(8):2778–2779.
- Malar RJJT, Sushna SL, Johnson M. Bio-efficacy of the leaves extracts of *Hyptis suaveolens* (L.) Poit against fish pathogens. *International Journal of live science and Pharma Research*. 2012;2(1):128–133.
- Agarwal K, Varma R. Antioxidant activity and phytochemical analysis of *Hyptis suaveolens* (L.) Poit. *Journal of Advanced Pharmacy Education & Research*. 2013;3(4):541–550.
- Noudogbessi JP, Agbangnan P, Yehouenou B. Physico-chemical properties of *Hyptis suaveolens* essential oil. *International Journal Med Aromatic plants*. 2013;3(2):191–199.
- Islam AKMM, Noguchi HK. Plant growth inhibitory activity of medicinal plant *Hyptis suaveolens*: could allelopathy be a cause? *Emir. Journal of Food Agriculture*. 2013;25(9):692–701.
- Umedum NL, Ugochukwu N, Ifeoma PU, et al. The efficacy of *Hyptis suaveolens*: A review of its nutritional and medicinal applications. *European Journal of Medicinal plants*. 2014;4(6):661–674.
- Islam AKMM, Ohno O, Suenaga K, et al. *Suaveolic Acid*: A Potent Phytotoxic Substance of *Hyptis suaveolens*. *The Scientific World Journal*. 2014;6(2014):1–6.
- Bulugahapitiya VP, Arachchige PTK. Mosquito Repellent Compounds from the Seeds of *Nigella sativa* L (Black cumin). Proceedings of the fourth academic sessions. *Technical session*. 2007;5:151–158.
- Raizada P. Ecological and vegetative characteristics of a potent invader, *Hyptis suaveolens* (Poit) from India. 2010;11(2):115–120.
- Iwalokun BA, Oluwadun A, Otunba A, et al. Chemical Composition and Antimicrobial Activity of a New Chemotype of *Hyptis suaveolens* (Poit) from Nigeria. *Current Research Journal of Biological Sciences*. 2012;4(3):265–272.
- Okonogi S, Chansakaow S, Vejabhikul S, et al. Antimicrobial activity and pharmaceutical development of essential oil from *Hyptis suaveolens*. *Acta Hort*. 2005;678:163–169.
- Mandal SM, Mondal KC, Dey S, et al. Antimicrobial activity of the leaf extracts of *Hyptis suaveolens* (L.) poit (2007). *Indian Journal Pharmaceutical Science*. 2010;69:568–569.
- Bachheti RK, Rai I, Joshi A. Physicochemical properties and elemental analysis of some non-cultivated seed oils collected from Garhwal region, Uttarakhand (India). *International Journal of ChemTech Research*. 2013;5:232–236.
- Bala CJ, Shenei VD, Momoh IS. Characterization of essential oil from *Hyptis suaveolens* leaves by gas chromatography-mass spectroscopy and fourier transform infra-red spectroscopy. *International Journal of Molecular Biology*. 2020;5(3):125–133.
- Olotuah OF. Laboratory Evaluation of Pesticidal Activities of *Hyptis suaveolens* in Pest Management. *International Journal of Agricultural Research*. 2013;8:101–106.
- Adda C, Atachi P, Hell K, et al. Potential use of the bushmint, *Hyptis suaveolens*, for the control of infestation by the pink stalk borer, *Sesamia calamistis* on maize in Southern Benin, West Africa. *Journal of Insect Science*. 2011;11:33.
- Conti B, Benelli G, Flamin G, et al. Larvicidal and Repellent activity of *Hyptis suaveolens* (Lamiaceae) essential oil against the mosquito *Aedes Albopictus skuse* (Diptera: culicidae). 2011;110(5):2013–2021.
- Arivoli S, Samuel T. Mosquitocidal activity of *Hyptis suaveolens* (L. Poit) (lamiaceae) extracts against *Aedes aegypti*, *Anopheles stephensi* and *Culex quinquefasciatus* (diptera: culicidae). *International Journal of Recent Scientific Research*. 2011;2(5):143–149.

29. Ghaffari H, Ghassam BJ, Nayaka SC, et al. Antioxidant and Neuroprotective Activities of *Hyptis suaveolens* (L.) Poit. Against Oxidative Stress-Induced Neurotoxicity. *Cell Mol Neurobiol.* 2014;34:323–331.
30. Lutz A, Sfara V, Alzogaray RA. Repellence Produced by Monoterpenes on *Rhodnius prolixus* (Hemiptera: Reduviidae) Decreases after continuous exposure to these compounds. *Journal of Insect Science.* 2014;14(254):1–6.
31. Shaikat MZH, Hossain MT, Azam MG. Phytochemical Screening and Antidiarrhoeal Activity of *Hyptis Suaveolens*. *International Journal of Applied Research in Natural Products.* 2012;5(2):1–4.
32. WHO. *Guidelines for Laboratory and Field Testing of Mosquito Larvicides.* WHO/CDS/WHOPES/GCDPP/2005 (13). 2005. p. 9.