

Contamination at scale: risk amplification, system vulnerabilities, and integrated mitigation in mass food production

Abstract

Mass food production improves food availability, affordability, standardization, and distribution efficiency, but it can also magnify the consequences of a localized food-safety failure. This structured qualitative integrative review examines how biological hazards, chemical contaminants, allergens and emerging contaminants enter, persist and spread through large-scale food-production systems, and how these hazards can develop into high-consequence public-health events. The synthesis draws on 60 peer-reviewed publications from 2020 through September 2026, supplemented by Codex and ISO framework documents for benchmarking established food-safety approaches. The evidence indicates that scale is best understood as a consequence multiplier rather than an independent hazard: large batches, interconnected equipment, persistent environmental reservoirs, cross-contamination, centralized distribution and delayed detection can expand both the number and geographic reach of exposures. Effective mitigation therefore requires a defence-in-depth architecture that integrates supplier assurance, hygienic design, validated process controls, sanitation, allergen management, environmental monitoring, rapid diagnostics, digital traceability, recall preparedness and organizational learning. Emerging technologies, including biosensors, whole-genome sequencing, high-pressure processing, cold plasma, Internet-of-Things monitoring and data-driven analytics, can strengthen these layers, but their evidential maturity and industrial readiness vary and each application requires matrix-specific validation and governance. The review proposes an integrated risk-amplification framework that links hazard introduction, persistence, transfer, batch amplification, distribution, detection delay, exposure and consequence containment while explicitly positioning the framework as complementary to HACCP and food-safety management systems.

Keywords: mass food production, food contamination, foodborne illness, food poisoning, cross-contamination, biofilms, food safety management systems, environmental monitoring, rapid detection, food traceability

Volume 9 Issue 1 - 2026

Loso Judijanto
IPOSS Jakarta, Indonesia

Correspondence: Loso Judijanto, IPOSS Jakarta, Indonesia

Received: September 15, 2026 | **Published:** September 30, 2026

Introduction

Modern food systems increasingly depend on industrialized processing, centralized manufacturing, large production batches, extended ingredient networks, complex logistics, and wide geographical distribution. These arrangements generate substantial economic and social benefits because they allow manufacturers to produce standardized foods at high throughput and distribute them rapidly to large consumer populations. At the same time, however, this architecture changes the consequences of food-safety failure. A microorganism introduced into a small artisanal production process may affect a comparatively limited number of units, whereas contamination introduced into a high-throughput production line can be transferred across equipment, incorporated into large batches, and distributed across multiple geographical areas before the problem is recognized. Consequently, the food-safety challenge associated with mass production is not simply whether contamination occurs but whether production and distribution structures amplify its consequences.^{1,2}

In this review, mass food production is defined operationally as a production-and-distribution configuration rather than by a single numerical threshold. Five dimensions are considered: (1) throughput and batch or lot size; (2) degree of production centralisation and automation; (3) connectivity among equipment, rework streams and successive production runs; (4) breadth and complexity of the supplier

and logistics network; and (5) geographic reach of distribution. An operation can therefore exhibit high amplification potential without being equally large on every dimension. This definition matters because contamination persistence, transfer and detectability differ across product types and process configurations; scale becomes safety-relevant when these dimensions increase the number of units, lots, locations or consumers connected to a single control failure.

The distinction between **hazard generation** and **risk amplification** is central to this review. Large-scale manufacturing does not necessarily generate fundamentally different pathogens from those found elsewhere in the food chain. *Salmonella*, pathogenic *Escherichia coli*, *Listeria monocytogenes*, *Campylobacter*, *Bacillus cereus*, toxin-producing bacteria, allergens, mycotoxins, chemical migrants, heavy metals and other hazards can originate at agricultural, processing, storage, packaging, transportation, retail or food-service stages. What large-scale manufacturing may change is the number of products, production lots and consumers affected once control fails. Shared equipment, conveyor systems, processing water, employee movement, difficult-to-clean niches, recirculated materials and long production runs may connect initially separate contamination pathways into a single amplification network.³⁻⁵

Microbiological hazards remain particularly important when considering acute food poisoning and foodborne disease. Pathogens can survive or adapt within processing environments, adhere to

equipment, establish biofilms, withstand environmental stress, and recontaminate foods after a validated lethality step. *Listeria monocytogenes* is especially relevant to ready-to-eat food production because its environmental adaptability facilitates persistence in processing facilities, while *Salmonella* and other organisms can persist even in relatively dry production environments. Spore-forming organisms such as *B. cereus* present a different challenge because spores may survive processes that effectively control vegetative organisms and subsequently germinate if storage conditions permit growth.^{6–8}

Food contamination nevertheless extends well beyond classical infectious agents. Mycotoxins, pesticide residues, heavy metals, process contaminants, substances migrating from food-contact materials, persistent environmental contaminants, endocrine-disrupting chemicals, PFAS and micro- or nanoplastics increasingly occupy the food-safety literature. Some of these contaminants are associated primarily with chronic or cumulative exposure rather than rapid outbreaks of vomiting or diarrhoea. It is therefore scientifically important not to use “food poisoning” as a universal synonym for every form of food contamination. Acute microbial intoxication, foodborne infection, IgE-mediated allergic reactions and long-term chemical exposure represent different risk pathways requiring different toxicological, epidemiological and regulatory responses.^{9–12}

The growth of global food supply chains creates another layer of vulnerability. Ingredients may pass through multiple countries, companies and intermediaries before entering a final product. Information fragmentation can delay source attribution when a safety incident occurs. Digital traceability, blockchain architectures, smart sensors and integrated data platforms have therefore attracted considerable attention as mechanisms for improving visibility, information integrity and recall speed. Their potential is significant, but traceability itself does not make unsafe food safe. A perfect digital record can rapidly identify a contaminated batch yet does not compensate for inadequate hygienic design, a failed cooking step or ineffective sanitation. Digital technologies should thus be treated as components of a broader prevention and containment system rather than substitutes for preventive control.^{13–15}

Similarly, compliance documents alone cannot guarantee food safety. Hazard Analysis and Critical Control Point approaches, Good Manufacturing Practices, sanitation standard operating procedures, supplier assurance, environmental monitoring and formal Food Safety Management Systems provide an essential organizational structure, but their effectiveness depends on implementation. Research on food-safety culture increasingly demonstrates that leadership, employee behaviour, communication, incentives, training, and organizational priorities influence whether written procedures become routine practice. Production pressure can compete with sanitation time, deviations may go unreported, employees may normalize recurring failures, and managers may underestimate weak signals before a serious event occurs.^{16–18}

This article therefore asks a broader question than “what contaminants can occur in mass-produced food?” It asks how **hazard introduction, persistence, cross-contamination, production scale, distribution scale, and detection delay interact to transform contamination into high-consequence food-safety events**. It further asks which combinations of conventional and emerging controls can interrupt this amplification pathway most effectively.

Accordingly, the objectives are to identify major contamination hazards relevant to mass food production; examine the pathways through which hazards enter, persist, and spread; explain how

production scale magnifies exposure; evaluate conventional food-safety controls; assess emerging detection, processing, and traceability technologies; integrate technical and organizational dimensions; and propose a multilayer framework for prevention and consequence containment. This framing contributes to the literature by treating mass-production food safety as a systems problem rather than a collection of unrelated pathogen-, contaminant- or technology-specific problems.

The proposed risk-amplification framework is not intended to replace established risk analysis, HACCP, prerequisite programmes, food-safety management systems, traceability or recall management. Instead, it adds a consequence-propagation lens: conventional frameworks identify and control hazards, whereas the amplification lens asks how far a failure can persist, transfer, multiply across production, spread through distribution and remain undetected before containment. This distinction is used throughout the review to clarify the framework’s incremental contribution.^{9–21}

Literature review

From contamination to foodborne risk

A hazard is not equivalent to risk. Hazard describes the intrinsic capacity of a biological, chemical or physical agent to cause harm, whereas risk reflects the likelihood and severity of adverse outcomes under actual exposure conditions. The same microorganism can therefore represent different levels of risk depending on its concentration, food matrix, storage conditions, processing treatment, intended consumers and opportunity for growth. Likewise, the same chemical contaminant can create different concerns depending on concentration, exposure frequency, toxicokinetics and duration of consumption.

This distinction is crucial when evaluating mass food production. A large factory does not automatically produce riskier food than a small operation. Large manufacturers may have greater technical expertise, validated processes, automated monitoring and sophisticated laboratories. However, if a critical failure occurs, production volume and distribution reach can enlarge the number of exposed consumers. The relevant conceptual relationship is therefore between **probability of control failure** and **magnitude of consequences conditional on failure**. Formal FSMS implementation can reduce outbreak occurrence, illustrating that industrialization can also provide powerful risk-reduction capabilities when systems are correctly designed and executed.¹

A second distinction concerns time horizon and contamination origin. Acute risks include foodborne infection, microbial intoxication and IgE-mediated allergic reactions that can produce clinical effects soon after exposure, whereas chronic risks involve repeated or cumulative exposure to contaminants such as some heavy metals, PFAS, endocrine-disrupting chemicals or process contaminants. Source-related contamination is introduced with raw materials, water, packaging or environmental inputs; process-related contamination is generated or transferred during manufacturing, for example through thermal reactions, shared equipment, rework, post-lethality exposure or sanitation failures. These categories overlap, but they imply different detection windows, exposure models and control strategies.^{9,12,22}

Biological hazards and acute foodborne illness

Foodborne bacteria differ substantially in ecology, infective dose, toxin production, thermal resistance and environmental persistence. *Campylobacter* is strongly associated with poultry chains and can be

introduced or transferred at multiple points from primary production through slaughter and processing. Control consequently benefits from interventions applied across the chain rather than reliance on a single terminal measure.²³ *Salmonella* similarly possesses multiple persistence mechanisms and can survive in processing environments, including low-moisture environments where sanitation procedures require careful design.⁸

Listeria monocytogenes represents an archetypal environmental-persistence hazard. Its ability to tolerate refrigeration, environmental stress and processing conditions creates concern in ready-to-eat production because contamination after a lethality step may persist until consumption. Stress responses may increase environmental survival, while established biofilms and persistent resident strains can serve as recurring contamination sources. Contemporary studies therefore emphasize environmental monitoring, hygienic zoning, niche identification, effective cleaning and sanitation, and trend analysis rather than sole reliance on finished-product testing.^{6,24,25}

Toxin-mediated food poisoning requires separate consideration. *Bacillus cereus* is ubiquitous, forms spores and produces toxins associated with emetic and diarrhoeal syndromes. The organism can persist despite processing where spores survive and subsequently germinate under favourable time-temperature conditions. The hazard is therefore not controlled solely by initial reduction in vegetative cells; effective management must also consider spore survival, cooling, storage, reheating and opportunities for toxin formation.^{26,7,27} Its ecology, virulence determinants and host interactions reinforce the importance of treating *B. cereus* as more than a minor quality defect.^{28,29}

Antimicrobial resistance further complicates biological food safety. Resistant organisms and resistance determinants can move through food chains and environmental interfaces. Food is not the only driver of antimicrobial resistance, but agricultural production, animal systems, processing environments and food handling can participate in broader transmission networks. This makes surveillance and mitigation a One Health concern extending beyond classical contamination counts.³⁰

Biofilms, environmental reservoirs and persistent contamination

The traditional image of contamination as a transient deposit on a visibly dirty surface underestimates the complexity of microbial persistence. Biofilms consist of structured microbial communities associated with surfaces and extracellular matrices that can increase tolerance to environmental stress and sanitation. In food plants, biofilms may develop on stainless steel, plastics, rubber, drains, joints, conveyor components, fillers, slicers and other difficult-to-clean locations. Established biofilms can intermittently release cells, converting environmental niches into recurring contamination sources.^{3,31}

The problem is not limited to wet facilities. Research on dry-surface biofilms demonstrates that apparently dry processing environments may still contain microscopic surface moisture sufficient to support survival and localized biofilm processes. This is particularly important for low-moisture foods because inappropriate introduction of water during sanitation may itself create additional risk. Consequently, hygienic design and cleaning strategies must be matched to the ecology of the facility rather than mechanically applying wet-cleaning routines across all industries.⁵

Persistent contamination also exposes the limitation of finished-product testing. Even extensive sampling examines only

a small fraction of total production, while microorganisms can be heterogeneously distributed. Environmental monitoring therefore provides information about the conditions under which contamination could occur before contaminated finished products are necessarily detected. Effective Environmental Monitoring Programs require risk-based site selection, appropriate indicators or pathogen targets, suitable methods, trend analysis and predefined corrective-action rules. They should be dynamic systems updated as equipment, products and risk profiles change.⁴

Chemical contamination and emerging hazards

Chemical hazards can enter foods through agricultural inputs, contaminated soil and water, industrial emissions, processing reactions, cleaning chemicals, equipment, packaging and storage environments. Unlike many acute microbiological outbreaks, chemical hazards often involve repeated low-dose exposure, making source attribution and causal assessment more difficult. Recent synthesis of chemical-contamination pathways therefore emphasizes farm-to-fork prevention, monitoring and regulatory surveillance rather than exclusive reliance on end-product testing.²²

Mycotoxins illustrate the interface between biological production and chemical toxicity. Fungal contamination may develop before harvest or during storage, and climate, moisture and temperature influence fungal growth and toxin production. Aflatoxin contamination remains especially important in cereals, nuts and other commodities, while maize can contain multiple mycotoxins across different stages of the food chain. Prevention therefore requires agricultural, drying, storage, testing and supply-chain controls rather than waiting until final manufacturing.^{32–34} Analytical reliability also depends on representative sampling and validated extraction and detection procedures, particularly where multiple or modified mycotoxins may co-occur.³⁵

Process contaminants introduce a different logic. Acrylamide, for example, can form during high-temperature processing of carbohydrate-rich foods. Mitigation must consequently balance microbial lethality, sensory properties and chemical-contaminant formation rather than assuming that “more processing” always means “safer processing.” Product formulation, precursor concentration, time-temperature conditions and technological interventions can influence formation.⁹

Food-contact materials constitute another pathway. Chemical substances may migrate from packaging, equipment or related materials into foods depending on material composition, food properties, temperature, contact duration and barrier characteristics. Even dry foods cannot automatically be assumed to experience negligible migration, and analytical identification remains challenging because food-contact materials can contain complex mixtures, including non-intentionally added substances.^{36,10}

Environmental contaminants further complicate the risk landscape. Heavy metals can move through food webs, creating exposure through contaminated food commodities.³⁷ PFAS are persistent chemicals found in environmental and food-related contexts, with analytical challenges arising from their diversity and low-level occurrence.^{38,39} Endocrine-disrupting chemicals similarly require attention to dietary exposure pathways and chronic toxicological effects.¹²

Micro- and nanoplastics represent a rapidly developing field in which occurrence is better established than many precise human-health risk estimates. Foods may become contaminated from environmental sources, packaging, processing or handling, yet analytical methodologies remain heterogeneous and susceptible to

contamination and measurement bias. The appropriate scientific response is therefore neither to dismiss the issue nor to overstate certainty. Standardized measurement, exposure characterization and mechanistic toxicology remain important priorities.^{12,40,41}

Allergen cross-contact

Food allergens differ fundamentally from microbial hazards because very small quantities may provoke clinically significant reactions in susceptible individuals, while the same material is harmless to most consumers. In mass production, allergen risk is strongly associated with shared equipment, rework, ingredient handling, scheduling, dust transfer, inadequate cleaning, incorrect labels and packaging mix-ups.

Cross-contact control therefore requires mapping allergen transmission vectors across the production environment. Dedicated equipment may be appropriate for some operations, while others rely on validated cleaning, production sequencing, controlled rework, physical segregation and analytical verification. Dzwolak⁴² demonstrates the value of structured cross-contact plans, while Jia and Evans⁴³ emphasize that management failures are often behavioural and organizational as well as technical.

Quantitative allergen risk assessment can improve decision-making by relating unintended allergen presence to exposure and clinically relevant reference doses. Nevertheless, it requires reliable data concerning concentrations, consumption amounts and transfer mechanisms. Precautionary allergen labelling should not become a substitute for effective control because indiscriminate “may contain” statements can reduce the usefulness of risk communication. Inconsistent precautionary labels can confuse both industry and consumers, supporting calls for more harmonized, risk-based approaches.^{44,45}

Positioning the amplification framework relative to established food-safety approaches

Codex HACCP and ISO 22000 provide established architectures for hazard analysis, prerequisite programmes, operational control,

verification, corrective action and management-system governance. The risk-amplification framework proposed here uses these approaches as its baseline rather than as competing models. Its added unit of analysis is the propagation of consequences after a control deviation: how persistence, process connectivity, lot size, distribution reach and detection delay interact to enlarge exposure. In that sense, HACCP focuses primarily on preventing or controlling significant hazards at defined process steps, while the amplification framework makes the downstream consequence pathway explicit and links prevention to surveillance and containment.^{19–21}

Risk analysis, traceability and recall management also remain distinct but complementary. Risk analysis estimates the likelihood and severity of adverse outcomes; traceability reconstructs product and information movement; and recall management removes or contains affected product. The amplification framework connects these functions temporally by identifying where a local initiating event can become a multi-lot, multi-location or population-level event. This makes the framework particularly useful for high-throughput or highly connected production systems in which the probability of an initiating failure may be low but the conditional consequence of an undetected failure can be large.

Accordingly, the framework’s components are defined as follows: hazard introduction is the initial entry or generation of a harmful agent; persistence is survival or retention in a reservoir; transfer is movement to food or another product stream; production amplification is the multiplication of affected units, lots or time periods through batch size and process connectivity; distribution amplification is expansion across locations or customers; detection delay is the interval during which exposure can continue before a signal prompts action; and consequence containment comprises traceability, withdrawal, recall, investigation and learning actions that limit or prevent recurrence. These definitions establish the relationships later depicted in Figure 1.




Figure 1 Contamination-to-Consequences Risk-Amplification Framework.

Methods

This study used a structured qualitative integrative literature review with critical, interpretive and thematic synthesis. The design was selected because the research question spans several evidence domains - food microbiology, toxicology, food engineering, analytical detection, organisational behaviour and supply-chain management - that use heterogeneous methods and outcomes. The purpose was therefore explanatory integration rather than exhaustive study retrieval, pooled effect estimation or meta-analysis. The review is not presented as a systematic review and does not claim PRISMA-style completeness; instead, the revised method makes the evidence assembly and synthesis process sufficiently transparent for readers to judge scope, relevance and limitations.

The evidence audit was updated on 29 September 2026. Information sources included PubMed and targeted DOI/publisher-indexed searches across major journal platforms used by the field, including ScienceDirect, Wiley Online Library, Taylor & Francis, MDPI and Frontiers, supplemented by backward citation checks from high-relevance reviews. The publication window was January 2020 through September 2026, and English-language peer-reviewed journal articles were prioritised. Earlier or non-journal materials were used only when they represented normative frameworks needed to position the analysis, specifically Codex HACCP guidance and ISO 22000.

Searches were conceptually organised in three term clusters. The first captured production context (for example, 'mass food production', 'industrial food processing', 'large-scale food manufacturing', 'high-throughput', 'centralised production' and 'food supply chain'). The second captured hazards and amplification mechanisms (for example, 'food contamination', 'foodborne disease', 'biofilm', 'cross-contamination', 'allergen cross-contact', 'mycotoxin', 'food-contact material', 'PFAS', 'microplastic', 'persistence', 'batch', 'distribution' and 'detection delay'). The third captured controls and technologies (for example, 'HACCP', 'food safety management system', 'environmental monitoring', 'rapid detection', 'biosensor', 'whole-genome sequencing', 'high-pressure processing', 'cold plasma', 'traceability', 'blockchain', 'IoT' and 'food safety culture'). Terms were combined iteratively with AND/OR operators as new mechanisms or technologies emerged during synthesis.

Articles were included when they provided empirical evidence, a contemporary review, or a directly relevant analytical framework for at least one of five domains: hazard architecture; persistence and cross-contamination; production- or distribution-scale amplification; prevention, detection and response; or organisational and supply-chain reliability. Articles were excluded when they were duplicates, published before 2020 (except the two normative framework documents), unavailable in English, lacked sufficient relevance to food-production safety, or addressed a technology without information useful for hazard control, detection, implementation or industrial applicability. The final analytical corpus comprised 60 peer-reviewed journal publications. Codex Alimentarius Commission (2023) and ISO 22000:2018 were used as normative comparators and were not counted among the 60 journal publications.

The synthesis used two-stage thematic coding. First-cycle coding assigned descriptive labels to evidence on hazard entry, source-

related and process-related contamination, environmental persistence, biofilm, allergen cross-contact, process failure, time-temperature control, shared equipment, batch size, production-run connectivity, distribution reach, detection delay, supplier assurance, sanitation, environmental monitoring, culture, diagnostics, genomic surveillance, traceability, recall and organisational learning. In second-cycle coding, related codes were clustered into higher-order themes corresponding to the risk-amplification chain. Papers could contribute to more than one theme, and the synthesis emphasised causal relationships and boundary conditions rather than the frequency with which a term appeared.

Credibility was strengthened through domain triangulation and explicit evidence-maturity distinctions. Claims about persistence were checked against biofilm and environmental-monitoring literature; technology claims were considered alongside food-engineering, analytical-validation and implementation evidence; and established controls were benchmarked against Codex/ISO principles. Emerging technologies were categorised qualitatively as (a) established or demonstrably industrialised for defined applications, (b) promising with pilot or application-specific validation, or (c) predominantly proof-of-concept or research-stage. This prevented technological potential from being presented as equivalent to demonstrated industrial effectiveness and informed the limitations stated in Section 5.10.

Results: Thematic findings

Theme 1: Mass production creates a multi-hazard architecture

The reviewed literature demonstrates that mass food production should not be analysed using a pathogen-only framework. The evidence supports four broad hazard families - biological hazards, chemical contaminants (including process contaminants), allergens, and physical/emerging contaminants - but their risk pathways are not interchangeable. They differ in route of entry, ability to multiply or persist, exposure time horizon, detectability, uncertainty and the point at which prevention, surveillance or corrective action is most effective.

Biological hazards are distinguished by their capacity to survive, multiply or produce toxins. Their risk is strongly affected by time, temperature, water activity, pH and ecological interactions. Chemical contaminants generally do not multiply in food but may persist through processing and accumulate in the diet. Allergens can remain biologically active at very small concentrations and require particularly rigorous cross-contact and labelling controls. Emerging contaminants such as microplastics add analytical uncertainty because measurement methods and health-risk characterization continue to evolve.^{7,12,42}

This heterogeneity suggests that "zero contamination" is an insufficiently precise operational objective. Food-safety systems instead require hazard-specific acceptance criteria, preventive controls, detection strategies and escalation procedures. A factory capable of controlling Salmonella through a validated heat treatment may simultaneously require entirely different controls for allergens, acrylamide or packaging migrants. Table 1 makes these differences explicit and shows why a single amplification mechanism or control hierarchy cannot be applied uniformly to every hazard class.

Table 1 Differential applicability of the risk-amplification framework across hazard families

Hazard class	Typical origin / route	Exposure profile	Scale-amplification mechanism	Priority control logic	Evidence / uncertainty
Biological pathogens / toxins	Raw materials; people; water; environment; post-lethality recontamination	Usually acute; organisms may multiply or produce toxins	Persistence in niches, shared equipment, long runs, large lots, wide distribution, delayed detection	Preventive + surveillance: supplier control, lethality validation, hygienic design, sanitation, EMP, rapid testing	Relatively mature for major pathogens, but organism- and matrix-specific
Food allergens	Ingredient handling; shared lines; rework; packaging or label mix-up	Acute reaction in susceptible consumers; very low doses may matter	Cross-contact across runs; incorrect labels applied to large batches; broad distribution	Preventive + corrective: segregation, sequencing, validated cleaning, label verification, controlled rework, rapid recall	Dose-response and individual susceptibility create residual uncertainty
Source-related chemical contaminants	Agricultural inputs; soil/water; packaging; environmental contamination	Often chronic or cumulative; some acute exceptions	Large sourcing pools and blending can spread low-level contamination across many units	Supplier assurance + surveillance: specifications, sampling, targeted analysis, source reduction	Analytical coverage and chronic exposure attribution may be uncertain
Process-related chemical contaminants	Thermal processing; cleaning chemicals; equipment/material interactions	Often chronic; depends on dose and frequency	Uniform process conditions can generate contaminant across an entire production run	Preventive process optimisation + verification: formulation and time-temperature control, migration testing	Trade-offs may exist between microbial lethality, product quality and chemical formation
Emerging contaminants / novel signals	Environment, packaging, processing or measurement discovery	Often uncertain or chronic; health relevance may still be evolving	High-volume production can broaden exposure, but magnitude is difficult to quantify when methods are not harmonised	Surveillance + research: validated sampling, method standardisation, exposure characterisation	Higher uncertainty; avoid equating occurrence with established health-risk magnitude

Theme 2: Contamination enters through multiple nodes

Raw materials are frequently the first pathway. Agricultural commodities can arrive with microbial pathogens, spores, mycotoxins, chemical residues or environmental contaminants. Animal-derived ingredients may carry zoonotic pathogens, while dry ingredients can introduce organisms into facilities where subsequent wet cleaning may be undesirable. Supplier approval, specifications, certificates, testing and risk-based verification therefore represent the first layer of industrial protection.

However, incoming ingredients are only one node. Water, air, employee movement, utensils, maintenance activities, pests, packaging materials, lubricants and equipment can introduce hazards after ingredients have passed initial controls. Even validated thermal processing can be defeated by post-process contamination if exposed ready-to-eat products encounter contaminated surfaces or environments. This observation shifts attention from individual critical control points to the connectivity of the whole plant.^{4,25}

The processing environment is especially important when contamination becomes persistent. Biofilms, drain reservoirs, damaged seals, hollow rollers, cracked surfaces and inaccessible equipment components can create niches from which microorganisms repeatedly enter production. The significance of these niches is disproportionate because they can contaminate successive production lots. One contaminated ingredient batch may be removed; a persistent environmental reservoir can contaminate products repeatedly until the underlying source is found.^{3,5,31}

Theme 3: Production scale operates as a consequence multiplier

The central finding of this review is that scale should be conceptualized primarily as a **risk amplifier**, not automatically as

a hazard generator. Several mechanisms explain this amplification. First, large batch sizes increase the amount of product associated with a single control failure. If contamination occurs before blending or mixing, distribution within a large batch can expose far more units than a small isolated process. Second, continuous production lines create temporal connectivity: equipment contamination can persist across hours or production shifts. Third, shared equipment creates product connectivity because contamination or allergens can transfer from one product to another. Fourth, centralized distribution creates geographical connectivity, allowing contaminated products to reach consumers in widely separated locations. Fifth, detection delay increases temporal exposure because product may continue to leave the facility while investigation is ongoing.

These mechanisms can interact. A persistent niche on shared equipment can contaminate multiple production runs, which are subsequently distributed through a national supply network before epidemiological signals trigger investigation. The problem therefore becomes multiplicative rather than additive.

Within the framework, the relationships are directional but not strictly linear. Persistence increases the probability of repeated transfer; transfer can connect products and production runs; batch size and run duration enlarge the number of affected units; distribution reach converts production amplification into geographic amplification; and detection delay determines how long these processes continue before intervention. Feedback is also possible: surveillance or traceability can shorten detection delay, while root-cause analysis can reduce future persistence or transfer. The framework therefore represents a causal organising map rather than a single deterministic sequence.

Figure 1 conceptualizes this process as an amplification chain. Its purpose is to distinguish the initial contamination event from

the downstream mechanisms that determine how severe the event eventually becomes.

Figure 1 separates initiating hazard control from consequence propagation. The upper modifiers - throughput, process connectivity, centralisation, supply-chain breadth and distribution reach - do not create a hazard by themselves, but they can increase the conditional consequence of an undetected failure. The lower control bands show where established HACCP/FSMS prevention and verification act, and where surveillance, traceability, recall and organisational learning limit consequences. This makes the framework complementary to established food-safety systems rather than a replacement for them.

Table 2 Applied examples of amplification pathways and interruption points

Applied example	Initiating mechanism	Amplification pathway	Critical interruption points
Ready-to-eat chilled food: persistent <i>Listeria</i>	Post-lethality environmental reservoir / niche	Persistence -> repeated transfer -> multi-lot production -> wide distribution -> delayed detection	Hygienic design; sanitation; risk-based EMP; rapid confirmation/WGS; lot traceability; recall
Shared-line allergen / mislabelling	Cross-contact during changeover or wrong packaging/label	Transfer or label error -> large batch affected -> broad distribution; no organism growth required	Segregation; sequencing; validated cleaning; line clearance; label verification; traceability; rapid recall

Theme 4: Persistence converts episodic hazards into systemic hazards

Biofilms and environmental persistence are among the most important transitions in the amplification pathway. A transient contamination event can theoretically disappear after cleaning or removal of the affected product. Once organisms become established in persistent niches, however, the hazard changes from episodic to systemic.

Contemporary biofilm research shows why routine sanitation may fail. Extracellular matrices can reduce penetration of antimicrobial agents, microorganisms within biofilms may display altered physiology, and surface properties influence adhesion. Mixed-species biofilms can add further complexity. The growing literature on *L. monocytogenes* emphasizes that eliminating mature environmental reservoirs may require a combination of mechanical cleaning, validated chemical sanitation, hygienic redesign, intensified environmental sampling and root-cause investigation.^{24,25} Because mature biofilms can show reduced susceptibility to disinfectants, hurdle approaches that combine complementary physical, chemical and process interventions may provide more robust control than reliance on a single antimicrobial mechanism.⁴⁶

Salmonella persistence similarly illustrates why low-moisture environments cannot be considered intrinsically safe. Organisms may survive for extended periods under dry conditions, while introduction of moisture can support localized growth. Pang et al.⁸ therefore emphasize multiple strategies for controlling resistant biofilms rather than dependence on a single sanitizer. Environmental monitoring becomes particularly valuable in this context because it converts invisible conditions into actionable information. Trends in indicator organisms or repeated pathogen findings in particular zones can reveal deteriorating hygienic control before finished-product positives become frequent. However, an EMP is useful only when results trigger meaningful investigation; repeated positives that are merely “cleaned and re-tested” without source elimination can normalize failure rather than control it.⁴

Theme 5: Human and organizational factors determine whether controls function

Food-safety plans operate through people. Training can improve knowledge and practices, but knowledge does not automatically

Two applied examples illustrate the framework’s differential utility. In a ready-to-eat chilled-food facility, a persistent *Listeria* niche can repeatedly transfer cells after a lethality step, contaminate multiple lots and reach geographically dispersed consumers before environmental or epidemiological detection. In contrast, an allergen event may begin with an incorrect label or inadequate changeover; no microbial growth is required, yet one packaging error can expose an entire batch and make traceability and rapid market withdrawal decisive. Table 2 maps these examples to the amplification chain.

produce safe behaviour, particularly when organizational incentives reward speed, yield or output more strongly than careful compliance. Meta-analytic evidence shows benefits from food-safety and hygiene training, yet sustained behavioural change requires reinforcement and appropriate organizational conditions.⁴⁷

Food-safety culture literature extends this observation. Zanin et al.¹⁶ found value in combining qualitative and quantitative approaches because formal survey responses can overestimate actual culture. Recent reviews further emphasize leadership, organizational commitment, communication and multicultural workforce considerations as determinants of how food-safety expectations are interpreted and enacted.¹⁸ Operationalization is particularly important because “strong food-safety culture” must translate into observable actions such as stopping production during deviations, reporting near misses, following hand-hygiene rules, respecting allergen changeover procedures and allowing sufficient sanitation time.¹⁷

The significance for mass production is substantial. High-throughput systems frequently operate under time and performance pressure. Small deviations can be repeated hundreds or thousands of times before detection. Consequently, organizational tolerance of minor deviations may become a mechanism of risk amplification.

Theme 6: Detection speed determines how long amplification continues

Traditional culture methods remain important because they provide trusted microbiological confirmation, but the time required for enrichment, isolation and identification can conflict with the speed of modern production and logistics. By the time a conventional result is available, large quantities of product may have been manufactured or distributed.

Rapid detection technologies attempt to reduce this information gap. Portable biosensors, nucleic-acid amplification, immunological platforms, microfluidics and point-of-care systems can shorten analytical turnaround and support decentralized testing. Microfluidic lab-on-chip platforms are particularly relevant because they can integrate sample handling and detection into miniaturized systems, although matrix effects, enrichment requirements and industrial validation remain important constraints.⁴⁸ More generally, rapid methods still face limitations related to sample preparation, detection

limits, cost, method validation and implementation at industrial scale.^{49–51}

Biosensor research targeting *Listeria*, *Salmonella*, *Campylobacter* and *E. coli* illustrates the transition toward more rapid and potentially on-site analysis.⁵² High-throughput sensing platforms further offer the possibility of simultaneously examining multiple targets or large sample numbers,⁵³ while CRISPR-based and other advanced molecular approaches are increasingly discussed as components of future detection systems.⁵⁴ Whole-genome sequencing adds a complementary surveillance capability by enabling high-resolution pathogen characterization, source tracing and reconstruction of transmission pathways during food-safety investigations.⁵⁵

Predictive microbiology provides a complementary function. Instead of identifying contamination after it occurs, mathematical modelling estimates microbial responses to environmental conditions. Incorporating food microstructure and matrix effects remains an important development because microbial behaviour in actual foods cannot always be predicted adequately from simplified laboratory media.⁵⁶ Figure 2 positions detection within a broader surveillance architecture. The key concept is that detection should occur at multiple points rather than being concentrated exclusively at finished-product release.




Figure 2 Multi-Layer Surveillance Architecture for Mass Food Production.

Figure 2 illustrates the difference between testing and surveillance. Testing is an individual measurement; surveillance integrates repeated measurements into a decision system. The greatest value of rapid diagnostics therefore emerges when laboratory information is

connected with process history, environmental trends, supplier data and distribution records rather than treated as an isolated result.

Theme 7: Emerging processing technologies add barriers but require validation

Nonthermal and novel processing technologies can strengthen pathogen control while preserving sensory and nutritional quality, but evidence maturity differs markedly by technology and application. High-pressure processing has demonstrated commercial industrial use for selected refrigerated and ready-to-eat products, whereas cold plasma remains more application-specific and sensitive to treatment geometry, food composition, oxidation, sensory effects and regulatory requirements. Therefore, technological capability should be distinguished from validated process efficacy in a defined product and from routine industrial usefulness.^{57–59}

The same distinction applies to monitoring technologies. Whole-genome sequencing is already established in many surveillance and source-attribution contexts, while biosensors, CRISPR-enabled detection, AI-supported analytics, smart packaging and IoT architectures span a continuum from laboratory proof-of-concept to pilot deployment and routine use. Recent reviews emphasise continuing constraints involving matrix interference, calibration, reproducibility, standardisation, interoperability, cost and field robustness. Accordingly, this review treats these technologies as enabling layers whose practical value depends on validated performance and integration into decision processes, not as stand-alone guarantees of food safety.^{60,61} The broader lesson is that emerging technology adds barriers to food-safety systems; it does not remove the need for supplier assurance, sanitation, hygienic design, storage control, competent personnel or management discipline. A technology should be adopted only when its target hazard, decision threshold, validation status and role in the control architecture are explicit. This qualification avoids conflating technological promise with demonstrated reduction in industrial risk.

Theme 8: Traceability limits consequences after prevention fails

Food traceability historically focused on identifying product movement between suppliers, processors and customers. Digital technologies increasingly extend this function by integrating sensor data, cloud platforms, IoT devices, immutable records and potentially genomic information. Blockchain research emphasizes transparency, security and tamper resistance, but implementation must be evaluated against cost, interoperability, governance and the quality of input data. Blockchain cannot correct inaccurate information entered into the system. Therefore, its value depends on integration with reliable identification, sensor and verification systems.^{13–15}

Smart traceability provides a broader concept. Yu et al.² describe systems combining portable detection, intelligent indicators, sensors, IoT and data-assisted methods. Within the risk-amplification framework, the major benefit is reduction of **time-to-intervention**. If contaminated products can be identified and geographically located rapidly, recalls can become more targeted and faster, reducing the duration and size of population exposure.

Figure 3 integrates preventive, detective and responsive controls into a defence-in-depth structure. The arrangement emphasizes that prevention, verification, response and organizational learning should operate as mutually reinforcing barriers rather than as isolated programs.



Figure 3 Defence-in-Depth Architecture for Contamination Mitigation.

The figure demonstrates why food safety should not depend on one “perfect” critical control point. Each layer addresses different failure modes. Supplier controls reduce hazard entry; hygienic design reduces persistence; processing reduces hazard concentration; monitoring reveals loss of control; traceability limits downstream exposure; and root-cause analysis prevents recurrence. The architecture becomes resilient when failure of one barrier is detected or compensated for by another.

Discussion and analysis

From contamination control to amplification control

The principal conceptual implication is not that conventional food-safety thinking should be replaced, but that consequence propagation should be made more explicit within it. Established risk analysis, HACCP and FSMS approaches are designed to identify hazards, establish controls, verify performance and manage deviations. The amplification lens adds a systems-level question: conditional on a control failure, which features of the production and distribution architecture determine how many lots, locations and consumers can become connected to that failure before detection and containment? This emphasis is especially relevant when low-frequency failures can have high-consequence outcomes.¹⁹⁻²¹

A simplified conceptual expression can therefore be proposed to summarize the interacting dimensions of amplification risk. The expression is intended as a conceptual organizing device rather than as a validated quantitative risk model.

$$R_{\text{sys}} \propto P(H) \times I_{\text{pers}} \times I_{\text{trans}} \times S_{\text{batch}} \times R_{\text{dist}} \times D_{\text{det}} \times V_{\text{pop}} \quad (1)$$

In Equation (1), R_{sys} denotes systemic food-safety risk; $P(H)$ represents the probability of hazard occurrence; I_{pers} and I_{trans} represent persistence and transfer intensities; S_{batch} represents batch-scale exposure; R_{dist} represents distribution reach; D_{det} represents detection delay; and V_{pop} represents population vulnerability. The proportionality sign is used deliberately because the expression is conceptual and does not imply that the factors have been empirically calibrated or are dimensionally equivalent. Preventive controls, sanitation, hygienic zoning, production planning, distribution architecture and rapid diagnostics can each reduce one or more components of the amplification pathway.

The model helps explain why rare failures can produce major events. Industrial food safety cannot be judged solely by the frequency of deviations. A very low-probability event can justify substantial investment when consequence magnitude is extremely large.

Prevention and consequence containment are different functions

The literature supports a useful distinction between **prevention controls** and **consequence-containment controls**. Supplier assurance, hygienic design, validated lethality treatments, allergen segregation, sanitation and preventive maintenance seek to stop unsafe food from being produced. Environmental monitoring and rapid detection identify emerging loss of control. Traceability, recall and outbreak investigation limit consequences once unsafe products may have entered commerce.

Organizations frequently prioritize prevention and underestimate containment because recall systems are activated only during crises. Yet mass production creates circumstances in which no preventive system can be assumed infallible. Recall readiness should therefore be tested in the same manner as emergency-response plans in other high-reliability industries.

Digital traceability can shorten the interval between problem identification and market withdrawal. However, effective recall also requires accurate lot coding, distribution records, communication protocols, decision authority and coordination with regulators and customers. A technologically sophisticated blockchain without accurate lot segregation offers limited practical protection.

HACCP/FSMS as the baseline, amplification analysis as a complementary lens

HACCP remains foundational because it directs attention toward significant hazards, preventive controls, critical limits, monitoring and corrective action. ISO 22000 and broader FSMS architectures extend this logic through prerequisite programmes, communication, verification, management responsibility and continual improvement. The proposed amplification framework begins from these established controls and asks an additional question: what is the consequence trajectory if a control is missed, bypassed, defeated by an environmental reservoir, or detected only after product release?

That additional question is important because important failure modes may be distributed across organisational and supply-chain boundaries rather than concentrated at a single critical control point. Environmental persistence, allergen changeover, packaging accuracy, upstream chemical contamination, hygienic design, data integrity and recall readiness can all shape the scale of consequences. These functions are already compatible with modern FSMS; the contribution of the amplification lens is to connect them within one consequence-propagation model rather than to establish a parallel compliance system.

Accordingly, the framework should be used as a stress-testing layer during hazard analysis, management review and recall preparedness. A facility can ask which single failure could contaminate many units, which persistent source could affect multiple runs, which data delay could allow continued release, and which distribution pattern would make a recall difficult. Evidence that formal FSMS implementation can reduce outbreak occurrence supports retaining established management architecture while strengthening attention to consequence connectivity.^{1,21}

Environmental monitoring should function as an early-warning system

EMP design should move beyond periodic compliance swabbing toward risk-based surveillance. High-risk sites should be selected according to product exposure, traffic patterns, hygienic zoning and historical findings. Trend analysis is at least as important as individual results. An increase in indicator organisms or repeated positives near a high-care area may represent a deteriorating system even when finished-product tests remain negative.

The literature also implies that “seek and destroy” philosophies are preferable to cultures in which positive environmental results are discouraged. If employees fear that finding contamination will be treated as personal failure, organizations inadvertently incentivize under-sampling and superficial investigation. Strong food-safety culture instead treats early detection as evidence that the monitoring system is functioning.

This illustrates the interaction between technical and cultural controls. Sampling plans are scientific instruments, but the quality of surveillance depends on organizational willingness to discover uncomfortable information.

Allergen management demonstrates the importance of cross-system thinking

Allergens reveal limitations of microbial analogies. A validated cooking process that eliminates bacterial hazards does not necessarily remove allergenicity. Therefore, allergen control is driven more strongly by segregation, changeover cleaning, formulation control, rework management and labelling accuracy.

Mass production can amplify allergen risk through shared equipment and packaging lines. A single incorrect packaging roll or formulation error may mislabel a large quantity of otherwise microbiologically safe product. Cross-contact assessment must therefore extend through equipment, personnel, airborne powders, scheduling and supply-chain information.^{42,43}

Quantitative risk assessment offers a route toward more proportionate management because it attempts to relate unintended presence to clinically meaningful exposure rather than treating every detectable molecule as equivalent. Nevertheless, uncertainties remain, and precautionary labelling practices require harmonization to avoid both under-warning and excessive warning.^{44,45}

Chemical risk requires a longer time horizon

Mass-food-safety discussions often focus on outbreaks because outbreaks produce visible clusters of illness. Chemical risk frequently develops differently. Exposure may be widespread but low-level, consequences may be delayed, and individual cases may not be easily linked to a single meal or product.

The appropriate control architecture therefore differs. Mycotoxin prevention requires field and storage interventions; packaging migrants require material compliance and migration testing; heavy

metals require environmental and ingredient monitoring; PFAS requires sophisticated targeted and non-targeted analytical strategies; and microplastics require improved methodology before precise quantitative health-risk estimates can be made.

A mature food-safety system should consequently distinguish acute incident management from long-term exposure management while connecting both within an integrated chemical-risk surveillance program. This distinction is especially important because chronic chemical exposure may require longitudinal monitoring, source reduction and regulatory intervention even when no discrete outbreak is observed.²²

Technology should shorten the food-safety information cycle

The most important strategic contribution of new detection technology may be understood through the concept of the **information cycle**. Traditional systems often follow the sequence: sample → transport → enrichment → laboratory analysis → reporting → investigation → intervention. Each stage takes time. In a high-throughput factory, production continues while information is being generated.

Portable biosensors, microfluidics and molecular methods attempt to reduce this delay, but their strategic value depends on where they sit on the evidence-maturity continuum. A fast assay that remains matrix-sensitive or insufficiently validated may be useful for screening but not for final release decisions; conversely, mature laboratory or genomic methods may support definitive confirmation and source attribution even if they are not instantaneous. The relevant performance question is therefore whether a technology shortens the actionable information cycle without sacrificing analytical reliability, interpretability or governance.

However, rapid results that are inaccurate can cause unnecessary recalls or false reassurance. Industrial implementation therefore requires analytical validation, quality assurance, matrix-specific evaluation and predefined decision thresholds. “Faster” must not be confused with “better” unless sensitivity, specificity and operational reliability are appropriate.

Building a digitally enabled food-safety system

Digital food safety should be designed as an information architecture rather than as a collection of fashionable technologies. Sensors can capture temperature, humidity and other process parameters; laboratory systems can capture analytical results; enterprise systems contain supplier and production information; traceability systems record lot movement; and external surveillance may identify outbreak signals. Artificial intelligence and machine-learning tools can support anomaly detection, predictive control and supply-chain traceability, but their value depends on data quality, governance and context-appropriate validation.⁶² The central challenge is therefore to connect these data streams so that meaningful anomalies generate timely, reviewable action.

Figure 4 presents a conceptual architecture for digitally enabled contamination prevention and response. It shows how heterogeneous data streams can be integrated into analytics, expert review, corrective action and organizational learning without eliminating human accountability. The central feature of Figure 4 is the human decision layer. Automation can identify deviations, but governance remains necessary to determine escalation thresholds, evaluate uncertainty and decide whether production should stop or products should be recalled. Digitalization therefore complements rather than eliminates professional judgement.




Figure 4 Integrated Digital Food-Safety Intelligence Architecture.

Implications for developing economies

The proposed model is especially relevant to rapidly industrializing food systems. Developing economies may experience simultaneous growth in industrial processing, modern retail, central kitchens, catering systems and long-distance distribution while laboratory capacity, surveillance infrastructure and regulatory enforcement develop unevenly. Importing the most sophisticated technology is not necessarily the optimal first investment. A facility with weak sanitation or poor equipment design may gain more safety from correcting fundamental hygienic failures than from installing blockchain infrastructure. Prioritization should therefore follow risk hierarchy.

A pragmatic sequence begins with safe water, sanitation, hygienic design, supplier controls, basic process validation, training and management accountability. Environmental monitoring and rapid detection should then be strengthened according to risk. Digital traceability becomes particularly valuable as distribution systems expand. Advanced analytics and novel processing technologies can be adopted where they address clearly identified residual risks. Resource limitations also increase the importance of prevention because large recalls can impose severe economic losses on firms and public authorities. Investments in basic food-safety systems should therefore be regarded not merely as compliance expenditure but as risk-reduction capital.

Limitations of the review and proposed framework

This review has several limitations. First, it is a structured qualitative integrative review rather than a systematic review; although the revised search process is transparent, the iterative and concept-driven approach does not guarantee exhaustive retrieval and may favour well-developed literatures or highly cited review articles.

The English-language and 2020-September 2026 publication window may also exclude relevant earlier, regional or non-English evidence. The final corpus of 60 peer-reviewed publications should therefore be interpreted as a purposive contemporary evidence base rather than a complete census of food-safety research.

Second, the evidence is heterogeneous in design, food matrix, geography and technological maturity, which limits direct comparison across hazards. Biological outbreaks, allergen events and chronic chemical exposure do not share a common time scale or endpoint, and emerging contaminants may have substantial measurement uncertainty. Similarly, evidence for high-pressure processing or genomic surveillance is more mature in defined applications than evidence for many AI-enabled, CRISPR, smart-packaging or next-generation biosensor applications. The review therefore avoids treating technological potential as equivalent to demonstrated industrial effectiveness.

Third, the risk-amplification framework and Equation (1) are conceptual rather than empirically calibrated. The variables are not assumed to be dimensionally equivalent or multiplicatively independent, and no universal threshold is proposed for what constitutes ‘mass’ production. The framework should be used to structure comparative reasoning, stress testing and research design; quantitative validation is still required before it can support prediction, scoring or regulatory decision thresholds.

Future research agenda

Several research priorities emerge from the synthesis. First, future studies should develop quantitative models linking production scale to outbreak consequences. Existing microbial risk assessments often model dose and exposure, but greater attention could be paid to connectivity among equipment, batches and distribution networks.

Second, multi-hazard frameworks are needed. Food-safety scholarship remains strongly compartmentalized between microbiology, allergens, toxicology and packaging science. Industrial managers, however, operate one production system in which these hazards coexist.

Third, more research is required on environmental-persistence thresholds and predictive indicators. Monitoring programs generate large quantities of data, but methods for converting repeated low-level findings into actionable early warnings require refinement. Fourth, rapid diagnostics need stronger evidence under real industrial conditions. Laboratory detection limits do not automatically translate into reliable performance in complex food matrices. Research should therefore emphasize sample preparation, enrichment reduction, multiplexing, field robustness and economic feasibility.

Fifth, digital technologies require evaluation based on measurable safety outcomes rather than adoption alone. Studies should test whether IoT, blockchain or AI actually shorten recall times, improve source attribution, reduce contamination or prevent outbreaks. Sixth, emerging contaminant research should improve standardization. Micro- and nanoplastic studies particularly require harmonized sampling, contamination control, extraction and measurement methods before robust comparisons across foods and populations are possible.

Finally, organizational research should investigate how production incentives influence food-safety decisions. Technical failures often become serious because employees, supervisors or managers respond inadequately to weak signals. Food-safety culture should therefore be studied as a causal component of system reliability, not merely as an employee-attitude survey.

Conclusion and recommendations

Mass food production is not inherently unsafe. Industrial facilities can deploy process validation, automation, monitoring and professional food-safety management at a level that may be difficult to reproduce in fragmented production. The synthesis nevertheless shows that high throughput, process connectivity and broad distribution can increase the conditional consequences of a control failure, particularly when persistence, cross-contact or detection delay connect many units or production periods to the same initiating event.

The analysis demonstrates that biological contamination remains the dominant concern for acute outbreaks and conventional food poisoning, but a comprehensive food-safety strategy must also incorporate allergens, mycotoxins, processing contaminants, packaging migrants, heavy metals, PFAS and emerging contaminants. These hazards differ in exposure mechanisms and health consequences, making hazard-specific controls essential.

The most effective strategy is therefore defence in depth. Prevention should begin with supplier assurance and hygienic design and continue through validated process controls, sanitation, allergen management and environmental monitoring. Rapid diagnostics should shorten the time required to identify emerging problems, while traceability and recall systems should minimize population exposure when preventive controls fail. Novel technologies such as cold plasma, high-pressure processing, advanced biosensors and digital traceability can strengthen the system but should not substitute for fundamental hygienic and managerial controls.

Food-safety culture forms the organizational infrastructure connecting these technical layers. Employees must be able and

expected to report deviations, halt unsafe production and escalate abnormal findings. Management should reward early detection rather than concealment of problems.

The central recommendation is therefore to complement contamination-control practice with explicit amplification analysis. Food producers and regulators should evaluate not only whether a hazard can enter or arise in food, but also whether it can persist, transfer across production, affect multiple lots, travel through distribution networks and remain undetected long enough to expand exposure. This lens is designed to sit within HACCP and FSMS processes, strengthening stress testing, surveillance design, traceability and recall preparedness rather than replacing established food-safety frameworks.

Future food-safety systems should increasingly combine preventive engineering, microbial ecology, toxicological surveillance, validated rapid analytics, digital traceability and organisational learning. However, technology choices should follow evidence maturity and application-specific validation: commercially established controls and surveillance tools should be distinguished from promising pilot systems and research-stage concepts. The proposed framework is therefore best interpreted as a conceptual organising tool for prevention and consequence containment, with quantitative calibration and comparative validation remaining priorities for future research.

Acknowledgements

None.

Conflicts of interest

The author declares there is no conflict of interest.

References

1. Aik J, Robin M Turner, Martyn D Kirk, et al. Evaluating food safety management systems in Singapore: A controlled interrupted time-series analysis of foodborne disease outbreak reports. *Food Control*. 2020;117:107324.
2. Yu Z, Dongyun J, Soyoun P, et al. Smart traceability for food safety. *Crit Rev Food Sci Nutr*. 2022;62(4):905–916.
3. Carrascosa C, Raheem D, Ramos F, et al. Microbial biofilms in the food industry—A comprehensive review. *Int J Environ Res Public Health*. 2021;18(4):2014.
4. Mota JDO, Géraldine B, Hervé P, et al. Environmental monitoring program to support food microbiological safety and quality in food industries: A scoping review of the research and guidelines. *Food Control*. 2021;130:108283.
5. Alonso VPP, Gonçalves MPMBB, Francisca AE de Brito, et al. Dry surface biofilms in the food processing industry: An overview on surface characteristics, adhesion and biofilm formation, detection of biofilms, and dry sanitization methods. *Compr Rev Food Sci Food Saf*. 2023;22(1):688–713.
6. Wiktorczyk-Kapischke N, Krzysztof Skowron, Katarzyna Grudlewska-Buda, et al. Adaptive response of *Listeria monocytogenes* to the stress factors in the food processing environment. *Front Microbiol*. 2021;12:710085.
7. Jovanovic J, Ornelis VFM, Madder A, et al. *Bacillus cereus* food intoxication and toxicoinfection. *Compr Rev Food Sci Food Saf*. 2021;20(4):3719–3761.
8. Pang X, Xin H, Xueying D, et al. Biofilm formation in food processing plants and novel control strategies to combat resistant biofilms: The case of *Salmonella* spp. *Food Sci Biotechnol*. 2023;32(12):1703–1718.

9. Sarion C, Codină GG, Dabija A. Acrylamide in bakery products: A review on health risks, legal regulations and strategies to reduce its formation. *Int J Environ Res Public Health*. 2021;18(8):4332.
10. Tsochatzis ED. Food contact materials: Migration and analysis—Challenges and limitations on identification and quantification. *Molecules*. 2021;26(11):3232.
11. Udovicki B, Mirjana A, Tanja CV, et al. Microplastics in food: Scoping review on health effects, occurrence, and human exposure. *Food Saf Risk*. 2022;9:7.
12. Peivasteh-Roudsari L, Raziye Barzegar-bafrouei, Kurush AS, et al. Origin, dietary exposure, and toxicity of endocrine-disrupting food chemical contaminants: A comprehensive review. *Heliyon*. 2023;9(7):e18140.
13. Feng H, Wang X, Duan Y, et al. Applying blockchain technology to improve agri-food traceability: A review of development methods, benefits and challenges. *J Clean Prod*. 2020;260:121031.
14. Patelli N, Mandrioli M. Blockchain technology and traceability in the agrifood industry. *J Food Sci*. 2020;85(11):3670–3678.
15. Xu Y, Li X, Zeng X, et al. Application of blockchain technology in food safety control: Current trends and future prospects. *Crit Rev Food Sci Nutr*. 2022;62(10):2800–2819.
16. Zanin LM, Pieterne AL, Diogo Thimoteo da C, et al. Influence of educational actions on transitioning of food safety culture in a food service context: Part 1—Triangulation and data interpretation of food safety culture elements. *Food Control*. 2021;119:107447.
17. Liggins G, Kim SR. How food safety culture is operationalized for retail food settings. *SAGE Open*. 2024.
18. Pai AS, Jaiswal S, Jaiswal AK. A comprehensive review of food safety culture in the food industry: Leadership, organizational commitment, and multicultural dynamics. *Foods*. 2024;13(24):4078.
19. Codex Alimentarius Commission. General Principles of Food Hygiene (CXC 1-1969). Rome: Food and Agriculture Organization of the United Nations and World Health Organization; 2023.
20. International Organization for Standardization (ISO). ISO 22000:2018 Food safety management systems - Requirements for any organization in the food chain. Geneva: ISO; 2018. Current edition confirmed in 2023; Amendment 1:2024.
21. Awuchi CG. Advancing HACCP, QMS, and FSMS in modern food and agricultural production; a systematic review on improving safe and sustainable production. *Cogent Food Agric*. 2026;12(1):2703900.
22. Onyeaka H, Ghosh S, Obileke K, et al. Preventing chemical contaminants in food: Challenges and prospects for safe and sustainable food production. *Food Control*. 2024;155:110040.
23. Soro AB, Whyte P, Bolton DJ, et al. Strategies and novel technologies to control *Campylobacter* in the poultry chain: A review. *Compr Rev Food Sci Food Saf*. 2020;19(4):1353–1377.
24. Liu X, Xuejuan X, Yangtai L, et al. Recent advances on the formation, detection, resistance mechanism, and control technology of *Listeria monocytogenes* biofilm in food industry. *Food Res Int*. 2024;180:114067.
25. Hua Z, Zhu MJ. Comprehensive strategies for controlling *Listeria monocytogenes* biofilms on food-contact surfaces. *Compr Rev Food Sci Food Saf*. 2024;23(3):e13348.
26. Choi W, Kim SS. Outbreaks, germination, and inactivation of *Bacillus cereus* in food products: A review. *J Food Prot*. 2020;83(9):1480–1487.
27. Dietrich R, Jessberger N, Ehling-Schulz M, et al. The food poisoning toxins of *Bacillus cereus*. *Toxins*. 2021;13(2):98.
28. Tuipulotu DE, Mathur A, Ngo C, et al. *Bacillus cereus*: Epidemiology, virulence factors, and host–pathogen interactions. *Trends Microbiol*. 2021;29(5):458–471.
29. Cayemite PE, Raymond P, Aider M. *Bacillus cereus* as an underestimated foodborne pathogen and new perspectives on its prevalence and methods of control: Critical and practical review. *ACS Food Sci Technol*. 2022;2(8):1196–1212.
30. Samtiya M, Matthews KR, Dhewa T, et al. Antimicrobial resistance in the food chain: Trends, mechanisms, pathways, and possible regulation strategies. *Foods*. 2022;11(19):2966.
31. Elafify M, Liao X, Feng J, et al. Biofilm formation in food industries: Challenges and control strategies for food safety. *Food Res Int*. 2024;190:114650.
32. Kumar A, Hardik P, Seema B, et al. Aflatoxin contamination in food crops: Causes, detection, and management: A review. *Food Prod Process Nutr*. 2021;3:17.
33. Jallow A, Huali Xie, Xiaoqian Tang, et al. Worldwide aflatoxin contamination of agricultural products and foods: From occurrence to control. *Compr Rev Food Sci Food Saf*. 2021.
34. Leite M, Freitas A, Silva AS, et al. Maize food chain and mycotoxins: A review on occurrence studies. *Trends Food Sci Technol*. 2021;115:307–331.
35. Iqbal SZ. Mycotoxins in food, recent development in food analysis and future challenges: A review. *Curr Opin Food Sci*. 2021;42:237–247.
36. Urbelis JH, Cooper JR. Migration of food contact substances into dry foods: A review. *Food Addit Contam Part A*. 2021;38(6):1044–1073.
37. Sarker A, Jang-EK, Abu Reza Md Towfiqul Islam, et al. Heavy metals contamination and associated health risks in food webs. *Environ Sci Pollut Res*. 2021;29(3):3230–3245.
38. Torres FG, De-la-Torre GE. Per- and polyfluoroalkyl substances (PFASs) in consumable species and food products. *J Food Sci Technol*. 2023;60(9):2319–2336.
39. Ogunbiyi OD, Timothy OA, Elizabeth OO, et al. Analytical approaches for screening of per- and polyfluoroalkyl substances in food items: A review of recent advances and improvements. *Environ Pollut*. 2023;329:121705.
40. Yoganandham ST, Hamid N, Junaid M, et al. Micro(nano)plastics in commercial foods: A review of their characterization and potential hazards to human health. *Environ Res*. 2023;236:116858.
41. Sohail M, Zunaira U, Sobia N, et al. Micro- and nanoplastics: Contamination routes of food products and critical interpretation of detection strategies. *Sci Total Environ*. 2023;891:164596.
42. Dzwolak W. Allergen cross-contact control plan supporting the implementation of food allergen management in small food businesses. *Food Control*. 2022;135:108777.
43. Jia L, Evans S. Improving food allergen management in food manufacturing: An incentive-based approach. *Food Control*. 2021;129:108246.
44. Remington BC, Joseph B, W. Marty Blom, et al. Allergen quantitative risk assessment within food operations: Concepts towards development of practical guidance based on an ILSI Europe workshop. *Food Control*. 2022;138:108917.
45. Linders YFM, Leo RL, W. Marty Blom, et al. Precautionary allergen labeling: Current communication problems and potential for future improvements. *Food Control*. 2023;147:109561.
46. Yuan L, Sadiq FA, Wang N, et al. Recent advances in understanding the control of disinfectant-resistant biofilms by hurdle technology in the food industry. *Crit Rev Food Sci Nutr*. 2021;61(22):3876–3891.
47. Insfran-Rivarola A, Diego Tlapa, Jorge Limon R, et al. A systematic review and meta-analysis of the effects of food safety and hygiene training on food handlers. *Foods*. 2020;9(9):1169.

48. Lonchamps PL, He Y, Wang K, et al. Detection of pathogens in foods using microfluidic “lab-on-chip”: A mini review. *J Agric Food Res.* 2022;10:100430.
49. Quintela IA, Vasse T, Lin CS, et al. Advances, applications, and limitations of portable and rapid detection technologies for routinely encountered foodborne pathogens. *Front Microbiol.* 2022;13:1054782.
50. Liu S, Kaixuan Z, Meiyuan H, et al. Research progress on detection techniques for point-of-care testing of foodborne pathogens. *Front Bioeng Biotechnol.* 2022;10:958134.
51. Panwar S, K Siddaardha Duggirala, Pooja Yadav, et al. Advanced diagnostic methods for identification of bacterial foodborne pathogens: Contemporary and upcoming challenges. *Crit Rev Biotechnol.* 2023;43(7):982–1000.
52. Cossettini A, Jasmina V, Michela M, et al. Rapid detection of *Listeria monocytogenes*, *Salmonella*, *Campylobacter* spp., and *Escherichia coli* in food using biosensors. *Food Control.* 2022;137:108962.
53. Zhao Y, Yuting S, Zhengzheng W, et al. The recent advances of high-throughput biosensors for rapid detection of foodborne pathogens. *TrAC Trends Anal Chem.* 2024;176:117736.
54. Gao R, Liu X, Xiong Z, et al. Research progress on detection of foodborne pathogens: The more rapid and accurate answer to food safety. *Food Res Int.* 2024;193:114767.
55. Zhang R, Yang T, Zhang Q, et al. Whole-genome sequencing: A perspective on sensing bacterial risk for food safety. *Curr Opin Food Sci.* 2022;47:100888.
56. Verheyen D, Van Impe JFM. The inclusion of the food microstructural influence in predictive microbiology: State-of-the-art. *Foods.* 2021;10(9):2119.
57. Varilla C, Marcone M, Annor GA. Potential of cold plasma technology in ensuring the safety of foods and agricultural produce: A review. *Foods.* 2020;9(10):1435.
58. Aganovic K, Christian Herte, Rudi F Vogel, et al. Aspects of high hydrostatic pressure food processing: Perspectives on technology and food safety. *Compr Rev Food Sci Food Saf.* 2021;20(4):3225–3266.
59. Laroque DA, Seó ST, Valencia GA, et al. Cold plasma in food processing: Design, mechanisms, and application. *J Food Eng.* 2022;312:110748.
60. Dhar P, Bhowmik A, Nath PC, et al. Smart technologies in food safety and quality monitoring: A review of technological advances, opportunities and challenges. *J Food Process Eng.* 2026;49(2):e70367.
61. Yu M, Jiang W, Cui X, et al. Biosensors with enzymatic amplification strategies for the detection of foodborne pathogenic microorganisms. *Biotechnol Adv.* 2026;92:108974.
62. Gbashi S, Njobeh PB. Enhancing food integrity through artificial intelligence and machine learning: A comprehensive review. *Appl Sci.* 2024;14(8):3421.