

Total laparoscopic hysterectomy versus total abdominal hysterectomy in the treatment of benign gynaecological disease: a retrospective review over 5 years

Abstract

Hysterectomy remains one of the most common gynaecological procedures performed in the UK. However unlike other parts of Europe and America, where laparoscopic hysterectomy (LH) rates have significantly increased, in the UK abdominal hysterectomy (AH) rates remain high and often the first choice for many surgeons. The minimal access route offers significant patient benefits over open surgery and the purpose of this study was to evaluate the role of total laparoscopic hysterectomy (TLH) versus total abdominal hysterectomy (TAH) in the management of benign gynaecological conditions. This retrospective study was carried out over a 5-year period and 296 procedures were included. Outcome measures included operating time, estimated blood loss (EBL), intraoperative and postoperative complications, postoperative analgesia requirements and length of hospital stay. TLH was associated with a significantly lower mean operating time (63.4 versus 75.3min $P<0.001$) and reduced EBL (145.1 versus 277.0ml $P<0.001$). Intraoperative complications were significantly less in the TLH group (1.9 versus 7.0% $P=0.029$) with no ureteric injuries noted. Analgesia requirements were also significantly less with fewer requiring breakthrough analgesia (6.2 versus 26.6% $P<0.001$). TLH was also associated with a significantly shorter inpatient hospital stay (1.7 versus 3.0 days $P<0.001$) and lower postoperative complication rates (6.8 versus 15.6% $P=0.016$). The results from our study highlight that TLH is superior to TAH in all operative outcome measures. With adequate training and experience TLH is a safe, reproducible technique that should be offered to all women requiring a hysterectomy for a normal sized uterus in the absence of significant adhesions.

Keywords: Hysterectomy, Laparoscopy

Volume 5 Issue 7 - 2016

Rebecca Mallick,¹ James English,¹ Natasha Waters¹

¹Brighton and Sussex University Hospitals (BSUH), UK

²Western Sussex Hospitals NHS Trust, UK

Correspondence: Rebecca Mallick, Department of Obstetrics and Gynaecology, Brighton and Sussex University Hospitals, Royal Sussex County Hospital, Eastern Road, Brighton, East Sussex, BN2 5BE, Tel 7411617143
Email rmallick@doctors.org.uk

Received: September 10, 2016 | **Published:** December 28, 2016

Background

Hysterectomy remains the most common gynaecological procedure performed in the UK, with on average 55,000 hysterectomies undertaken each year. Since the first laparoscopic hysterectomy (LH) was described by Reich et al.¹ in 1989, LH rates have increased significantly across parts of Europe and America. In Germany, for example, between 2007-2012, the rate of total laparoscopic hysterectomy (TLH) increased to approximately 30% while total abdominal hysterectomy (TAH) rates fell significantly to 7%. By comparison however, abdominal hysterectomy (AH) rates in the UK remain high and were 62% for the years 2011/2012.² However trends in the UK are changing and there is a growing acceptance of the role of LH by many surgeons, despite relatively recent recommendations from bodies such as NICE in 2007,³ ACOG in 2009⁴ and Cochrane in 2015⁵ advocating the vaginal route as the mode of first choice for hysterectomy.

Many safety concerns regarding LH stem from the evaluate study,⁶ which failed to show any real advantage for a laparoscopic approach and reported high major complication rates of 11.1%. However increasingly more studies have shown LH to be a safe, reproducible technique associated with low complication rates^{7,8} and significant patient benefits including reduced blood loss⁹ and hospital stay.¹⁰

We report on a series of 296 consecutive hysterectomies, performed

for the treatment of benign gynaecological disease in our department between 2009-2014. The aim of this study was to evaluate the role of TLH versus TAH in the management of benign gynaecological conditions specifically comparing operative outcomes, such as operating time and estimated blood loss (EBL), and complication rates.

Methods

This study was carried out over a 5-year period in a teaching hospital and tertiary referral centre for endometriosis. Total abdominal and total laparoscopic hysterectomies performed for the treatment of benign gynaecological disease during that period were included. Exclusion criteria included; malignancy, uterine size greater than 12 weeks, hysterectomy performed primarily for prolapse, hysterectomy performed in conjunction with the resection of deep infiltrating endometriosis including rectal resections and all subtotal hysterectomies including conversions from TAH.

Outcome measures included; operating time, EBL, intraoperative and postoperative complications, postoperative analgesia requirements and length of hospital stay. Intraoperative complications included bladder, bowel and ureteric injury and blood loss greater than 500ml. Postoperative complications were subdivided into minor and major complications. Minor complications included urinary tract infections (UTI), postoperative ileus, wound infection,

postoperative pyrexia $>38^{\circ}\text{C}$ and vault haematomas conservatively managed. Major postoperative complications included significant bleeding requiring return to theatre, vault/wound dehiscence and vault haematomas requiring surgical intervention. Data was analysed using SPSS (version 22). Descriptive statistical testing was utilised and a comparison of data made using the Mann-Whitney U test for continuous data and χ^2 analysis for nominal data. P values of <0.05 were considered statistically significant.

Findings

From 2009 to 2014, 296 hysterectomies were performed. Of these 161(54.4%) were performed laparoscopically, 128 (43.2%) were performed abdominally and 7(2.4%) vaginally. During this period the rate of TLH increased approximately sevenfold from 10% to 75%, while the rate of TAH fell from 87% to 25%. The rate of vaginal hysterectomy (VH) remained low at 0 to 5.7% (Figure 1). Due to the low VH rates, the data was excluded from comparison and further statistical analysis. Baseline characteristics between the 2 groups were comparable (Table 1). Pain was the most common indication for surgery in the laparoscopic group (42.2%) while heavy menstrual bleeding (HMB)/irregular bleeding predominated in the abdominal group (68.8%) (Table 2).

Table 1 Baseline patient characteristics

	Laparoscopy	Abdominal
Age (years)	43.7 (25-77)	45.3 (30-75)
BMI	30.0 (18-51)	28.1 (17-47)
Current smoker	30 (18.6%)	20 (15.6%)
Nulliparous	36(24.3%)	21 (17.2%)

Data presented as mean (range) or absolute numbers (%)

Table 2 Indications for surgery

	Laparoscopy	Abdominal
Pain	68 (42.2%)	20 (15.6%)
Hmb/irregular bleeding	59 (36.6%)	88 (68.8%)
Pain & bleeding	14 (8.7%)	8 (6.3%)
Atypical hyperplasia/ postmenopausal bleeding	11 (6.8%)	6 (4.7%)
Premenstrual tension	3 (1.9%)	3 (2.3%)
Severe dyskaryosis	4 (2.5%)	1 (0.8%)
Family history of ovarian cancer	1 (0.6%)	0 (0.0%)
Fibroids	1 (0.6%)	0 (0.0%)
Ovarian cysts	0 (0.0%)	2 (1.6%)

Data presented as absolute numbers (%)




Figure 1 Hysterectomy rates (%) over time.

TLH was associated with a significantly lower mean operating time (63.4 versus 75.3min $P<0.001$) and reduced EBL (145.1 versus

277.0ml $P<0.001$). Intraoperative complications were significantly less in the TLH group (1.9 versus 7.0% $P=0.029$). Two bladder injuries were noted in the TLH group compared to one bladder injury and one ureteric injury in the TAH group. No bowel injuries were noted in either group. EBL greater than 500mls was significantly greater in the TAH group (0.6 versus 5.5% $P=0.024$) (Table 3). The conversion to laparotomy rate was 1.2% ($n=2$): one was converted due to excessive bleeding and the other due to extensive adhesions. Return to theatre rates were lower in the TLH group (1.9 versus 3.1% $P=0.704$), with the most common reason in both groups being intra-abdominal bleeding. Analgesia requirements were significantly less in the TLH group. Oral analgesia was sufficient for 74.5% in the TLH group compared to 10.9% in the TAH group in whom PCA and/or epidural analgesia was utilised. Significantly fewer women in the TLH group required breakthrough analgesia (6.2 versus 26.6% $P<0.001$). TLH was associated with a significantly shorter inpatient hospital stay (1.7 versus 3.0 days $P<0.001$). Postoperative readmission rates were also lower in the TLH group (3.1 versus 4.7% $P=0.54$). The postoperative parameters are summarised in Table 4.

Table 3 Intraoperative parameters and complications

	Laparoscopy	Abdominal	P value
Operating times (min)	63.4 (20-147)	75.3 (34-155)	<0.001
Estimated blood loss (ml)	145.1 (0-800)	277.0 (50-1300)	<0.001
Intraoperative complications	3 (1.9%)	9 (7.0%)	0.029

Data presented as mean (range) or absolute numbers (%)

Table 4 Postoperative parameters

	Laparoscopy	Abdominal	P value
Return to theatre	3 (1.9%)	4 (3.1%)	0.704
Breakthrough analgesia needs	10 (6.2%)	34 (26.6%)	<0.001
Inpatient duration (days)	1.7 (1-6)	3.0 (1-11)	<0.001
Postoperative readmission	5 (3.1%)	6 (4.7%)	0.546

Data presented as mean (range) or absolute numbers (%)

TLH was associated with significantly lower total postoperative complication rates (6.8 versus 15.6% $P=0.016$) for both minor (5.0 versus 13.3% $P=0.013$) and major complications (1.9 versus 2.3% $P=1.00$), however for major complications the difference was not statistically significant. The postoperative complications are summarised in Table 5.

Table 5 Postoperative Complications

	Laparoscopy	Abdominal	P value
Total postoperative Complications	11 (6.8%)	20 (15.6%)	0.016
Major Complications	3 (1.9%)	3 (2.3%)	1.00
Intra-abdominal Bleed	2 (1.2%)	2 (1.6%)	1.00
Vault Haematoma requiring RTT	1 (0.6%)	0 (0.0%)	1.00
Abdominal wound breakdown	0 (0.0%)	1 (0.8%)	0.443
Minor Complications	8 (5.0%)	17 (13.3%)	0.013
UTI	1 (0.6%)	3 (2.3%)	0.325
Wound Infection	1 (0.6%)	2 (1.6%)	0.586
Ileus	2 (1.2%)	3 (2.3%)	0.658
Pyrexia >38	1 (0.6%)	1 (0.8%)	1.00
Urinary Retention	1 (0.6%)	1 (0.8%)	1.00
LRTI	0 (0.0%)	3 (2.3%)	0.086
Vault Haematoma	2 (1.2%)	3 (2.3%)	0.658

Data presented as absolute numbers (%)

Discussion

There are numerous clear benefits of LH yet in the UK, for benign gynaecological disease, AH rates remain significantly higher than both LH and VH. The most recent Cochrane review⁵ continues to advocate the use of VH over both AH and LH, however it does highlight the benefits of laparoscopy, which include a more rapid return to normal activity and less febrile episodes postoperatively. Nonetheless it reports a longer operating time and an increased risk of urinary tract injuries with LH.

When discussing LH complication rates, the most concerning research, particularly with regards to urinary tract injury, stems from the evaluate study by Garry et al.⁶ It quoted significantly higher risks of urinary tract injuries with LH (OR 2.61, 95% CI 1.22–5.60) and a high major complication rate of 11.1%. However since its publication there has been significant criticisms of this study^{7,11} namely the varied experience of the 43 surgeons performing the procedures and the unvalidated assumption that the learning curve for LH is approximately 20 cases. With this varied surgical experience, it can be hypothesised that the increased complication rates may have been a consequence of the relative inexperience of the surgeons rather than the technique of LH. Recent evidence suggests that the learning curve may require substantially more cases than 20 per year and the number of hysterectomies performed is likely to significantly impact complication rates.^{12–16} Evidence regarding the learning curve published by Twijnstra et al.¹⁵ suggests that there is a significant improvement in surgical outcome for up to 125 procedures, considerably higher than assumed in the evaluate study. When looking at the effect of surgical volume on outcome, Wallenstein et al.¹⁴ reported that the overall complication rates decreased from 6.2% for low volume surgeons to 4.2% for high volume surgeons.

In our study the intraoperative complication rates were significantly less in the TLH group (1.9 versus 7.0% $P=0.029$) with no ureteric injuries noted. This low complication rate is replicated in the wider literature.^{17–19} Doganay et al.¹⁸ reported no significant differences in the rates of bladder or ureteric injury associated with LH and VH; however there were significant differences in the urinary tract injury rates when compared to AH. Thus, current evidence tends to suggest that AH is associated with the highest rate of urinary tract injury with the rates of injury for LH and VH being equivalent. We also found a significant reduction in EBL in the TLH group (145.1 versus 277.0ml $P<0.001$) in keeping with other studies.^{19,20} Specifically for TLH, blood loss has been found to increase with uterine size⁸ and increasing BMI.²¹

Historically it was suggested that the operating time associated with performing LH was likely to be increased when compared with open hysterectomy and this is the conclusion of the most recent Cochrane review.⁵ However in our study, TLH was associated with a significantly lower mean operating time (63.4 versus 75.3min $P<0.001$). One key feature to take into account is surgeon experience and as described by Pather et al.²² there does not appear to be any difference in operating times once the initial learning curve has been passed.

The anticipated decrease in postoperative pain associated with minimally invasive surgery is supported by the current literature^{23,24} and our data is in keeping with this. We found overall analgesia requirements to be significantly less in the TLH group with oral analgesia sufficient for 74.5%, compared to 10.9% in the TAH group. Significantly fewer women in the LH group required breakthrough analgesia (6.2 versus 26.6% $P<0.001$). Interestingly Ghezzi et al.²⁴

also described a significant advantage with regards to postoperative pain when comparing LH to VH.

One would also expect that hospital stay would be reduced when surgery is performed by the minimally invasive route and this is borne out by the literature.^{9,22,25} Our data is again in keeping with this with the mean inpatient stay being significantly shorter in the TLH group (1.7 versus 3.0 days $P<0.001$). When compared to VH, Ghezzi et al.²⁴ reported that TLH was also associated with a shorter hospital stay.

With regards to overall postoperative complications we found the rates significantly lower in the TLH group (6.8 versus 15.6% $P=0.016$). Major complications were less (1.9 versus 2.3% $P=1.00$), although this was not statistically significant, however minor complication rates were significantly less (5.0 versus 13.3% $P=0.013$). These complication rates are comparable to the wider literature. Karaman et al.²⁶ described a major complication rate of 1% observed in a series of 1120 laparoscopic hysterectomies, while in another prospective series of 3190 laparoscopic hysterectomies, Donnez et al.⁷ described a similarly low major complication rate of 0.37–0.51%. Furthermore Wright et al.^{27–29} and Kondo et al.²⁵ reported similarly significant benefits of laparoscopic over open hysterectomy for both benign and malignant disease. Leiserkowitz et al.³⁰ reported that vascular and bowel injuries, pulmonary embolism and wound problems were all more common with TAH.

For many years isolated case reports suggested that TLH carried an increased risk of vault dehiscence with commentators blaming suturing techniques and the use of energy to transect the vaginal vault thus delaying healing. This was further highlighted by Hur et al.³¹ who reported that LAVH was associated with a four fold increase in the risk of vault dehiscence when compared with VH. However a more recent large study of 9,973 hysterectomies by Koo et al.³² demonstrated the highest vault dehiscence risk to be associated with TAH (0.6%) and the lowest after TLH (0.2%) with VH at 0.4% ($p=0.016$). In our data no cases of vault dehiscence was noted in either group.

Conclusion

There is now substantial evidence that routine AH is inferior to both LH and VH in all outcome measures. Since this is so, it is no longer feasible, or one can argue ethically correct, to support the routine use of AH to remove a normal size uterus. Our study highlights the significant benefits of LH including a shorter operating time, reduced EBL, less intraoperative and postoperative complications, less analgesia requirements postoperatively and a shorter inpatient stay. The real choice is between LH and VH, which seem equal in many outcome measures however there is increasing evidence that pain may be less and hospital stay shorter with the laparoscopic approach. Also a narrow vagina, lack of prolapse or the presence of abdominal pathology including adhesions, endometriosis and adnexal disease, largely preclude a vaginal approach. Also with the increasing practice of prophylactic salpingectomy at the time of hysterectomy, a laparoscopic approach makes this technically more feasible. In effect, it should be as unacceptable to perform a routine TAH as it is currently to perform a routine open cholecystectomy, and women, dependent on the nature of the pathology, should be offered a minimally invasive procedure. In experienced hands, TLH is a safe, reproducible technique with low complication rates; however significant training is required to attain this level of expertise. In order to bring this reality into effect in the UK, a major change in terms of surgical training and mentorship will be required.

Ethical approval

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Acknowledgments

None.

Conflicts of interest

None.

References

- Reich H, Decaprio J, McGlynn F. Laparoscopic Hysterectomy. *J Gynaecol Surg.* 1989;5(2):213–216.
- NHS. UK hysterectomy rates. 2013
- NICE. Laparoscopic techniques for hysterectomy. NICE interventional procedure guidance [IPG239]. 2007
- ACOG. Choosing the route of hysterectomy for benign disease. ACOG Committee Opinion No. 444. *Obstet Gynecol.* 2009;114(5):1156–1158.
- Aarts JW, Nieboer TE, Johnson N, et al. Surgical approach to hysterectomy for benign gynaecological disease. *Cochrane Database Syst Rev.* 2015;Cd003677.
- Garry R, Fountain J, Mason S, et al. The eVALuate study: two parallel randomised trials, one comparing laparoscopic with abdominal hysterectomy, the other comparing laparoscopic with vaginal hysterectomy. *BMJ.* 2004;328:1229–1236.
- Donnez O, Jadoul P, Squifflet J, et al. A series of 3190 laparoscopic hysterectomies for benign disease from 1990 to 2006: evaluation of complications compared with vaginal and abdominal procedures. *BJOG.* 2009;116(4):492–500.
- Uccella S, Cromi A, Bogani G, et al. Systematic implementation of laparoscopic hysterectomy independent of uterus size: clinical effect. *J Minim Invasive Gynecol.* 2013;20(4):505–516.
- Yi YX, Zhang W, Zhou Q, et al. Laparoscopic-assisted vaginal hysterectomy vs abdominal hysterectomy for benign disease: a meta-analysis of randomized controlled trials. *Eur J Obstet Gynecol Reprod Biol.* 2011;159(1):1–18.
- Warren L, Ladapo JA, Borah BJ, et al. Open abdominal versus laparoscopic and vaginal hysterectomy: analysis of a large United States payer measuring quality and cost of care. *J Minim Invasive Gynecol.* 2009;16(5):581–588.
- Donnez J, Squifflet J, Jadoul P, et al. Results of eVALuate study of hysterectomy techniques: High rate of complications needs explanation. *BMJ.* 2004;328(7440):643.
- Wattiez A, Soriano D, Cohen SB, et al. The learning curve of total laparoscopic hysterectomy: comparative analysis of 1647 cases. *J Am Assoc Gynecol Laparosc.* 2002;9(3):339–345.
- Twijnstra AR, Blikkendaal MD, Kolkman W, et al. Implementation of laparoscopic hysterectomy: maintenance of skills after a mentorship program. *Gynecol Obstet Invest.* 2010;70(3):173–178.
- Wallenstein MR, Ananth CV, Kim JH, et al. Effect of surgical volume on outcomes for laparoscopic hysterectomy for benign indications. *Obstet Gynecol.* 2012;119(4):709–716.
- Twijnstra AR, Blikkendaal MD, van Zwet EW, et al. Predictors of successful surgical outcome in laparoscopic hysterectomy. *Obstet Gynecol.* 2012;119(4):700–708.
- Boyd LR, Novetsky AP, Curtin JP. Effect of surgical volume on route of hysterectomy and short-term morbidity. *Obstet Gynecol.* 2010;116(4):909–915.
- Gendy R, Walsh CA, Walsh SR, et al. Vaginal hysterectomy versus total laparoscopic hysterectomy for benign disease: a metaanalysis of randomized controlled trials. *Am J Obstet Gynecol.* 2011;204(5):388.e1–388.e8.
- Doganay M, Yildiz Y, Tonguc E, et al. Abdominal, vaginal and total laparoscopic hysterectomy: perioperative morbidity. *Arch Gynecol Obstet.* 2011;284(2):385–389.
- Candiani M, Izzo S, Bulfoni A, et al. Laparoscopic vs vaginal hysterectomy for benign pathology. *Am J Obstet Gynecol.* 2009;200(4):368.e1–368.e7.
- Jugnet N, Cosson M, Wattiez A, et al. Comparing vaginal and coelioscopic total or subtotal hysterectomies: prospective multicentre study including 82 patients. *Gynaecol Endoscop.* 2001;10(5–6):315–321.
- Siedhoff MT, Carey ET, Findley AD, et al. Effect of extreme obesity on outcomes in laparoscopic hysterectomy. *J Minim Invasive Gynecol.* 2012;19(6):701–717.
- Pather S, Loadsman JA, Mansfield C, et al. Perioperative outcomes after total laparoscopic hysterectomy compared with fast-track open hysterectomy – a retrospective case-control study. *Aust N Z J Obstet Gynaecol.* 2011;51(5):393–396.
- Naik R, Jackson KS, Lopes A, et al. Laparoscopic assisted radical vaginal hysterectomy versus radical abdominal hysterectomy—a randomised phase II trial: perioperative outcomes and surgicopathological measurements. *BJOG.* 2010;117(6):746–751.
- Ghezzi F, Uccella S, Cromi A, et al. Postoperative pain after laparoscopic and vaginal hysterectomy for benign gynecologic disease: a randomized trial. *Am J Obstet Gynecol.* 2010;203(2):118.e1–118.e8.
- Kondo W, Bourdel N, Marengo F, et al. Is laparoscopic hysterectomy feasible for uteri larger than 1000 g? *Eur J Obstet Gynecol Reprod Biol.* 2011;158(1):76–81.
- Karaman Y, Bingol B, Gunenc Z. Prevention of complications in laparoscopic hysterectomy: experience with 1120 cases performed by a single surgeon. *J Minim Invasive Gynecol.* 2007;14(1):78–84.
- Wright JD, Ananth CV, Lewin SN, et al. Robotically assisted vs laparoscopic hysterectomy among women with benign gynecologic disease. *JAMA.* 2013;309(7):689–698.
- Wright JD, Neugut AI, Wilde ET, et al. Use and benefits of laparoscopic hysterectomy for stage I endometrial cancer among medicare beneficiaries. *J Oncol Pract.* 2012;8(5):e89–e99.
- Wright JD, Herzog TJ, Neugut AI, et al. Comparative effectiveness of minimally invasive and abdominal radical hysterectomy for cervical cancer. *Gynecol Oncol.* 2012;127(1):11–17.
- Leiserowitz GS, Xing G, Parikh-Patel A, et al. Laparoscopic versus abdominal hysterectomy for endometrial cancer: comparison of patient outcomes. *Int J Gynecol Cancer.* 2009;19(8):1370–1376.
- Hur HC, Donnellan N, Mansuria S, et al. Vaginal cuff dehiscence after different modes of hysterectomy. *Obstet Gynecol.* 2011;118(4):794–801.
- Koo YJ, Kim DY, Kim JH, et al. Vaginal cuff dehiscence after hysterectomy. *Int J Gynaecol Obstet.* 2013;122(3):248–352.