

Incidental breast lesion on chest CT scan: a review

Abstract

Most breast cancers are diagnosed by conventional breast imaging which includes mammography, ultrasound and MRI. However, CT is an often overlooked modality in the detection and diagnosis of breast cancer. Incidental findings of breast cancer by CT are not so rare. This article aims to aware radiologists and gynaecologists to the potential of breast cancer detection with CT. In this brief review of the literature, we will discuss the morphological characteristics of breast tumours on CT and the imaging management of suspicious lesions discovered on CT.

Volume 13 Issue 1 - 2022

Quoc Duy Vo, Thierry Molteni

FMH specialist in radiology, EHC Hospital Morges, Switzerland

Correspondence: Dr Quoc Duy Vo, FMH specialist in radiology, EHC Hospital Morges, Chemin du Cret 2, Switzerland, Tel 0041797782164, Email quocduy.vo@ehc.vd.ch

Received: January 04, 2022 | **Published:** January 14, 2022

Introduction and background

Breast cancer is the most frequent cancer amongst women and it represents a global public health problem, with 1.7 million new cases diagnosed every year¹ and a worldwide annual prediction of 3,2 million new cases per year by 2050.² It is also the leading cause of cancer death in women aged between 20 and 59 years, with approximately 0.5 million deaths per year in 2013.¹ Today, breast cancer is a well-known disease with readily recognized risk factors, which include family history, genetic abnormalities, alcohol consumption, nutrition, female's hormonal context, obesity and thoracic radiotherapy.³

Breast cancer diagnosis relies on clinical examination and radiological investigations, which include a plethora of medical imaging techniques, most commonly mammography. Ultrasound and magnetic resonance imaging (MRI) have been frequently employed as complementary tools to mammography, and have repeatedly proved their efficiency.⁴ Currently, CT scan has become an exceedingly common imaging modality, used in nearly all areas of medicine. The steady increase in utilization of this technique in the past few decades has led to an increased detection rate of "incidental" breast lesions, which may be of major health concern.⁵⁻⁸ Therefore, radiologists must be familiar with breast findings on CT scans and must be able to propose an adequate workup.

Conventional breast imaging

Mammography is the most common imaging modality used to assess breast disease and it is capable of detecting subtle abnormalities such as microcalcifications, architectural distortion and opacities with spiculated margins with a specificity of 55% and sensibility of 70%.⁹ Recently, computer-aided detection (CAD) has been developed to help radiologists in identifying breast anomalies. It employs a specific computer algorithm capable of individualizing and marking suspicious mammogram findings, assisting the radiologist during the interpretation of the exam.^{10,11} However, there are some limitations to mammograms, especially breast density that can provoke superimpositions, which are able to hide pathological lesions and avoid detection, even with the use of CAD.^{4,10} For these reasons, digital breast tomosynthesis has been extensively used in conjunction with mammography. It acquires multiple low dose mammographic projections of the breasts which are organized in stacks of slices after tomographic reconstruction, similar to CT, reducing tissue superimposition and improving detection of masses or architectural distortion (Figure 1).^{4,12}




Figure 1 A. Mammography of the right breast revealed an irregular and spiculated opacity in the outer upper quadrant (white arrow). B. Tomosynthesis performed on the cranio-caudal view (left image) improves characterization of the lesion compared to standard mammography (right image), with better visualization of the spiculated aspect of the lesion.

Breast ultrasound is usually used conjointly with mammography, particularly in patients with dense breasts. It was first used to differentiate solid from cystic lesions during the 70's.^{4,10} Nowadays, with technological advancement, modern ultrasound devices are able to characterize breast lesions (Figure 2) and to improve diagnosis

of breast cancer with a negative predictive value of 99.5% and a sensitivity of 98.4%,¹³ thus avoiding unnecessary biopsies. In spite of its technical improvements, ultrasound still bears some limitations: first of all, it is an operator-dependent modality, therefore physicians's experience may play a crucial role in disease detection; secondly, it is unable to detect microcalcifications, one of the early hallmarks of ductal carcinoma in situ (DCIS), which is believed to herald infiltrating ductal carcinoma.^{4,10,13}




Figure 2 A. Hypoechoic lesion with spiculated borders associated with a hyperechoic halo and an acoustic shadow. B. Hypoechoic lesion with irregular and angulated borders. C. Well delimited hypoechoic lesion presenting a "taller than wide" shape. D. Simple cyst presenting as a well delimited anechoic lesion with posterior enhancement. E. Complex cyst with septations and mural nodule. F. Well delimited hypoechoic lesion containing small cysts with ovoid shape.


During the past two decades, breast MRI has emerged as one of the most valuable imaging techniques in the diagnosis of breast cancer. It is considered the most sensitive breast imaging modality, providing a sensitivity of 90% for breast cancer detection with variable specificity ranging from 37% to 100%.¹⁴⁻¹⁶ Its use allows detection of multifocal cancer, recognition of invasive components in DCIS, detection of cancer in excessively dense breasts, evaluation of tumor response in neoadjuvant chemotherapy¹⁷ and screening for high risk women including those with BRCA mutations.⁴ Its main disadvantages lie on its availability and on its lengthy scan times which can be difficult to manage, especially for claustrophobic patients.

Studies and frequency

As mentioned earlier, a dramatic rise in the use of CT scan, normally performed for indications other than breast disease, has led to an increased detection of incidental breast abnormalities. Consequently, a considerable number of studies have been conceived to evaluate the incidence and possible outcomes of incidental breast lesions detected on unenhanced or contrast-enhanced CT, with variable results depending on the use of contrast media. In a large study, Shojaku et al. performed unenhanced chest CT scans on 1'008 female patients to investigate the frequency of breast tumors. Breast anomalies were detected in 6 patients, in which 3 were diagnosed with breast cancer (0,3%).⁸ In another large-scale study, Swensen et al. assessed the performance of low-dose chest CT to evaluate the presence of incidental pulmonary nodules in 1'520 individuals aged 50 years or older, who had smoked 20 pack-years or more. In 3 female patients, a diagnosis of breast cancer was found (0,1%).¹⁸ Lin et al. reported 23 female patients with incidental breast findings in a series of 2'250 contrast-enhanced chest CT scan. Among these 23 patients, 16 had breast cancer which was confirmed by further investigations (0,7%),⁵ suggesting that the use of contrast media may improve breast cancer detection in CT Scan.¹⁹

CT-based protocols and characterization of malignant breast lesions

Some authors advocate that contrast-enhanced CT improves detection and characterization of breast lesions. Perrone et al. propose its use as an alternative to MRI in patients with usual contraindications such as incompatible pacemakers or surgical clips, severe dyspnea and claustrophobia. They recommend to acquire CT images at 1 minute after contrast administration and to apply a cutoff attenuation value of 90 Hounsfield units that would allow to differentiate between benign and malignant lesions (Figure 3).²⁰ Inoue et al. demonstrate that the use of dynamic contrast-enhanced CT (DCE-CT) like DCE-MRI, can be useful to identify and to characterize breast lesions. In their protocol, acquisitions were obtained at 1, 3, and 8 minutes after intravenous contrast injection. On time-density curve analysis, the washout pattern had a positive predictive value for breast malignancy of 93%, with a sensitivity of 91%.²¹ Morphological criteria can also be implemented in order to distinguish malignant from benign lesions. Spiculated or hazy margins, irregular shape and rim enhancement can all be looked upon as reliable features of malignancy.^{6,21} Moreover, axillary lymphadenopathies, pectoralis muscle infiltration and pleural effusion strongly suggest tumoral invasion.⁶ In summary, contrast-enhanced CT scan seems to be a reliable alternative to MRI, with the advantages of being faster, particularly for dynamic studies, as well as less susceptible to movement artifacts.²⁰ However, due to radiation exposure concerns, chest CT scan must be reserved for elderly patients.²²






Figure 3 A. Follow-up contrast-enhanced chest CT in a patient with lung carcinoma shows an incidental right breast lesion with contrast enhancement up to 102 HU. B. Same image with smaller FOV (field-of-view) evidencing the spiculated and irregular borders. C. Digital mammography of the same patient showing an irregular lesion with blurred borders. D. Ultrasound imaging reveals a hypoechoic lesion with angulated borders associated with a hyperechoic halo.

Conclusion

Nowadays, chest CT scan is widely performed for a variety of indications other than breast disease, though its relentless increasing use over the years has resulted in an ever-growing detection of incidental breast findings. Therefore, radiologists must be able to recognize and to assess these findings. Breast tumors morphological criterias on CT are the same as in MRI and can therefore be reliably used. However, any incidental breast lesion on CT requires conventional breast imaging assesment including mammography and echography, breast MRI can be performed for dense breasts or in case of multicentric lesions.

Acknowledgments

None.

Funding

None.

Conflicts of interest

The author declares that they have no competing interests.

References

1. Winters S, Martin C, Murphy D, et al. Breast cancer epidemiology, prevention, and screening. *Prog Mol Biol Transl Sci.* 2017;151:1–32.

2. Tao Z, Shi A, Lu C, et al. Breast cancer: epidemiology and etiology. *Cell Biochem Biophys.* 2015;72(2):333–338.
3. Angahar LT. An overview of breast cancer epidemiology, risk factors, pathophysiology, and cancer risks reduction. *MOJ Biol Med.* 2017;1(4):00019.
4. Joe BN, Sickles EA. The evolution of breast imaging: past to present. *Radiology.* 2014;273(2 Suppl):S23–S44.
5. Wen-Chiung Lin, Hsian-He Hsu, Chao-Shiang Li, Jyhet al. Incidentally detected enhancing breast lesions on chest computed tomography. *Korean J Radiol.* 2011;12(1):44–51.
6. Jung Hee Son, Hyun Kyung Jung, Jong Woon Song, et al. Incidentally detected breast lesions on chest CT with US correlation: a pictorial essay. *Diagn Interv Radiol.* 2016;22(6):514–518.
7. P Moyle, L Sonoda, P Britton, R Sinnatamby. Incidental breast lesions detected on CT: what is their significance? *Br J Radiol.* 2010; 83(987):233–240.
8. Hiroko Shojaku, Hikaru Seto, Hisakazu Iwai, et al. Detection of incidental breast tumors by noncontrast spiral computed tomography of the chest. *Radiation medicine.* 2008;26(6):362–367.
9. Pisano ED, Yaffe MJ. Digital mammography. *Radiology.* 2005;234(2):353–362.
10. Ioana Andreea Gheonea, Zoia Stoica, Simona Bondari. Differential diagnosis of breast lesions using ultrasound elastography. *Indian J Radiol Imaging.* 2011;21(4):301–305.
11. Freer TW, Ulissey MJ. Screening mammography with computer-aided detection: prospective study of 12,860 patients in a community breast center. *Radiology.* 2001;220(3):781–786.
12. Skaane P, Bandos AI, Gullien R, et al. Comparison of digital mammography alone and digital mammography plus tomosynthesis in a population-based screening program. *Radiology.* 2013;267(1):47–56.
13. Hooley RJ, Scoutt LM, Philpotts LE. Breast ultrasonography: state of the art. *Radiology.* 2013;268(3):642–659.
14. Gurpreet S. Dhillon, Nick Bell, et al. Breast MR imaging: what the radiologist needs to know. *J Clin Imaging Sci.* 2011;48.
15. Macura KJ, Ouwkerk R, Jacobs MA, et al. Patterns of enhancement on breast MR images: interpretation and imaging pitfalls. *Radiographics.* 2006;26(6):1719–1734.
16. Carol H. Lee. Problem solving MR imaging of the breast. *Radiologic Clinics of North America.* 2004;42(5):919–934.
17. Gisela LG Menezes, Floor M Knuttel, Bertine L Stehouwer, et al. Magnetic resonance imaging in breast cancer: A literature review and future perspectives. *World J Clin Oncol.* 2014;5(2):61–70.
18. Swensen SJ, Jett JR, Hartman TE, et al. Lung cancer screening with CT: mayo clinic experience. *Radiology.* 2003;226(3):756–761.
19. Nicolas D Prionas, Karen K Lindfors, Shonket Ray, et al. Contrast-enhanced dedicated breast CT: initial clinical experience. *Radiology.* 2010;256(3):714–723.
20. A Hussain, A Gordon-Dixon, H Almusawy, et al. The incidence and outcome of incidental breast lesions detected by computed tomography. *Ann R Coll Surg Engl.* 2010;92(2):124–126.
21. Inoue M, Sano T, Watai R, et al. Dynamic multidetector CT of breast tumors: diagnostic features and comparison with conventional techniques. *AJR Am J Roentgenol.* 2003;181(3):679–686.
22. Johannes Gossner. Intramammary findings on CT of the chest – a review of normal anatomy and possible findings. *Pol J Radiol.* 2016;81:415–421.