

Extratropical cyclone in the South Georgia and south sandwich islands and double spiral galaxies

Abstract

The work is focused on the comparative analysis of the shape of spiral galaxies and the subtropical cyclone that formed north of Georgia Island and passed north of the South Sandwich Islands, in the South Atlantic Ocean. Subtropical cyclones with double spirals appear to be common in these areas of the South Atlantic. A subtropical cyclone is a weather system that has some characteristics of a tropical cyclone and some characteristics of an extratropical cyclone. They can form between the equator and the 50th parallel. In mathematics, a spiral is a curve, which emanates from a point, moving farther away as it revolves around the point. The characteristic shape of hurricanes, cyclones, typhoons is a spiral. The characteristic equation of which spiral the Extratropical Cyclone (EC) Its double spiral shape, whose mathematical equation has already been defined as Cote's spiral, Gobato et al. (2022) and similarly Lindblad¹ show shape of double spiral galaxies, already studied among others is discussed here.

Keywords: Galaxy, South Georgia and the South Sandwich Islands, Subtropical cyclone, Double spiral galaxy.

Volume 7 Issue 3 - 2023

Ricardo Gobato,¹ Abhijit Mitra,² Poulomi Mullick³

¹Green Land Landscaping and Gardening, Seedling Growth Laboratory, Brazil

²Department of Marine Science, University of Calcutta, India

³Department of Management, Techno India University, India

Correspondence: Ricardo Gobato, Green Land Landscaping and Gardening, Seedling Growth Laboratory, 86130-000, Parana, Brazil, Email ricardogobato@hotmail.com, ricardogobato@gardener.com

Received: May 10, 2023 | **Published:** July 12, 2023

Introduction

The South Georgia Group lies about 1,390 km (860 mi; 750 mi) east-southeast of the Falkland Islands, at

54°-55°S, 36°-38°W. It comprises South Georgia Island¹ itself, by far the largest island in the territory, and the islands that immediately surround it and some remote and isolated islets to the west and east-southeast. It has a total land area of 3,756 square kilometers (1,450 sq. mi), including satellite islands, but excluding the South Sandwich Islands, which form a separate island group.² A cyclone is a large air mass that rotates around a strong center of low atmospheric pressure, counterclockwise in the Northern Hemisphere and clockwise in the Southern Hemisphere as viewed from above (opposite to an anticyclone).³⁻⁸ A subtropical cyclone is a weather system that has some characteristics of a tropical cyclone and some characteristics of an extratropical cyclone. They can form between the equator and the 50th parallel.^{3,7,9-14}

These storms usually have a radius of maximum winds that is larger than what is observed in purely tropical systems, and their maximum sustained winds have not been observed to exceed about 32 m/s (64 knots). Subtropical cyclones sometimes become true tropical cyclones, and likewise, tropical cyclones occasionally become subtropical storms. Subtropical cyclones in the Atlantic basin are classified by their maximum sustained surface winds: Subtropical depressions have surface winds less than 18 m/s (35 knots), while subtropical storms have surface winds greater than or equal to 18 m/s.^{7,8,13,15-26}

¹South Georgia and the South Sandwich Islands is a British Overseas Territory in the southern Atlantic Ocean. It is a remote and inhospitable collection of islands, consisting of South Georgia and a chain of smaller islands known as the South Sandwich Islands. South Georgia is 165 kilometers (103 mi) long and 35 km (22 mi) wide and is by far the largest island in the territory. The South Sandwich Islands lie about 700 km (430 mi) southeast of South Georgia. The territory's total land area is 3,903 km² (1,507 sq. mi). The Falkland Islands are about 1,300 km (810 mi) west from its nearest point.

Tropical cyclones are compact, circular storms, generally some 320 km (200 miles) in diameter, whose winds swirl around a central region of low atmospheric pressure. The winds are driven by this low-pressure core and by the rotation of the Earth, which deflects the path of the wind through a phenomenon known as the Coriolis force. As a result, tropical cyclones rotate in a counterclockwise (or cyclonic) direction in the Northern Hemisphere and in a clockwise (or anticyclonic) direction in the Southern Hemisphere.^{7,8,13,15-26} The occurrence of cyclones is relatively common for the region at this time of year, but the recent phenomenon has been exacerbated by other meteorological and atmospheric factors. This phenomenon, with this feature to lower the pressure inside quickly generates very strong winds and so that name of explosive cyclones.^{10-13,16} For large-scale occurs, the subtropical cyclones influence and are influenced by the weather and other atmospheric phenomena point of view, the call synoptic condition.²⁷

In mathematics, a spiral is a curve, which emanates from a point, moving farther away as it revolves around the point.²⁸⁻³⁰ The characteristic shape of hurricanes, cyclones, typhoons is a spiral,^{7,8,13,27,31-37} are several types of turns, and determining the characteristic equation of which spiral the cyclone bomb (CB)³⁸ fits into is the goal of the work. Spiral galaxies form a class of galaxy originally described by Edwin Hubble in his 1936 work *The Realm of the Nebulae* and, as such, form part of the Hubble sequence. Most spiral galaxies consist of a flat, rotating disk containing stars, gas and dust, and a central concentration of stars known as the bulge. These are often surrounded by a much fainter halo of stars, many of which reside in globular clusters.³⁹

Spiral galaxies are named by their spiral structures that extend from the center into the galactic disc. The spiral arms are sites of ongoing star formation and are brighter than the surrounding disc because of the young, hot OB stars that inhabit them.³⁹⁻⁴²

The core of cyclone presents the form of a double spiral, Figures 1-3, in the same way the study of the spiral of the galaxies of Lindblad.^{1,43} This spiral is denoted from Cotes Spiral Gobato et al.^{11-16,23-25,28-30}




Figure 1 Image of Georgia, scale 1:200, on April 11, 2003,PM, and nucleus at the coordinates given in the image.⁴⁵




Figure 5 Image of Georgia, scale 1:100, in surface wind model generated by the Zoom Earth system, on April 11, 2003, 12:00,with 5km/h WSW, and nucleus at the coordinates given in the image.⁴¹




Figure 2 Image of Georgia, at scale 1:20, on April 11, 2003,AM, whose nucleus was at the approximate coordinates of those in the image.⁴⁵

Spiral galaxies

The Figure 6 show spectacular spiral galaxies using the impressive power of the HAWK-I^{2,50,51}




Figure 3 Image of Georgia, scale 1:20, on April 11, 2003,PM and nucleus at the coordinates given in the image.⁴⁵




Figure 6 Image of the nearby galaxy Messier 83 was taken in the infrared part of the spectrum with the HAWK-I instrument on ESO's Very Large Telescope. The very fine image quality of this camera, coupled with the huge light-collecting power of the VLT, reveals vast numbers of stars within the galaxy. The images were taken in three different parts of the infrared spectrum and the total exposure time was eight and a half hours, split into more than five hundred exposures of one minute each. The field of view is about 13 arcminutes across.^{50,59}

Messier 83 (The Southern Pinwheel)

The Figure (7) show Hubble image captures hundreds of thousands of individual stars, thousands of star clusters and hundreds of supernova remnants in the spiral galaxy M83. Also known as the




Figure 4 Image of Georgia, scale 1:100, in surface wind model generated by the Zoom Earth system, on April 11, 2003, 12:00,for atmospheric pressure at sea level in the cyclone vortex^{1,36-39}, and nucleus at the coordinates given in the image.⁴⁵

²The HAWK-I instrument mounted on the telescope's Nasmyth (side) port. HAWK-I is attached on Yepun, Unit Telescope number 4 of ESO's Very Large Telescope and saw First Light on the night of 31 July 2007. HAWK-I covers about 1/10th the area of the Full Moon in a single exposure. It is uniquely suited to the discovery and study of faint objects, such as distant galaxies or small stars and planets. HAWK-I is one of the newest and most powerful cameras on ESO's Very Large Telescope (VLT). It is sensitive to infrared light, which means that much of the obscuring dust in the galaxies spiral arms becomes transparent to its detectors. Compared to the earlier, and still much-used, VLT infrared camera ISAAC, HAWK-I has sixteen times as many pixels to cover a much larger area of sky in one shot and, by using newer technology than ISAAC, it has a greater sensitivity to faint infrared radiation. Because HAWK-I can study galaxies stripped bare of the confusing effects of dust and glowing gas it is ideal for studying the vast numbers of stars that make up spiral arms

Southern Pinwheel, this galaxy is located 15 million light-years away from Earth in the constellation Hydra. It was discovered in 1752 by the French astronomer Nicolas Louis de Lacaille. With an apparent magnitude of 7.5, M83 is one of the brightest spiral galaxies in the night sky. It can be observed using a pair of binoculars most easily in May. [49], [50]

Hubble's image reveals interstellar bubbles produced by nearly 300 supernovas. By studying these supernova remnants, astronomers can better understand the nature of the stars that exploded and dispersed their nuclear processed chemical elements back into the galaxy, contributing to the next generation of new stars.



Figure 7 Hubble image captures hundreds of thousands of individual stars, thousands of star clusters and hundreds of supernova remnants in the spiral galaxy M83. Also known as the Southern Pinwheel, this galaxy is located 15 million light-years away from Earth in the constellation Hydra.⁵⁰

A. NGC 1566

NGC 1566³, sometimes known as the Spanish Dancer, is an intermediate spiral galaxy in the constellation Dorado, positioned about 3.5° to the south of the star Gamma Doradus. It was discovered on May 28, 1826 by Scottish astronomer James Dunlop. At 10th magnitude, it requires a telescope to view. The distance to this galaxy remains elusive, with measurements ranging from 6 Mpc up to 21 Mpc.^{53,54} The small but extremely bright nucleus of NGC 1566 is clearly visible in this image, a telltale sign of its membership of the Seyfert class of galaxies. The centers of such galaxies are very active and luminous emitting strong bursts of radiation and potentially harboring supermassive black holes that are many millions of times the mass of the sun.^{53,54}

NGC 1566 is not just any Seyfert galaxy; it is the second brightest Seyfert galaxy known. It is also the brightest and most dominant member of the Dorado Group, a loose concentration of galaxies that together comprise one of the richest galaxy groups of the southern hemisphere. This image highlights the beauty and awe-inspiring nature of this unique galaxy group, with NGC 1566 glittering and glowing, its bright nucleus framed by swirling and symmetrical lavender arms.^{53,54}

³This galaxy forms a member of the NGC 1566 subgroup of the Dorado Group, of which it is dominant and brightest member (although Kilborn and colleagues (2005) listed it as second brightest member of the NGC 1566 group after NGC 1553). The X-ray emission from the group is dominated by the hot gas halo of this galaxy, which extends out to 29 kpc before merging with the background radiation. The galaxy appears to be interacting with smaller members of its subgroup. Radio emissions suggest the disk is asymmetrical and the neutral hydrogen gas shows a mild warp.

The HAWK-I instrument mounted on the telescope's Nasmyth (side) port. HAWK-I is attached on Yepun, Unit Telescope number 4 of ESO's Very Large Telescope and saw FirstLight on the night of 31 July 2007. HAWK-I covers about 1/10th the area of the Full Moon in a single exposure. It is uniquely suited to the discovery and study of faint objects, such as distant galaxies or small stars and planets.



Figure 8 Hubble image shows NGC 1566, a beautiful galaxy located approximately 40 million light-years away in the constellation of Dorado (The Dolphinfish). NGC 1566 is an intermediate spiral galaxy, meaning that while it does not have a well-defined bar-shaped region of stars at its center like barred spirals it is not quite an unbarred spiral either.^{53,54}

Analysis

The Figure 1 show the image of Georgia, scale 1:200, on April 11, 2003, PM, and nucleus at the coordinates given in the image.

The Figure 2 show the image of Georgia, at scale 1:20, on April 11, 2003, AM, whose nucleus was at the approximate coordinates of those in the image.

The Figure 3 the image of Georgia, scale 1:20, on April 11, 2003, PM and nucleus at the coordinates given in the image. The Figure 4 show the image of Georgia, on a 1:100 scale, in an atmospheric pressure gradient model generated by the Zoom Earth system, on April 11, 2003, 12:30, with 951 mbar, and whose core was located at approximate coordinates of image.

The Figure 5 the image of Georgia, scale 1:100, in surface wind model generated by the Zoom Earth system, on April 11, 2003, 12:00, with 5km/h WSW, and nucleus at the coordinates given in the image.

The mathematical model⁴⁴ for the atmospheric pressure gradient in Figure 5, used by Zoom Earth⁴² matches the correct way to scale the atmospheric pressure, as can be seen in the comparison of the satellite images in Figures 1 and 2.

The model of wind currents Figure 4, for the displacement of air masses observed in the images is consistent with that observed in Figures 2-3, which presents a great turbulence in the vortex.

The highlighted cyclone vortex Figures 2 and 3 still in turbulent formation presents two linear containment barriers, in an L shape.

The subtropical cyclone that formed northwest of South Georgia & South Sandwich Island is here called Georgia. It moved 237 km in

12 h towards the West, when it was 589km from South Georgia Island, to 809 km from the center of the coast of the South Georgia Island. During this time interval, it maintained an atmospheric pressure at sea level at its vortex close to 951 hPa. It presented rotational winds of 5 km/h approximately 8 km from the central vortex.

With an approximate dimension of 1,000,000 km², and an area of direct influence of 3,500,000 km², the subtropical cyclone Georgia moved at an average speed of 19.75 km/h.

The mathematical model for the atmospheric pressure gradient in Figure 5, used by ZoomEarth⁴² matches the correct way to scale the atmospheric pressure, as can be seen in the comparison of the satellite images in Figures 1 and 2.

The model of wind currents Figure 4, for the displacement of air masses observed in the images is consistent with that observed in Figures 2 and 3, which presents a great turbulence in the vortex.

The highlighted cyclone vortex Figures 2 and 3 still in turbulent formation presents two linear containment barriers, in an L shape.




Figure 9 Image of Georgia, scale 1:100, in surface wind model generated by the Zoom Earth system, on April 11, 2003, 12:00, with 5km/h WSWV, and nucleus at the coordinates given in the image.

Figure 9 shows Georgia’s double spiral shape at 1:200 scale. The graph was constructed from satellite images, Figures 1-3, with cloud streams formed by wind currents in Georgia.

The analogous shape of Georgia and the galaxies Messier 83 and NGC 1566, studied here, is clear. These present a double spiral, as studied by Lindblad,¹ but with the Cote’s spiral form, Gobato et al.^{12,13,16}

Table 1 shows the coordinates of Georgia, between April 11, 2023, AM and PM, which Georgia maintains, a central vortex pressure of 951 mbar, with an approximate dimension of 5 km.

Conclusion

The subtropical cyclone that formed northwest of South Georgia & South Sandwich Island is here called Georgia. It moved 237 km in 12 h towards the West, when it was 589km from South Georgia Island, to 809 km from the center of the coast of the South Georgia Island. During this time interval, it maintained an atmospheric pressure at sea level at its vortex close to 951 hPa. It presented rotational winds of 5 km/h approximately 8 km from the central vortex.

Table 1 Subtropical Cyclone Georgia: Location/Pressure

April 11, 2023	Coordinates	Pressure (hPa)
AM	53° 13'09" S 27° 45'05" W	951
PM	53° 16'42" S 24° 00'38" W	951

With an approximate dimension of 1,000,000 km², and an area of direct influence of 3,500,000 km², the subtropical cyclone Georgia moved at an average speed of 19.75 km/h.

The highlighted cyclone vortex still in turbulent formation presents two linear containment barriers, in an L shape.

The have Georgia’s double spiral Cote’s shape. The analogous shape of Georgia and the galaxies Messier 83 and NGC 1566, studied here, is clear. These present a double spiral, as studied by Lindblad,¹ but with the Cote’s spiral form, Gobato et al.^{12,13,16}

References

- Lindblad B. On the circulation theory of spiral structure. *Astrophysica norvegica*. (12). Stockholms Observatorium, Saltsjobaden. 1964.
- Parker MC, Jeynes C. Maximum Entropy (Most Likely) Double Helical and Double Logarithmic Spiral Trajectories in Space–Time. *Scientific Reports*. 2019;10779.
- Cyclone. Creative Commons. CC BY–SA 3.0. 2023.
- American Meteorological Society. Glossary of Meteorology: Cyclone, 2020.
- Landsea, C. Subject: (A6) What is a sub–tropical cyclone? 2009.
- Atlantic Oceanographic and Meteorological Laboratory.
- Vasquez T. Weather Forecasting Handbook (5th Edition). Weather Graphics Technologies. 2002.
- Gobato R, Heidari A. Cyclone Bomb Hits Southern Brazil in 2020. *Journal of Atmospheric Science Research*. 2020;3(3):8–12.
- Armentrout D, Armentrout P. Tornadoes, Series: Earth’s Power. Rourke Publishing (FL), 2007, ISBN: 1600442331,9781600442339.
- Edwards R. The Online Tornado. Storm Prediction Center. National Oceanic and Atmospheric Administration. 2006.
- Gobato R, Mitra A, Valverde L. Tornadoes analysis Concordia, Santa Catarina, Southern Brazil, 2022 season. *Aeronautics and Aerospace Open Access Journal*. 2022;6:184–188.
- Gobato R, Mitra A, Gobato MRR, et al. Cote’s Double Spiral of Extra Tropical Cyclones. *Journal of Climatology & Weather Forecasting*. 2022;10:1–5.
- Gobato R, Mitra A, Heidari A, et al. Spiral galaxies and powerful extratropical cyclone in the Falklands Islands. *Physics & Astronomy International Journal*. 2022;6:48–51.
- de Leão M, Rodrigues P R. Methods of Differential Geometry in Analytical Mechanics, Series: Mathematics Studies. Elsevier Science Ltd. 1989.
- Gobato R, Heidari A, Mitra, A, et al. Spiral Galaxies and Powerful Extratropical Cyclone in the Falklands Islands.
- Gobato R, Heidari A, Mitra, A, et al. Extratropical Cyclone in the Falklands Islands and the Spiral Galaxies. *Sumerianz Journal of Scientific Research*. 2022;32–43.
- Gobato R, Mitra A, Heidari A, et al. Spiral galaxies and powerful extratropical cyclone. 2022.

18. Gobato R, Mitra A. Vortex Storms in the West of Santa Catarina. *Biomedicine and Chemical Sciences*. 2022;1:41–46.
19. Gobato R, Heidari A, Mitra A. Mathematics of the Extra– Tropical Cyclone Vortex in the Southern Atlantic Ocean. *Journal of Climatology & Weather Forecasting*. 2021;9:1–5.
20. Bluestein HB. Severe Convective Storms and Tornadoes: Observations and Dynamics, Series: Springer Praxis Books Springer– Verlag Berlin Heidelberg. 2013.
21. Gobato R, Gobato MRR, Heidari A. Evidence of Tornadoes Reaching the Countries of Rio Branco do Ivaí and Rosario de Ivaí, Southern Brazil on June 6, 2017. *Climatol Weather Forecasting*. 2018;6(4).
22. Gobato R, Gobato MRR, Heidari A. Evidence of Tornadoes Reaching the Countries of Rio Branco do Ivaí and Rosario de Ivaí, Southern Brazil on June 6, 2017. *Sci Lett*. 2019;7(1):32–40.
23. Gobato R, Gobato MRR, Heidari A. Storm Vortex in the Center of Paraná State on June 6, 2017: A Case Study. *Sumerianz Journal of Scientific Research*. 2019;2(2):24–31.
24. Gobato R, Gobato MRR, Heidari A. Vortex Cote’s Spiral in an Extratropical Cyclone in the Southern Coast of Brazil. *Archives in Biomedical Engineering and Biotechnology*. 2020;4(5):1– 4.
25. Gobato R, Heidari A. Vortex Cote’s Spiral in an Extratropical Cyclone in the Southern Coast of Brazil. *J Cur Tre Phy Res App*. 2020;1(2):109–112.
26. Gobato R, Heidari A, Mitra A, et al. Cotes’s Spiral Vortex in Extratropical Cyclone Bomb South Atlantic Oceans. *Aswan University Journal of Environmental Studies (AUJES)*. 2020;1(2):147–156.
27. Whittaker ET. A Treatise on the Analytical Dynamics of Particles and Rigid Bodies: With an Introduction to the Problem of Three Bodies. New York: Dover. 2022:83.
28. Vossle DL. Exploring Analytical Geometry with Mathematica. Academic Press. 1999.
29. Casey J. A treatise on the analytical geometry of the point, line, circle, and conic sections, containing an account of its most recent extensions, with numerous examples. University of Michigan Library. 2001.
30. Sharipov R A. Course of Analytical Geometry. Bashkir State University (Russian Federation). 2011.
31. Whittaker ET, McCrae Sir W. Treatise on analytical dynamics of particles and rigid bodies. Cambridge Mathematical Library, Cambridge University Press. 1989.
32. George JJ. Weather Forecasting for Aeronautics. Elsevier Inc. 1960.
33. Yorke S. Weather Forecasting Made Simple. Countryside Books Reference. Countryside Books. 2010.
34. Anderson JD. Fundamentals of Aerodynamics. McGraw–Hill Companies. 1984.
35. Weisstein EW. Cotes’s Spiral Cotes’s Spiral. Wolfram MathWorld. 2023.
36. Gobato R, Heidari A, Mitra A, et al. Cotes’s Spiral Vortex in Extratropical Cyclone bomb South Atlantic Oceans. 2020.
37. Fischer R. Fibonacci Applications and Strategies for Traders: Unveiling the Secret of the Logarithmic Spiral. 1993.
38. Bluestein HB, Bosart LF. Synoptical of Dynamic Meteorology and Weather Analysis and Forecasting: A Tribute to Fred Sanderson, Series: Meteorological Monographs 3(55). American Meteorological Society. 2008.
39. South Georgia and the South Sandwich Islands. Creative Com– mons. CC BY–SA 3.0. 2023.
40. Toomre A. Theories of Spiral Structure. *Annual Review of Astronomy and Astrophysics*. 1977;15(1):437–478.
41. Becker W, Contopoulos G. The Spiral Structure of Our Galaxy, Series: Inter– national Astronomical Union. Becker, w. Contopoulos G.(eds.); Union Astronomique Internationale 38, Springer Netherlands. 1970.
42. Zoom Earth. NOAA/NESDIS/STAR, GOES–East, RainView. zoom. earth. 2010.
43. Saffman PG. Vortex dynamics. Series: Cambridge monographs on mechanics and applied mathematics. Cambridge University Press. 1992.
44. Gobato R, Gobato A, Fedrigo DFG. Study of tornadoes that have reached the state of Parana. *Parana J Sci Educ*. 2016;2(1):1–27.
45. Brazil’s Navy. Synoptic Letters. Brazil’s navy. Synoptic Letters. Zoom Earth. 2023.
46. Rafferty JP. Storms, Violent Winds, and Earth’s Atmosphere. Series: Dynamic Earth. Britannica Educational Publishing. 2010.
47. Krasny R. A study of singularity formation in a vortex sheet by the point vortex approximation. *J Fluid Mech*. 1986;167:65–93.
48. Saffman PG. Vortex dynamics. Series: Cambridge monographs on mechanics and applied mathematics. Cambridge University Press. 1992.
49. Sokolovskiy MA, Verron J. Four–vortex motion in the two layer approximation – integrable case. RDX. 2000.
50. Gobato R, Heidari A. Vortex hits southern Brazil in 2020. *J Cur Tre Phy Res App*. 2020;1(2):109.
51. NASA gov. Messier 83 (The Southern Pinwheel). 2017.
52. Spiral galaxy. Creative Commons. CC BY–SA 3.0.
53. NASA gov. Messier 83 (The Southern Pinwheel). 2017.
54. ESA/Hubble & NASA. NGC 1566. European Space Agency. 2020.