

Does Temporary Catheter Drainage of Urine Improves Detrusor Function in Chronic Urinary Retention in Patients with Bladder Outlet Obstruction?

Abstract

Chronic urinary retention (CUR) is one of the consequences of bladder outlet obstruction (BOO) that occurs due to impaired functions of detrusor muscles secondary to obstruction related changes in the bladder wall. The effect of catheter drainage (CD) on detrusor function in CUR in animals is well studied but in humans there are very few studies to see its effect in CUR.

Material and methods: This is a non-randomized, retrospective study wherein case records of 50 eligible patients attending the Department of Urology during January 2010 to March 2012 were evaluated for CUR. Two UDS, performed 6-12 weeks (average 9 weeks) apart and the parameters which were evaluated were bladder sensation, capacity and voiding detrusor pressure. These parameters were also evaluated against the age of the patient, duration of symptoms and the amount of urine drained on catheterization.

Results: The commonest symptoms observed were decrease flows of urine, frequency, straining and incomplete emptying. Bladder function improvement after CD was observed to be maximum in patients with following characteristics: age at presentation in the range of > 60 to < 70 years, duration of lower urinary tract symptoms (LUTS) less than 5 years and amount of urine drained at the time of catheterization being < 500 ml.

Conclusion: CUR secondary to BOO is a common problem in elderly males. UDS is the "gold standard" for its evaluation and helps to plan further management. CD of urine is an effective mode of temporary management in patients of CUR secondary to BOO as it provides rest to the detrusor muscle in humans.

Keywords: Bladder outlet obstruction; Urinary bladder; Chronic urinary retention; Catheter drainage; Urodynamic study

Research Article

Volume 4 Issue 1 - 2017

Mahendra Pal*, Vinod S Joshi and Sharad N Sagade

Department of Urology, P D Hinduja National Hospital and Medical Research Center, India

***Corresponding author:** Mahendra Pal, Department of Urology, P D Hinduja National Hospital and Medical Research Center, Mahim, Mumbai, India, Tel: 400016 9757091924; Email: mahen1197@gmail.com

Received: August 24, 2016 | **Published:** January 18, 2017

Abbreviations: CUR: Chronic Urinary Retention; BOO: Bladder Outlet Obstruction; CD: Catheter Drainage; LUTS: Lower Urinary Tract Symptoms; PVR: Post Void Residual; DM: Detrusor Muscle; DP: Detrusor Pressure; UDS: Urodynamic Study; SPC: Suprapubic Catheter; LUT: Lower Urinary Tract

Introduction

Chronic urinary retention (CUR) secondary to bladder outlet obstruction (BOO) in spite of being a frequently encountered clinical problem, there are lacunae in the understanding of its patho mechanisms and management protocols in humans. International Continence Society defined CUR as "a non-painful bladder, which remains palpable or percussible after the patient has passed urine" [1]. However the practiced definition of CUR is "post void residual (PVR) urine of >300 ml in men who are voiding, or >1000 ml in men who are unable to void" [2,3].

The pathophysiology in CUR secondary to BOO revolves around the decrease contractile strength of the detrusor muscle

(DM) which causes incomplete evacuation of urine. In initial phases of BOO, DM undergoes hypertrophy with increase collagen deposition in the stroma. This helps to increase the detrusor pressure (DP) there by compensating for obstruction and maintaining the urinary flow [4-6]. In due course of time, changes in vascular supply and neural innervations of DM occur, which eventually results in reduced DM sensitivity and contractility and leads to detrusor weakening or failure resulting in CUR [7-9]. Classification of CUR has been attempted on various basis [10-15] (Table 1). The key investigation in CUR is the "Urodynamic study" (UDS), and is also the 'gold standard' for measuring DP and guide to plan further management [14]. Most treatment options are based on "expert opinion," however there are no established guidelines for the management of CUR secondary to BOO due less number of evidences [16-18] (Figure 1). This study attempts to evaluate the effect of temporary catheter drainage (CD) of urine on detrusor function in CUR with the help of UDS in men with BOO along with the factors that influence detrusor recovery.




Figure 1: Incidence of clinical symptoms.

Table 1: Classification of CUR.

	By Clinical Evaluation
1.	Inability of the patient to pass urine: Complete or Partial Symptoms: Painful or Painless Etiology/Mechanism: Obstructive or Non-obstructive
	Urodynamic Study
2.	A) High pressure chronic retention (HPCR): The bladder pressure is more than 30 cm H ₂ O during micturition. B) Low pressure chronic retention (LPCR): The bladder pressure is less than 20 cm H ₂ O during micturition.

Patients and Methods

This is a retrospective, non-randomized study. 50 patients reporting to the department of urology during January 2010 and March 2012 were evaluated for the lower urinary tract symptoms (LUTS), including the clinical history and urodynamic evaluation with appropriate inclusion and exclusion criteria (Table 2). The study was approved by the institutional review board. Forty eight patients presented with indwelling Foley’s catheter with an average duration of 3 weeks (range 2-4 weeks) during their first visit, two patients had suprapubic catheter (SPC) of an average duration of 2.5 weeks (range 2-3 weeks). The amount of urine drained on initial catheterization was recorded from previous documents. UDS was performed within 4 weeks after catheterization recorded as first UDS study. A second UDS was done approx 6-12 weeks after the first. Standard multichannel UDS was followed in all case.

Definitions and criteria used

- i. CUR: PVR of >350 ml in men on ultrasound or > 350 ml of

urine drained on catheterization.

- ii. Bladder compliance, sensation, capacity and DP were measured at volume the perception and strong desire occur to pass urine respectively.
- iii. Detrusor function improvement: Minimum of 15 cm H₂O improvement in DP.

Statistical analysis

Descriptive statistics were described as means, standard deviations and ranges. For categorical variables percentages were used. A value of p <0.05 was considered significant. All tests were performed using SPSS version 16 statistical package under the guidance of a bio-statistician.

Observations

Patients were grouped into 4 different categories of age group for analysis purpose. What are the 4 categories? (It is unclear to follow in the following sentences/Paragraph. Please list the

4 groups first before the details). The mean age of presentation was 71 years (52-90 years). Majority of the men were seen in the age group of 70-80 years constituting 44% (n=22) of the study population (Table 3). The commonest symptoms were decrease flow of urine, frequency, straining and incomplete emptying observed in 46 patients, thin stream was the least common presentation observed in 4 patients only, 4 patients were asymptomatic (Figure 1). Patients were divided into three categories of duration of symptoms and most of the patients (n=23, 46%) had symptoms since 0-5 years. The average duration of symptoms was 8.5 years, ranging from 2-15years (Figure 2). Patients were divided in to three categories of amount of urine drained <500 ml (n=12), 501-1000 ml (n=31), >1000 ml (n=7). Bladder function was assessed in all patients by UDS evaluation.

Bladder compliance was normal in all patients. Overall improvement in UB sensation was noticed in all age groups and was statistically significant (p <0.001). However, the improvement in < 60 years was not statistically significant in spite of reduction in mean volume of urine being present during 2nd UDS evaluation (Table 4). There was an improvement noted in the capacity of UB

during UDS and was statistically significant (p <0.001) except for age group < 60 years (Table 5). DP was assessed in different age groups in which significant improvement was observed during 2nd UDS evaluation. This improvement in DP after catheterization was statistically significant (P<0.001). However no statistically significant change was seen in the age group of <60 years (Table 6). Grades of voiding detrusor function in various age groups: Most of the patients (n=31) showed improvement in detrusor function. Maximum improvement was noticed in age group >60 <70 years (78.5%), (Table 7).

Correlation between duration of symptoms and grades of detrusor function: Although improvement in detrusor function was seen with lesser duration of symptoms, it was statistically insignificant (p = 0.124) (Table 8). Comparison of grades of detrusor function and amount of urine drained: Improvement in detrusor function was observed in 11 patients (91.6%) in whom the amount of drained urine was < 500ml. The amount of urine drained did not show statistically significant influence on detrusor function (Table 9).




Figure 2: Duration of symptoms.

Discussion

CUR secondary to BOO occurs in elderly and DM is significantly damaged [16]. Although being a common clinical entity seen by clinicians there is sparse evidence of its pathophysiology and management guidelines in humans (Figure 3). A total of 50 study patients were evaluated in our study and most of the patients i.e. 22 (44 %) belonged to the age group of >70 to <80 years, followed by 14 (28%) patients in > 60 to <70 years age group. This was consistent with the clinical findings, as physiological changes

in lower urinary tract (LUT) along with prostatic enlargement occurs with aging and it is not uncommon in elderly men to ignore their urinary symptoms due to misbelieve that these are part and parcel of normal changes of aging which leads to delay in seeking treatment [11-13]. Also, many patients with BOO initially prefer medical management. However, a substantial number of these patients have disappointing results and coupled with reluctance to avoid surgery. These patients eventually present as CUR at a later stage thus a advanced age group presentation with CUR is observed [13, 16-27].

Table 2: Inclusion and exclusion criteria.

Inclusion Criteria	
1.	Male sex Age > 40 years Subjects with CUR due to BOO Subjects with significant post void urine (more than 350 ml of post void urine) on ultrasound or urine drained on catheterization. Subjects with asymptomatic urinary tract infection.
Exclusion Criteria	
2.	Subjects with following ailments: Spinal cord injury Cerebro-vascular accident Neurological disease Long term uncontrolled diabetes mellitus Peripheral neuropathy. Long term (> 4 months) of continuous urinary drainage in form of: a) Per urethra catheter b) Suprapubic catheter (SPC) c) Clean self intermittent catheterization (CSIC) Subjects unsuitable for operative treatment (i.e. requiring assistance for their daily work) scheduled for long term catheterization of urinary drainage due to Immobilization. Symptomatic urinary tract infection.

Table 3: Age-wise distribution of patients.

Age Group (yrs)	No. of Patients	Percentage (%)
< 60	03	6
≥ 60 < 70	14	28
≥ 70 < 80	22	44
≥ 80	11	22
Total	50	100

Table 4: Comparison of Sensation on UDS evaluation in various age groups after temporary CD.

Age Group (yrs)	1 st UDS Evaluation		2 nd UDS Evaluation(6-12 weeks)		Paired t-test p-value
	Mean (ml.)	SD	Mean (ml.)	SD	
< 60	337.67	196.64	276.67	92.18	0.440 (NS)
≥ 60 - < 70	260.142	113.10	158.79	74.00	0.001 (S)
≥ 70 - < 80	321.23	157.21	203.36	121.64	0.000 (S)
≥ 80	321.82	143.11	217.64	124.64	0.023 (S)
Total	305.24	143.41	198.42	110.58	P<0.001

Table 5: Age-wise improvement in bladder capacity after temporary CD.

Age Group (yrs)	1 st UDS Evaluation		2 nd UDS Evaluation (6-12 weeks)		Paired t-test p-value
	Mean (ml.)	SD	Mean (ml.)	SD	
< 60	466.33	57.93	304.00	142.79	0.085 (NS)
≥ 60 - < 70	442.50	177.01	270.00	147.74	0.000 (S)
≥ 70 - < 80	513.87	191.50	320.59	115.30	0.000 (S)
≥ 80	526.20	258.64	360.10	246.00	0.015 (S)
Total	493.50	195.97	313.18	157.98	p<0.001 (S)

Table 6: Comparison of Detrusor Pressure after temporary CD.

Age Group (yrs)	1 st UDS Evaluation		2 nd UDS Evaluation (6-12 weeks)		Paired t-test p-value
	Mean (in cm H ₂ O)	SD	Mean (in cm H ₂ O)	SD	
< 60	28.33	5.77	53.00	18.36	0.190 (NS)
≥ 60 - < 70	18.57	15.77	55.79	34.06	0.001 (S)
≥ 70 - < 80	15.87	14.24	46.35	27.36	0.000 (S)
≥ 80	19.10	12.92	42.50	13.81	0.000 (S)
Total	18.02	14.07	48.62	26.81	P < 0.001 (S)

Table 7: Age wise improvement in detrusor function after temporary CD.

Age Groups (years)	No. of Patients	Grades of Detrusor Function Improvement		
		Improved (>40 cm H ₂ O) n (%)	Mild Improvement (20-40 cm H ₂ O) n (%)	No Improvement (<20 cm H ₂ O) n (%)
< 60	3	2 (66.6)	0	1(33.3)
>60--<70	14	11(78.5)	1(7.1)	2(14.2)
>70--<80	22	13(59)	7(31.81)	2(4.5)
>80	11	5(45.5)	6(54.5)	0
Total	50	31	14	5

Table 8: Correlation between duration of symptoms and improvement in detrusor function after temporary CD.

Duration of Symptoms (years)	No of Patients (n=50)	Grades of Detrusor Function Improvement			χ ² test
		Improved (>40 cm H ₂ O) n (%)	Mild Improvement (20-40 cm H ₂ O) n (%)	No Improvement (<20 cm H ₂ O) n (%)	
0-5	23	18(78.2)	3(13)	2(8.6)	χ ² = 7.24 df=4 p=0.124 (NS)
5-10	19	9(47.3)	8(42)	2(10.5)	
>10	4	1(25)	2(50)	1(25)	
Total	50	28	13	5	

Table 9: Correlation between urine drained and grades of detrusor function after temporary CD.

Amount of Urine Drained (ml)	No. of Patients n (%)	Grades of Detrusor Function Improvement			χ ² test
		Improved (>40 cm H ₂ O) n (%)	Mild Improvement (20-40 cm H ₂ O) n (%)	No improvement (<20 cm H ₂ O) n (%)	
<500	12(24)	11(91.6)	-	1(8.3)	χ ² = 7.07 df=4 p=0.132 (NS)
501-1000	31(62)	16(51.61)	12(38.7)	3(9.6)	
>1000	7(14)	4(57.1)	2(28.5)	1(14.2)	
Total	50	31	14	5	




Figure 3: Expected management of CRU.

LTC: Long Term Catheterization; CISC: Clean Intermittent Self Catheterization

Underlining the significance of aging in CUR in their review, Miegs JB et al. [13] and Ian K Walsh et al. [15] concluded that the age was an important risk factor for developing CUR. They found that the incidence of CUR increases with age from 0.4 /1000 person in age group 45-49 years to 7.9/1000 person in age 70-80 years and also that the risk of having CUR was 1/10 in 7th decade which increases to 1/3 in 8th decade. A review by Davies JA [17] stated that the average age of developing CUR secondary to BOO was 66 years. It was associated with co morbidities contributing to urinary problems. They observed that the average age at which patient develops detrusor failure (overflow incontinence) was 71 years which is consistent with the findings of the present study.

In our study 46 (92%) patients gave prior history of LUT symptoms, 4 (8%) patients were asymptomatic. Commonest symptoms observed were decrease flow of urine, frequency, straining, incomplete emptying in 46 patients followed by intermittency, nocturia, hesitancy, incontinence and thinning of stream. This is in agreement with the observations made by Davies JA [17] in their review that the common symptoms of which the patients are aware of are frequency, decrease flow and hesitancy. A retrospective study by SA Kaplan et al. [19] concluded that most common symptom of BOO was nocturia followed by increase frequency of urine, decrease flow of urine, hesitancy, dribbling, straining, sensation of incomplete voiding and incontinence.

All patients in our study had painless retention of urine probably due to impaired sensitivity of bladder and detrusor contractility due to bladder distension in CUR. Many studies demonstrated that in BOO there is recruitment of neural-detrusor responses causing altered neural control of micturition resulting in reduced bladder contractility, impaired central processing, and altered sensation [28-32]. Gosling et al. [9] in a prospective study showed that BOO leads to decrease in the autonomic innervations

of human DM which in turn results in decrease sensitivity of bladder.

Greenland Je et al. [33] and Kershen et al. [34] studied in pig and human respectively and concluded that blood flow in bladder reduces with distension that leads to detrusor ischemia and its impaired function. It has been shown in animal model that after relieving the obstruction, these changes are reversible as muscle mass returns to near normal, sensation and compliance improves and bladder capacity decreases. To regain the normal bladder function, it is necessary to give rest to the fatigued detrusor muscle. This is achieved by CD of UB through per urethra, SPC or operative intervention. Continuous urinary drainage through catheter bypasses the BOO, keeps bladder in collapse state and provides rest to DM [35-39]. UDS is the gold standard for measuring detrusor contractile dysfunction and quantitating the degree of BOO [14, 40-46].

Limited data is available on determining the effect of temporary catheter drainage in patients with CUR secondary to BOO using UDS. Siroky et al. [47] in their prospective study of elderly population by UDS demonstrated that aging is associated with a reduced bladder capacity, an increase in uninhibited contractions, decreased urinary flow rate, diminished urethral pressure profile, and increased PVR urine volume. In our study the patients underwent UDS evaluation for parameters like compliance, sensation, capacity and bladder pressure on two occasions separated by duration of 6-8 weeks. The results analysed after the 2nd UDS evaluation revealed overall improvement in sensation, capacity and bladder pressure which was statistically significant ($p < 0.05$) in all age groups except for age group of less than 60 years. This could probably be attributed to less number of patients in this age group in our study.

All patients in our study had normal compliance. The overall improvement in UDS parameters observed in our study was consistent with the findings of Hautmann et al. [35] prospective study on UDS evaluation in CUR which concluded that temporary catheter drainage of the bladder results in improved compliance, sensation, and maximum bladder capacity after low pressure drainage through SPC for average 13.6 weeks however compliance in our patient was normal on presentation and after temporary CD. The patients that benefited most on catheter drainage had following characteristics on UDS evaluation: Age group on presentation in the range of > 70 to < 80 years, duration of symptoms less than 5 years, and amount of urine drained on presentation being < 500 ml.

Patients in whom least improvement was observed were in the age group of < 60 years, duration of symptoms > 10 years and amount of urine drained on presentation was being > 1000 ml. Many studies reveal that the duration of symptoms has adverse effect on bladder function as the long term bladder distension leads to irreversible ultrastructural changes in detrusor muscle and bladder wall but at what time period these irreversible changes occur, it is not clearly defined. Kershen RT [34] and Ohnishi N et al. [37] postulated that long term distension of bladder leads to DM ischemia which leads to decrease bladder contractility and impaired bladder emptying causing CUR.

Gabella G et al. [38] observed that in animal long term retention leads to detrusor damage due to increase collagen content in bladder wall and impaired contractility of detrusor. All these studies directly or indirectly indicate that more the duration, more the chance of detrusor damaged in CUR due to BOO. Bob Djavan et al. [46] reviewed that patients with age more than 80 years, residual urine of more than 1000 ml and initial voiding detrusor pressure less than 28 cm H₂O on presentation was associated with less chances of detrusor recovery after relieving the BOO.

Selius BA et al. [12] stated in their prospective study on urinary retention in adults stated that the first step in management of CUR is to use indwelling catheter to decompress the bladder for up to one month which contributes to reversing the impaired detrusor function. Ghalayini IF et al. [36] in a prospective study emphasized the usefulness of clean intermittent self catheterization (CISC) in the recovery of bladder function in men with CUR. He kept the patient on CISC before TURP and found good improvement in bladder function as per UDS findings in patients with CISC. N Ohnishi et al. [37] and Gabella et al. [38] in their study on rabbits and rats respectively concluded that changes observed in later stages of BOO like reduced detrusor contractility, increased muscle mass, decrease compliance secondary to collagen deposition were reversible after relieving the obstruction. They observed that the hypertrophied muscle weight and mass became near normal six weeks after releasing the obstruction and that no degeneration of muscle cell or nerve endings was seen. This study documents the great plasticity of the musculature in the reduction of muscle mass after de-obstruction. Releasing the obstruction was equal to CD and same results can be reproduced by CD.

The major limitations of the current study were its retrospective nature, small sample size and shorter duration of follow up. A prospective study with a larger sample size and a control group is required to confirm the findings of this study. But nonetheless

as there are very few studies in humans which have evaluated the detrusor function after CD in CUR patients secondary to BOO and almost none having evaluated the factors influencing the detrusor function after CD, we attempt to fill these lacunae through the present study.

Conclusion

Temporary CD in CUR has significant beneficial effect on DM recovery in patients with BOO and should be part of initial management. Age of presentation, duration of BOO and amount of urine drained at presentation have clinically significant influence on detrusor recovery in patients undergoing CD thus the management protocol should be individualized to have a favourable outcome. UDS remains the gold standard for diagnosis and evaluation of DM recovery in CUR in BOO after temporary CD. Prospective studies are needed to validate expected management of CUR.

References

1. Abrams P, Cardozo L, Fall M, Griffiths D, Rosier P, et al. (2002) The standardization of terminology of lower urinary tract function: report from the Standardization Subcommittee of the International Continence Society. *Neurourol Urodyn* 21(2): 167-178.
2. Negro CL, Muir GH (2012) Chronic urinary retention in men: How we define it, and how does it affect treatment outcome. *BJU Int* 110(11): 1590-1594.
3. Abrams PH, Dunn M, George N (1978) Urodynamic Findings In Chronic Retention Of Urine And Their Relevance To Result Of Surgery. *Br Med J* 2(6147): 1258-1260.
4. Goslin JA, Dixon JS (1980) Structure of trabeculated detrusor smooth muscle in cases of prostatic hypertrophy. *Urol Int* 35(5): 3351-3372.
5. Macarak EJ, Howard PS (1999) The role of collagen in bladder filling. *Adv Exp Med Biol* 462: 215-223.
6. Levin RM, Monson FC, Haugeard N, Buttyan R, Hudson A, et al. (1995) Genetic and cellular characteristics of bladder outlet obstruction. *Urol Clin North Am* 22(2): 262-283.
7. Azadzo KM, Tarcan T, Siroky MB, Krane RJ (1999) Atherosclerosis induced chronic ischemia causes bladder fibrosis and non-compliance in the rabbit. *J Urol* 161(5): 1626-1635.
8. Torrens M, Morrison JFB, Dixon JS, Gosling JA (1987) Structure and innervation in the human. In the physiology of the lower urinary tract. Springer-Verlag P. 3.
9. Gosling J, Gilpin S, Dixon JS, Gilpin CJ (1986) Decrease in the autonomic innervation of human detrusor muscle in outflow obstruction. *J Urol* 136(2): 501-504.
10. Mitchell JP (1984) Management of chronic urinary retention. *Br Med J (Clin Res Ed)* 289(6444): 515-516.
11. Odunayo Kalejaiye, Mark J Speakman (2009) Management of acute and chronic retention in men. *European Urology Supplements* 8: 523-529.
12. Brian A. Selius, DO, Rajesh Subedi (2008) Urinary retention in adults: Diagnosis and initial management: *American Family Physician*. 77(5): 643-650.
13. Meigs JB, Barry MJ, Giovannucci E, Rimm EB, Stampfer MJ, et al. (1999) Incidence rates and risks factor for acute urinary retention: The health professional follow up study. *J Urol* 162(2): 376-382.

14. Nitti VW (2005) Pressure flow urodynamic studies: The gold standard for diagnosing bladder outlet obstruction. *Rev Urol* 7 Suppl 6: S14-21.
15. IK Walsh, AR Stone. The role of urodynamics in assessing symptomatic benign prostate hyperplasia. prostateconditions.org
16. Mirone V, Imbimbo C, Longo N, Fusco F (2007) The detrusor Muscle : An Innocent Victim Of Bladder Outlet Obstruction . *Eur Urol* 51(1): 57-66.
17. Roehrborn CG (2005) Acute urinary retention: Risk and management. *Rev Urol* 7 Suppl 4: S31-S41.
18. DAVIES JA (1954) Retention of urine, mode of presentation and its surgical value. *Br Med J* 1(4865): 796-798.
19. Kaplan SA, Bowers DL, Te AE, Olsson CA (1996) Differential diagnosis of prostatism: A 12 year retrospective analysis of symptoms, urodynamics and satisfaction with therapy. *J Urol* 155(4): 1305-1308.
20. Levy BJ, Wight TN (1990) Structural Changes in the Aging Submucosa: New morphologic criteria for the evaluation of the unstable human bladder. *J Urol* 144(4): 1044-1055.
21. Resnick NM, Yalla SV (1987) Detrusor hyperactivity with impaired contractile function. An unrecognized but common cause of incontinence in elderly patients. *JAMA* 257(22): 3076-3081.
22. Malone Lee J, Wahedna I (1993) Characterisation of detrusor contractile function in relation to old age. *Br J Urol* 72(6): 873-880.
23. Fry CH, Bayliss M, Young JS, Hussain M (2011) Influence of age and bladder dysfunction on the contractile properties of isolated human detrusor smooth muscle. *BJU Int* 108(2 Pt 2): E91-96.
24. Jeong SJ, Kim HJ, Lee YJ, Lee JK, Lee BK, et al. (2012) Prevalence and clinical features of detrusor underactivity among elderly with lower urinary tract symptoms: A comparison between men and women. *Korean J Urol* 53(5): 342-348.
25. Pfisterer MH, Griffiths DJ, Schaefer W, Resnick NM (2006) The effect of age on lower urinary tract function: A study in women. *J Am Geriatr Soc* 54(3): 405-412.
26. Zhao W, Aboushwareb T, Turner C, Mathis C, Bennett C, et al. (2010) Impaired bladder function in aging male rats. *J Urol* 184(1): 378-385.
27. Nordling J (2002) The aging bladder--A significant but underestimated role in the development of lower urinary tract symptoms. *Exp Gerontol* 37(8-9): 991-999.
28. Cortivo R, Pagano F, Passerini G, Abatangelo G, Castellani I (1981) Elastin and collagen in the normal and obstructed urinary bladder. *Br J Urol* 53(2): 134-137.
29. Macarak EJ, Ewalt D, Baskin L, Coplen D, Koo H, et al. (1995) The collagens and their urologic implications. *Dv Exp Med Biol* 385: 173-177.
30. Uvelius B, Mattiasson A (1984) Collagen content in the rat urinary bladder subjected to infravesical outflow obstruction. *J Urol* 132(3): 587-590.
31. Rubinstein M, Sampaio FJ, Costa WS (2007) Stereological study of collagen and elastic system in the detrusor muscle of bladders from controls and patients with infravesical obstruction. *Int Braz J Urol* 33(1): 33-39.
32. Levin RM, Haugaard N, O'Connor L, Buttyan R, Das A, et al. (2000) Obstructive response of human bladder to BPH vs. rabbit bladder response to partial outlet obstruction: A direct comparison. *Neurourol Urodyn* 19(5): 609-629.
33. Greenland JE, Brading AF (1996) Urinary bladder blood flow changes during the micturition cycle in a conscious pig model. *J Urol* 156(5): 1858-1861.
34. Kershen RT, Azadzo KM, Siroky MB (2002) Blood flow, pressure and compliance in the male human bladder. *J Urol* 168(1): 121-125.
35. Hamann MF, van der Horst C, Naumann CM, Wiederholt C, Seif C, et al. (2008) Functional results after temporary continuous drainage of the hypocontractile bladder. *Urologe* 47(8): 988-993.
36. Ghalayini If, Al Ghazo Ma, Pickard RS (2005) A prospective randomized trial comparing transurethral prostatic resection and clean intermittent self-catheterization in men with chronic urinary retention. *BJU Int* 96(1): 93-97.
37. Ohnishi N, Horan P, Levin SS, Levin RM (2000) Intermittent catheterization limits rabbit bladder dysfunction in response to partial outlet obstruction. *J Urol* 163(1): 292-295.
38. Gabella G, Uvelius B (1999) Reversal of muscle hypertrophy in the rat urinary bladder after removal of urethral obstruction. *Cell Tissue Res* 277(2): 333-339.
39. Thomas Aw, Cannon A, Bartlett E, Ellis-Jones J, Abrams P (2004) The Natural History Of Lower Urinary Tract Dysfunction In Men: The Influence Of Detrusor Underactivity On The Outcome After Transurethral Resection Of The Prostate With A Minimum 10-Year Urodynamic Follow-Up. *Bju Int* 93(6): 745-750.
40. Toppercer A, Tetreault JP (1979) Compliance of the bladder: An attempt to establish normal values. *Urology* 14(2): 204-205.
41. Mc Kertich K (2011) Urodynamics. *Aust Fam Physician* 40(6): 389-391.
42. Homma Y, Batista J, Bauer S. Urodynamics Committee. 7: 317-372.
43. Schäfer W, Abrams P, Liao L, Mattiasson A, Pesce F, et al. (2002) Good urodynamic practices: Uroflowmetry, filling cystometry, and pressure-flow studies. *Neurourol Urodyn* 21(3): 261-274.
44. Chancellor MB, Blaivas JG, Kaplan SA, Axelrod S (1991) Bladder outlet obstruction versus impaired detrusor contractility: The role of outflow. *J Urol* 145(4): 810-812.
45. Abrams PH, Griffiths DJ (1979) The assessment of prostatic obstruction from urodynamic measurement and from residual urine. *Brit J Urol* 51(2): 129-134.
46. Djavan B, Madersbacher S, Klingler C, Marberger M (1997) Urodynamic assessment of patients with acute urinary retention: Is treatment failure after prostatectomy predictable?. *J Urol* 158(5): 1829-1833.
47. Siroky MB (2004) The aging bladder. *Rev Urol* 6(Suppl 1): S3-7.